



COVID-19 (Coronavirus)

TABLE OF CONTENTS

| | |
|---|----|
| 1. Summary of Updates | 3 |
| 2. Strength of Evidence Ratings | 4 |
| 3. Introduction | 5 |
| 3.1. Virus Characteristics | 7 |
| 3.2. Clinical Presentation | 9 |
| 3.3. Business Considerations | 12 |
| 3.4. Disability and Return-to-Work Considerations | 23 |
| 4. Vaccines | 24 |
| 5. Masks and Respirators | 30 |
| 6. Lockdowns and Shutdowns | 34 |
| 7. Diagnostic Approach | 34 |
| 8. Treatment Recommendations | 40 |
| 8.1. Overview | 40 |
| 8.2. Hydroxychloroquine or Chloroquine | 43 |
| 8.3. Azithromycin | 51 |
| 8.4. Favipiravir | 54 |
| 8.5. Lopinavir/Ritonavir | 56 |
| 8.6. Remdesivir | 59 |
| 8.7. Low-Molecular-Weight Heparin | 62 |
| 8.8. Fluvoxamine | 65 |
| 8.9. Paxlovid | 66 |
| 8.10. Molnupiravir | 68 |
| 8.11. IL-6 Receptor Antagonists | 69 |
| 8.12. Adalimumab | 71 |
| 8.13. Baricitinib | 71 |
| 8.14. Casirivimab plus Imdevimab (Regeneron) | 73 |
| 8.15. Bamlanivimab plus Etesevimab | 74 |
| 8.16. Sotrovimab | 78 |
| 8.17. Bebtelovimab | 81 |

| | |
|---|-----|
| 8.18. Ivermectin | 82 |
| 8.19. Convalescent COVID-19 Antibodies..... | 89 |
| 8.20. Glucocorticosteroids | 90 |
| 8.21. Interferon Beta-1b..... | 92 |
| 8.22. Ribavirin..... | 95 |
| 8.23. Zinc | 98 |
| 8.24. Vitamin D..... | 100 |
| 9. Rehabilitation..... | 102 |
| 9.2. Pulmonary Rehabilitation..... | 102 |
| 9.3. Cardiac Rehabilitation | 104 |
| 9.4. Exercise Therapy..... | 106 |
| 9.5. Memory and Cognition..... | 108 |
| 9.6. Joint Pain | 110 |
| 9.7. Mental Health..... | 110 |
| 10. Contributors | 111 |
| References | 117 |

1. SUMMARY OF UPDATES

Note: This guideline and its recommendations were last reviewed and updated on **April 29, 2022**. (prior versions: April 8, 2020; April 24, 2020; May 8, 2020; June 12, 2020; June 17, 2020; August 19, 2020; December 14, 2020; and March 29, 2021).

The total depth and breadth of quality literature for the treatment of COVID-19, although growing, still remains fairly limited, especially on certain important sections of guidance. Further, caution is warranted for the potential for changes in efficacy that may be present for current virus variants compared with the variant(s) studied in the existing publications. Some of the studies may continue to be particularly fluid due to the continuing pace of change in knowledge. Research data, especially those associated with treatments, continue to be published prior to peer review. Some newer treatments do not yet have published peer review papers; thus, reliance for those is necessarily on press releases, FDA documents, and other non-peer-reviewed sources. Under normal circumstances, such data would not be considered for an evidence-based guideline. However, the ongoing severity, urgency, and mortality associated with this pandemic do not allow the luxury of time to await the publication of randomized controlled trial data. The literature will continue to be monitored and this guideline will be updated as needed in response to new research reports, changes in prior reports caused by peer review, and any retractions of papers.

With this ninth version of this guideline, the document and updated guidance are transitioning to management of an endemic virus from an epidemic/pandemic, as the omicron variant was highly adept at worldwide dissemination in a matter of a few weeks.

The **April 29, 2022** update includes the following major changes:

- Transitioning of the guidance from managing an epidemic/pandemic to managing SARS-CoV-2 as an endemic virus
- Recommendations to promptly perform after-action reviews at the federal, state, and local levels. These reviews should define (in)effective responses to the pandemic and help to prepare for the next viral surge and/or epidemic/pandemic.
- Recommendation to develop multi-arm randomized trials of therapeutics for the early (24- to 48-hour) timeframe with existing evidence of potential efficacy to be implemented at the beginnings of the next variant surge
- Emphasis on management of the virus as being spread by aerosols
- De-emphasis on contact spread and recommendations against regular cleaning of surfaces purely for the purpose of control of the pandemic
- Recommendations against lockdowns
- Recommendations for selective use of N95/KN95 respirators for significantly immunosuppressed population
- Recommendation to discontinue masking in the general population
- Guidance for schools to remain open and without masking
- Recognition of the durability of natural immunity, while supporting vaccination

- New recommendations in support of fluvoxamine, ivermectin, paxlovid, molnupiravir, bebtelovimab, and sotrovimab
- Recommendations against convalescent antibodies, and casirivimab plus imdevimab (Regeneron)

2. STRENGTH OF EVIDENCE RATINGS

Strength of Evidence ratings are used to designate the quality and amount of evidence that supports a specific guideline recommendation, when taking into account the entire body of relevant evidence found in the literature search. The body of evidence on a topic consists of all studies found that were relevant to the specific clinical question and of acceptable quality. In general, the highest quality of evidence found should be used by the Panel as the basis for the guideline recommendation, unless other factors, such as the potential for harm, are an overriding consideration. When multiple studies of similar quality and relevance are found on a topic, these studies should be evaluated as a group; if results are generally consistent, they would be considered either Strong Evidence (for high-quality studies) or Moderate Evidence (for moderate-quality studies). In all cases, the rationale for each recommendation and scientific studies used as evidence, should be documented by the Panel.

- **A Strong evidence-base:** Two or more high-quality studies.¹
- **B Moderate evidence-base:** At least one high-quality study or multiple moderate-quality studies² relevant to the topic and the working population.
- **C Limited evidence-base:** At least one study of moderate quality.
- **I Insufficient Evidence:** Evidence is insufficient or irreconcilable.

For treatment, the criteria used by evidence reviewers to categorize the quality of individual randomized controlled trials as high, moderate, or low quality are: adequate randomization, concealed treatment allocation, baseline cohort comparability, patient blinded, provider blinded, assessor blinded, controlled for co-interventions, compliance acceptable, dropout rate acceptable, timing of assessments equivalent, data analyzed by intention to treat, and lack of bias.³ Each criterion receives a score of 0, 0.5, or 1. See [Table B in the Methodology](#) for a definition of each criterion and scoring level. Studies are considered of low quality if they are rated 3.5 or less, moderate quality if they are rated 4-7.5, and high quality if they are rated 8-11.

Please see our full [methodology](#) for more information.

¹For therapy and prevention, randomized controlled trials (RCTs) with narrow confidence intervals and minimal heterogeneity. For diagnosis and screening, cross-sectional studies using independent gold standards. For prognosis, etiology or harms, prospective cohort studies with minimal heterogeneity.

²For therapy and prevention, a well-conducted review of cohort studies. For prognosis, etiology or harms, a well-conducted review of retrospective cohort studies or untreated control arms of RCTs.

³ Van Tulder et al., 2003

3. INTRODUCTION

Novel coronavirus 2019 (COVID-19) is an acute respiratory infection caused by the coronavirus SARS-CoV-2. The disease it causes has been named “coronavirus disease 2019” (abbreviated “COVID-19”) (309).

The pandemic began in Wuhan, China in November 2019, then expanded markedly throughout the Wuhan region. The importance of the source is critical to prevent and/or for early identification of the next pandemic. Based on prior research and experience with coronavirus infections, the origin of this pandemic was initially suggested as having an animal host reservoir such as pangolins with origination from horseshoe bats in caves in Yunnan province, in southwestern China, which are located approximately 800 miles from Wuhan. However, an intermediate animal species (e.g., pangolins) has not been found (310) (311). The genetic sequencing of the virus uniquely includes furin cleavage sites typical of bioengineered viruses, and the Wuhan laboratory was conducting research with coronaviruses.

There is indirect and strongly disputed evidence suggesting that the epidemic may have begun in Wuhan earlier than November, including increased hospital traffic, web searches for potential COVID-related symptoms beginning in August 2019, and other information that suggested a potential virus laboratory shutdown in October 2019 (312) (313) (314) (315) (316). Yet, the source of the virus remains highly disputed (317) (318), with conflicting evidence surrounding gain-of-function experiments and viral alterations(319), and early publications denouncing the laboratory leak theory (318) were subsequently retracted (320). Regardless of the source, the Chinese New Year and lack of early travel cessation to/from Wuhan likely accelerated the spread of the virus through global travel hastening the development of a pandemic. COVID-19’s SARS-CoV-2 virus is now found on all continents and nearly all countries studied.

Quarantines were implemented early in the pandemic. However, they were likely ineffective at slowing or preventing the pandemic (321) for several reasons, including delays in implementing local, regional and global quarantining; late enactment of travel bans; early lack of rapid and accurate viral testing; early emphasis on contact instead of respiratory precautions; subsequent emphases on droplet spread with neglect of the potential for aerosol spread; ongoing delays in the institution of, and lack of attention to, aerosol precautions; large numbers of undiagnosed, mild, or asymptomatic patients spreading the virus a (322)(323); and inadequate detection of the spread of cases in regions prior to the recognition of COVID-19 within that area (324). An added fact later identified that precludes eliminating the virus from the global human population is the susceptibility of animals, although the importance of this potential factor regarding spread to humans is still poorly documented. Evidence of infections in many species of animals, presumably documenting the potential for 'animal reservoirs' (325) (326)(327,328,329), also precludes eliminating the virus from the global human population, without consideration of the other required criteria for viral eradication (e.g., no asymptomatic cases; no breakthrough cases among the vaccinated; ability to geographically restrict the virus; an effective method to interrupt disease transmission).

Public health management of this pandemic has varied markedly across countries, states, and jurisdictions. Because there was no quality evidence to support any of these measures early in the pandemic, expert opinion was relied upon for enactment of these measures. The initial guidance focused on handwashing and restricting travel to China (January-February 2020, which subsequently expanded to include other countries), varying degrees of closure for businesses and schools (March 2020), recommendations for personal masking (March-April 2020), and public masking orders (April-July 2020). Some states began to reopen most, if not all businesses, starting in May-June 2020. Subsequently, restrictions have waxed and waned largely in response to rising and falling COVID-19 incidence rates. Typically, a combination of approaches has been used, including the quarantine of affected individuals, contact tracing, isolation, stay-at-home orders, physical distancing, mask use, and the closure of non-essential businesses (66).

The pandemic subsided markedly in the summer of 2020 in northern latitudes. However, as fall/winter 2020–21 began, the pandemic surged in northern, cold climates where conditions of lower temperatures, lower humidity, less intense ultraviolet (UV) irradiation, and higher indoor population densities combined to likely contribute to record levels of cases in nearly all jurisdictions (330) (331) (332). Additionally, controversy regarding the efficacy and sustainability of various public health measures, especially the (re)closure of businesses and schools, has intensified as the case rates plummeted in 2021. Quality data are weak; some countries (e.g., Japan, South Korea) have instituted less stringent measures with seemingly somewhat comparable or better results (333) (334) (335) (336,337,338) (339) (340) (341) (342) (343). A metanalysis found lack of efficacy of lockdowns (5).

However, beginning in approximately October 2020, the delta variant (B.1.617.2) began to circulate in India (344). This variant was estimated to be approximately twice as contagious as the alpha variant (60), which was twice as contagious as the original virus. The delta variant was found in the United States in spring 2021 and subsequently was found in nearly all countries. While the original delta variant arose in a largely unvaccinated population, it has caused breakthrough infections among the vaccinated in the United States and elsewhere. The relatively low levels of viral spread in spring 2021 were replaced by surges by mid-summer that peaked in the United States in late August to early September.

Still, while there have been breakthrough fatalities, particularly among the immunocompromised (345) (346), most of the delta variant cases were among the non-vaccinated, as the vaccines still proved to be highly effective in preventing hospitalizations (346) (347) (348)(349,350,351) (352)(353). There were few moves to reinstate restrictions in the United States; however, Australia and some other countries re-instituted severe restrictions with no good evidence of commensurate benefits. Subsequently, there were at least two more variants of concern that were identified: lambda (which was identified in Peru in December 2020) and mu (which was identified in Columbia in January 2021) (344). However, these variants were eventually displaced by omicron.

In mid-November 2021, the omicron variant (B.1.1.529) was described in Botswana and South Africa (354), with the identification retrospectively from a sample from recently-arrived diplomats November 11, 2021 in Botswana (355). However, other samples in multiple countries (e.g., England, South Africa, Nigeria, USA) date the detection of omicron to November 1, 2021, thus assuring it had arisen weeks to months earlier although undetected. The origin of this markedly different variant with large numbers of mutations (approximately 4 times the spike protein mutations compared with the original strain; 3 times compared with the delta variant) is unclear, whether arising from a long-term host, going undetected over a series of mutations, arising from an animal reservoir, or originating from other source (356)(357). Omicron has been estimated to be more than twice as contagious as delta. It rapidly spread around the world, including to the United States by early December, being isolated from a person without a travel history on December 2, 2021. Omicron was so efficient at spreading that the peak case rate in South Africa occurred on approximately December 15, 2021. It displaced delta as the predominant variant in the United States in less than one month, by December 25, 2021 (358). The speed of spread has also effectively precluded proactive updating of vaccinations using existing techniques. The peak U.S. omicron-related COVID-19 case rate occurred on approximately January 14, 2022.

Worldwide, the pandemic continues to provide numerous challenges, especially in regards to the increasingly rapid rates of successive variant transmission, and among countries with lagging immunization rates or that relied on less effective vaccines. Challenges include surges, hotspot outbreaks, attempts at surge prevention, and mitigation; COVID-19 diagnostic testing availability, accuracy, capacities, and limitations; unique treatment challenges and sparse to varying evidence of efficacy; increasingly global public restlessness with restrictions; and increasing business/economic concerns. The economic damages include loss of employment, income, and housing; closure of in-

person school instruction; as well as worsening supply chain disruptions involving quite diverse sectors and products (359) (360) (361) (362).

Successes in reducing impacts of the first phases of the COVID-19 pandemic are primarily credited to rapid vaccination uptake, residual immunity due to prior infections, and initially falling infection rates. However, "termination" of the COVID-19 pandemic is being redefined, as we have come to realize that eradication is impossible. Omicron's substantially lower severity, including an ~91% lower risk of mortality (363), is facilitating the transition from pandemic to endemic management. Attention is now turning to issues such as vaccination breakthrough cases, the frequency of booster immunizations, the duration of vaccine-related immunity, the relative importance of vaccine-related compared with natural immunity, therapeutics, and subsequent viral variants.

3.1. VIRUS CHARACTERISTICS

Contagiousness

COVID-19's SARS-CoV-2 virus appears to be more contagious than the prior coronaviruses, and each successive widespread variant has been ever more successful at spreading (364). The Omicron variant has been estimated to be 2 to 3.3 times as infective as Delta and 10 times more infective than the original virus (365) (366).

Initially, the virus was thought to be primarily spread through direct contact, resulting in a primary national focus on handwashing (367) (368) (369). Then, the theory changed to droplets as the primary spread. These beliefs starkly changed and the science has now coalesced around this guideline's prior position early in 2020 that the virus is primarily spread by microdroplets/aerosols (defined as $<0.5\ \mu\text{m}$) followed by respiratory droplets (defined as $>100\ \mu\text{m}$ in size) (370) (371) (372,373,374,375,376,377,378,379,380) (381) (382,383,384,385,386,387,388,389,390)(391)(392).

Aerosols can remain suspended in the air potentially for hours (370), and well beyond the 6-feet (or 1-meter, per the World Health Organization) physical distancing guideline, with one estimate for need of up to 10-meter distances (390) (393)[51]. One experimental study estimated aerosolized virions retained infectivity and virion integrity for up to 16 hours (394), thus likely explaining the widespread and reproduced patterns of viral epidemic spread despite institution of physical measures. Still, to what extent an infectious dose can be generated, what is an infective dose, and what is present beyond 6-foot distances has yet to be clearly demonstrated (395,396,397,398,399,400,387). There is some evidence of a dose-response relationship between dose and clinical severity (401). Regardless, the recognition of aerosol spread has significant impacts on selecting the best preventive measures.

The contagiousness and virulence of the SARS-CoV-2 virus delta variant appear to be about 6-fold greater than that of influenza. Estimates of the contagiousness or transmission rate without interventions (e.g., physical distancing) of the original virus ranged from 2.0 to 3.9—that is, 2 to 3.9 new cases arise from each known case (i.e., R_0) (402), which is far higher than typical influenza transmission rate of ~1.3 (403). The R_0 for delta has been estimated at 5 and the R_0 for omicron is estimated at over 7. However, the reproduction number varies markedly and as the epidemic curve surges or tails off, the reproduction number rises or falls respectively. From a population standpoint, however, each case does not appear to be equally infectious. One analysis of 1,038 confirmed SARS CoV-2 infections in Hong Kong between January and April 2020 revealed that 80% of the infections were caused by just 19% of the initial cases; the majority of patients failed to infect anyone else. Most transmission occurred from household contacts, followed closely by external social events (404). Superspreading has continued to be a purported major mechanism of spread with successive variants (405) (406) (370) (407).

The asymptomatic infection rate of omicron in South African healthcare workers has been reported at 16%, which is approximately 6-fold higher than with the beta or delta variants (408). This striking asymptomatic carriage rate is theorized to help contribute to the rapid dissemination of omicron. Other estimates have ranged as high as 70-90% of cases worldwide being asymptomatic (409) (410). Evidence found prior variants also had evidence of high rates of asymptomatic cases, (411,412) (413,414,415,416,417,418,419,420,421,422,423,424,425,426,427,428,429,430,431,432,433); however, omicron has exceeded those rates.

The virus's survivability on surfaces varies depending on the material; it has been estimated with experimental methods to survive up to 9 days (434), although those experimental methods are limited by not including environmental settling rates, inactivation by UV light, or diffusion, and use of doses unlikely to be found under normal non-experimental conditions. Furthermore, a thin nanofilm of liquid from droplets has been reported to extend the viral survival on surfaces [91]. The total viable viral counts decline with time (390). The survival time of the virus was reported to differ by surface type, with approximate upper limits of detection being 4 hours on copper, 24 hours on cardboard, 48 hours on stainless steel, and 72 hours on plastic (434). Survival on human skin has been measured at 9.04 hours, which is much longer than the measured survival of influenza virus on skin (1.82 hours) (435). Survival of the virus in aerosols is thought to be as long as 16 hours (394). However, it is still unclear how much virus is needed to infect a human from either surfaces or aerosols. Many studies show detection of viral RNA that is likely inadequate for and/or incapable of transmitting an infection.

Preliminary experimental and epidemiological-ecological data suggest spread may be optimal in indoor and/or cooler climate conditions (330) (436,437,438,331), and prior data on the SARS coronavirus are corroborative (439). Experimental evidence suggests that simulated sunlight rapidly inactivates the virus. At a simulated sunlight intensity of the summer solstice at 40 degrees of latitude, the inactivation rate was 90% inactivated every 6.8 minutes (440). The ecological data indicate that there were slower rates of infection with higher temperatures in Delhi, India, and Pakistan (330) (331), although there was no correlation with humidity (330). The data from Pakistan also suggest an inverse relationship between COVID infection rates and UV light, although the UV data appear to be highly correlated with the heat indices (331). Other data suggest lower infections with higher humidity (332) (441). This suggests highly variable disease transmission risks based on seasonality and in indoor compared with outdoor environments.

Incubation and period of infectious viral shedding

The incubation period is the amount of time that occurs between exposure and the onset of symptoms. The incubation period of the SARS-CoV-2 virus was initially estimated to be approximately 5–6 days (323,442,443), while the estimate for delta was 4 days, and the estimates for omicron are 3 days (444). Previously, 97.5% of cases occurred by 11.5 days after exposure and infrequent cases of up to 14 days (445,371,446). There are some recent reports suggesting infectivity is still possible at 10 days (447) (448). Viral shedding may antedate symptoms by 1–2 days, peak viral load occurs 1-3 days before symptoms, (449) and viral titers are highest in the earliest phases of infection; however, a study has suggested the peak shedding for omicron infections may be 3-6 days after the onset of symptoms (450), although this study measured peak RNA material, which may not correlate directly with maximum viral replication (450).

The duration of infectious viral shedding remains controversial, primarily due to the ability to measure virus and/or virus particles in body fluids for long periods after the acute infection with sensitive techniques, such as polymerase chain reaction (PCR) (451,452). Yet, it is less clear whether these particles are infectious, and there are far fewer studies of viral shedding that relied on viral culture suggesting active virus. Even those few studies with viral culture results may not yield enough virus particles that are sufficient to provide an infectious dose (452).

A pooled study prior to omicron of 79 studies with 1,858 patients reported that pharyngeal virus shedding peaks prior to the onset of symptoms, averages 17 days, and lasts up to 83 days (453,454). The mean durations of viral shedding were 14 days in the lower respiratory tract, 16 days in stool, and 16 days in serum. Although replication-competent virus has not been isolated 3 weeks after symptom onset, recovered patients can continue to have SARS-CoV-2 RNA detected in their upper respiratory specimens for up to 12 weeks (455,456,457). Further study of 285 “persistently positive” persons, which included 126 persons who had developed recurrent symptoms, found no secondary infections among 790 contacts attributable to contact with these case patients. Efforts to isolate replication-competent virus from 108 of these case patients were unsuccessful, suggesting a lack of viable virus (455). No study detected live virus beyond the ninth day (453). These findings contrast with those of MERS and SARS, which peaked after symptom onset and lasted for shorter durations. It is also apparent that a positive PCR test is not a binary (yes or no) answer; the threshold cycle count should be reported, as counts above 28-30 are more likely to be false positive or have residual particles more likely to be incapable of transmitting infection.

There have been many reports of re-infections (458,459,460,461,462,463), which accelerated markedly with omicron. The risk of re-infection with omicron has been estimated at 5.4 times that of delta (464). Previously, a prior infection was estimated to have provided 85% protection against reinfection, but this estimate has now fallen to 19% (464). Yet, the cases are milder on average.

Due to the extent of re-infections and especially the problems with omicron, the concept of herd immunity is undergoing reevaluation. It is unclear if it will be resurrected for COVID-19.

3.2. CLINICAL PRESENTATION

There are at least six distinct types or clinical presentations of COVID-19’s SARS-CoV-2 virus infections, the first and third of which incur no healthcare visits; pre-symptomatic individuals may or may not incur healthcare visits (26):

- asymptomatic
- pre-symptomatic
- mild, subclinical infection (e.g., mild rhinorrhea, pharyngitis)
- upper respiratory tract infection (URI), which also may include gastrointestinal symptoms
- lower respiratory tract infection, including pneumonia
- acute respiratory distress syndrome (ARDS)

Treatments differ for each presentation (see [Treatment](#) section for more details).(465,466,467,468,469,470)(471)

Symptoms and Signs

The symptoms of COVID-19 vary but are generally typical of respiratory infections, such as fever and cough. The symptoms with the omicron variant are primarily rhinorrhea, headache, fatigue, sneezing, and sore throat (472); these symptoms differ compared with prior variants. COVID-19 symptoms may include the following, listed here for prior variants and omicron (407) (371) (473,474,475,476):

- fever, low or high grade (80–88%; omicron: 54%)
- dry cough (63–69%; omicron: 83%) (445,477)

- loss of appetite (39–84%; omicron: 33%) (478)
- fatigue (38–46%; omicron: 74%)
- sputum production (33–42%)
- chest pain or pressure (28–36%)
- dyspnea (shortness of breath) (19–35%)
- myalgia and/or arthralgia (muscle and joint pain; 15–33%; omicron: 58%)
- sore throat (12–14%; omicron: 72%)
- headache (11–15%; omicron: 68%)
- chills (6–11%)
- nausea or vomiting (5–10%)
- diarrhea (4–29%) [124]
- nasal congestion (4–5%)
- abdominal pain (4%; omicron: 6%)
- conjunctivitis (pink eye; 1%) (479)
- hemoptysis (1%)
- rhinorrhea (runny nose; omicron: 78%)
- sneezing (omicron: 43%)
- anosmia and dysgeusia (loss of smell and taste; 85% moderate/severe or anosmic; omicron: 12–23%) (480)

Severity of disease may be related to the inoculation dose (481). The wearing of masks has been theorized to increase the proportion of asymptomatic cases by lowering that inoculation dose (481,482).

Note: The following discussion of symptoms is based on reports of prior variants. It is premature to incorporate findings from omicron reports, although there is a dearth of reports of severe outcomes, unlike with prior variants.

Cardiovascular symptoms and signs were sometimes noted on initial presentation with prior variants (465,466,467,468,469,470). Immunothrombotic dysregulation associated with COVID-19 pneumonia has been described (483). Coagulopathy associated with antiphospholipid antibodies and multiple infarcts have been reported (484,485). Seizures have been reported as a presenting disorder (486). Young and old patients have presented with large-vessel strokes as an initial manifestation of COVID-19 infection (486,487). Pulmonary arterial thromboses have been reported (488). Among ICU patients, 31–59% of patients incurred venous or arterial thromboembolic event(s) (489,490), compared with 10–25% of patients hospitalized for other reasons (490,491). Heparin-induced thrombocytopenia (HIT) is a severe adverse reaction to heparin caused by heparin-dependent, platelet-activating anti-platelet factor 4 (PF4)/heparin antibodies frequently found in acute severe

COVID-19 infections and is associated with thrombotic events (492). Anti-phospholipid antibodies associated with thrombosis have also been identified in severe COVID-19 pneumonias (492). Recovering competitive athletes also have been found to have cardiac abnormalities on magnetic resonance imaging (MRI) (471).

Dermatological abnormalities such as urticaria, vasculitides, and pityriasis rosea have been described (493,494,495,496). The most common dermatological presentations have been polymorphic and erythematous, chilblain-like, and urticarial lesions (497). Various neurological and psychiatric presentations including stroke-like symptoms, altered mental status, dementia-like syndromes, and new or recurrent affective disorders have been reported (498,499,500,501,502,503,504,505)(506)(299)(298). Although the prevalence of direct kidney involvement in COVID-19 disease ranges from 3 to 15%, it is a marker for multiple organ failure and severe disease (507). Acute kidney injury is thought to be triggered by a cytokine storm. In addition, the ACE2 receptor, essential for viral uptake, is highly expressed on podocytes and tubule epithelial cells. Albuminuria and hematuria have been detected in COVID-19 infection (508), along with the isolation of viral RNA from urine (509). Most (71%) of those who die from COVID-19 have findings consistent with disseminated intravascular coagulation (510).

Because the symptoms for most patients are typical of nonspecific respiratory tract infections, they can be difficult to distinguish from other diseases (77,78). The disease commonly begins with mild symptoms for several days, which may readily facilitate its spread to other individuals. A minority of patients then develop more severe symptoms and may require ICU care (511). This appears to be most common at days 4–7 after symptom onset. These more severe cases of COVID-19 involve additional symptoms that typically accompany pneumonia, such as shortness of breath. Respiratory problems may further progress to severe dyspnea, require oxygen supplementation, and develop into acute respiratory distress syndrome (ARDS). Patients with pneumonia may have tissue hypoxia, tachypnea, tachycardia, and crackles on chest examination. Severe cases may present with shock and respiratory failure. The hallmarks of COVID-19 infection on thoracic imaging have been bilateral and peripheral ground-glass and consolidative pulmonary opacities (512).

The virus infection may also cause no symptoms; however, asymptomatic and pre-symptomatic individuals may still pass the virus to others, who may then develop symptoms (513,511,514). The CDC estimates that 40–45% of transmission occurs prior to symptom onset and that the infectiousness is comparable between asymptomatic and symptomatic individuals (513,323). Children tend to be asymptomatic or have milder symptoms, which suggests a mechanism that may accelerate disease transmission throughout the population (511), although this is not proven. It is also possible that the immune system of most children effectively detects the virus with resultant lower average viral loads and thus contagion; however, nasopharyngeal viral loads are not well correlated, whereas saliva viral loads have been correlated with severity (515,516). Regardless, one-third of hospitalized children require ICU stays (517). A pediatric multisystem inflammatory syndrome also has been reported in children who presented with persistent fever and features of Kawasaki disease or toxic shock. Most of those patients tested positive for the COVID-19 virus or for antibodies to the virus, suggesting a post-infection immune response. None of the children have died, but several have required mechanical ventilation (518).

Mortality

The mortality rate for COVID-19 has changed considerably over the course of the epidemic, being much lower through successive waves and variants (519) (61). The mortality risk for omicron was 15% of the risk of delta (520); in South Africa, there was a 90.7% reduction in the death rate with omicron compared with delta. Findings in South Africa also include a 76% reduction in oxygen therapy requirements and a 62.5% reduction in hospital stay (days) (520).

The mortality of COVID-19 was initially estimated to be approximately 10-fold higher than that of typical seasonal influenza (521). Subsequently, severity estimates have been reported as low enough

to be comparable with prior influenza epidemics (432,522,523,524), with a range of infection fatality rates of 0.03–0.5% and corrected rates of 0.02–0.4% (525). The CDC also estimated the overall *symptomatic* case fatality ratio is 0.004, or 1 in 250 (323). Mortality can be predicted based on risk factors and clinical findings on presentation (526).

Mortality risks prior to omicron increased sharply with age, with a prior symptomatic case fatality ratio of 1 in 2000 among those 0–49 years of age, 1 in 500 among those 50–64 years of age, and 1 in 77 among those 65+ years of age (323). The mortality rate for males is 57–64% higher than that for females. Nursing home residence is a particularly potent fatality risk (527,528,529,530,531). The risk of severe disease and/or death is also correlated with other underlying conditions, such as heart disease, hypertension, diabetes mellitus, chronic renal disease, dialysis, liver disease, chronic obstructive pulmonary disease (COPD), smoking, and obesity (380,532,533,534,535,536,537)(538); however, approximately 1% of fatalities occur in previously healthy patients (539). Auto-antibodies to type I Interferon-alpha were found to be more prevalent in patients older than 65 years old, and were associated with an increased risk of severe disease and death from COVID-19 pneumonia (540). The O and Rh- blood group types appear to have slightly lower risk of infection and severe disease(541). Genetic susceptibility (i.e., 3p21.31 gene cluster) has been reported in a large genome-wide association study, along with a 45% increased risk among those with type A blood (542). Past outbreaks of coronavirus infections had considerably higher mortality rates: 34% for MERS and 10% for SARS. However, the mortality rate is not the only factor in determining the seriousness of a disease; a high rate of infectivity and/or easy transmissibility could result in many more total deaths despite a lower case fatality rate.

3.3. BUSINESS CONSIDERATIONS

The actions an employer can take to mitigate the risk of COVID-19 infection are changing as the virus transitions to being endemic and highly contagious with omicron. The actions primarily center on the virus's aerosol and rapid spread combined with lowered morbidity in a largely immunized and/or immune population. Industry sectors with higher risks of workplace COVID-19 outbreaks have reportedly included manufacturing, agriculture, forestry, fisheries, transportation, and warehousing (2). Regardless and importantly, there appears to be a strong potential for a patchwork of different health department regulations in different jurisdictions during the transition from a pandemic to endemic state; naturally, it is important to adhere to those laws and regulations. Managing a company operating in multiple jurisdictions may accordingly become more complex.

One major difference between epidemic and endemic management is that modified **ring isolation** around those at high risk is likely to be effective at protecting those more vulnerable, particularly during or just before forecasted variant surges. This includes:

- greater caution around those at high risk (e.g., cancer, undergoing chemotherapy, other immunosuppressed state);
- institute N95/KN95 for those at high risk, if able to tolerate;
- institute N95/KN85 for those in close contact with those at high risk; and
- reduce and avoid contacts among those at high risk (e.g., avoid grocery stores, other large and densely populated retailers, restaurants, other locations with large numbers of people).

Because some individuals at high risk will not desire some, or any, of these measures (e.g., nearing end of life and/or risk/benefit tolerances/rewards), institution of these measures should only be done after a well-informed discussion and N95 fitting and assessment of tolerance.

There are multiple domains for an employer's actions. Please see the following sections on:

[Employee issues](#) (e.g., education and medical surveillance)

[Travel issues](#)

[Physical distancing methods](#)

[Personal protective equipment](#) (e.g., respirators, masks, gloves, face shields)

[Ventilation issues](#)

[Disinfection practices and contact spread measures](#)

[Policies and procedures](#)

[Industry-specific recommendations](#)

The education of workers in each of these areas is advised as appropriate.

A business with broad geographic interests may also wish to incorporate geographic- and workplace-specific risks (3). This is particularly given that the current vaccination rates vary more than 100-fold across the globe (4), and it can be anticipated that differences by northern/southern hemisphere and other environmental issues (i.e., heat, humidity, UV, use of air conditioning) may persist.

Should a variant arise with significantly greater morbidity arise than the omicron or delta variants, then consideration of re-institution of some public health measures may be indicated. Otherwise, most public health measures are proving of little if any use for omicron and were also of little if any help with the prior variants (5).

3.3.1. EMPLOYEE ISSUES

COVID-19 vaccination

Employers are recommended to strongly encourage vaccination of their entire workforce at the earliest date, including internationally (see also [Vaccination recommendations](#)). The CDC has produced many publications to support these efforts (543,544). Communication to employees regarding their eligibility is recommended. Encouraging household member vaccination also is recommended, as it helps protect the workforce. Other considerations include facilitation of vaccination appointments for workers (e.g., computers at the worksite to access scheduling platforms), incentives, and hosting on-site vaccination clinics.

As the omicron surge has occurred with most of the population protected by vaccines that are partially effective, it is unclear whether the updated vaccine boosters planned for release in Spring 2022 to cover omicron will be necessary. As the projected updated vaccination release will be after the omicron surge has been exhausted, adverse effects on a population basis may be greater due to widespread natural immunity, especially if given too quickly after the surge when the antibody levels are highest. Further, the data for booster benefit are largely in the elderly and those at high risk; there are limited data that boosters make a difference in transmission. Given mild clinical omicron variant presentations in most of the population and unknown long term effects of repeated boosters, further requirements for the young, healthy population currently appear premature

COVID-19 surveillance

Employers do not typically perform surveillance for influenza or other communicable diseases. As the US transitions to an endemic state with the now extraordinarily communicable SARS-CoV-2 virus variants, it may also transition away from employer-based surveillance for COVID-19. The following section retains prior recommendations for this period of transition and in the event that there is another variant with substantially higher morbidity than omicron or delta.

Prior to omicron, employers were recommended to have implemented a surveillance system that included education and screening to avoid having workers with potential asymptomatic, early,

and/or symptomatic but subclinical COVID symptoms enter the workplace premises. Options for larger employers and/or jobs with greater risks (e.g., mission-critical jobs; a workforce where one ill worker could infect an essential group of workers, which would shut down the workplace at least until herd immunity is largely achieved) include daily/periodic electronic questionnaires with or without temperature measurements. Electronic questionnaires are likely to be more effective than temperature measurements because 69% of seriously ill individuals are afebrile (545). Evidence suggests temperature screening is ineffective and detects only one case for every 40 cases missed (546); temperature measurements are also likely to miss all subclinical and many symptomatic cases (323). In the event of a serious surge, diagnostic testing should be performed on those with symptoms, most commonly through the local healthcare or public health systems. Diagnostic testing may also be performed to ascertain asymptomatic spread, especially among essential workers. Testing daily or every few days has been used in some workplaces and among mission-critical workers. However, testing without experienced medical judgment is ill-advised because the false-negative rates are reportedly 20–67% (371); thus, cases with high indices of clinical suspicion should typically be treated as presumptive cases (371). Home testing kits have not been found to be highly reliable for omicron; testing frequency, in order to make a difference for preventing spread, appears to require a minimum of thrice weekly testing of the entire target population. Considerations also include providing communications and expectations to subcontractors, suppliers, and others who may have significant interactions with the employer (e.g., assurance of policies to address symptomatic employees, surveillance).

Employees with possible COVID symptoms

Prior to omicron, sick employees were removed from work even if mildly ill. With the omicron surge, it has become necessary for some mildly affected workers to remain at work with adequate respiratory protection to attempt to prevent transmission to coworkers for a given industry to continue to function (e.g., health care, air travel) (547) (548). Accordingly, the CDC modified the return-to-work provisions to 5 days (549). This phenomenon is typical of an endemic disease where infected and asymptomatic or minimally ill workers are not removed (547) (548). As this is recommended to be a transition point to an endemic state, the remainder of this section addresses when medical removal may be used by employers wishing to remove such employees. Such policies may still be more supportable with the advent of a variant with higher morbidity/mortality than omicron or delta.

Sick employees (including those with minimal symptoms) should stay home from work, as it is important to eliminate all contact between the healthy workers in the workplace and anyone with potentially infectious symptoms (550). If there is believed to be SARS-CoV-2 virus transmission in the area (currently true of essentially all US urban and many rural areas, although the rates are now decreasing markedly), then anyone with even mild symptoms of a respiratory tract infection (e.g., cough, fever, fatigue) should stay home to be sure they do not progress to a clear, and potentially severe, COVID-19 infection (511), as well as to prevent transmission to others. Sick employees should also be encouraged to undergo testing if available. They should be instructed to call a provider or healthcare organization in advance, discuss the symptoms, seek testing if available (especially at outdoor tents), and wear a mask in public settings.

Any questions about potential COVID-19 infections should be directed to the local health department, which has the expertise and personnel to investigate outbreaks and perform contact tracings (provided they are not overwhelmed by the current epidemic). It is important to recognize that return-to-work recommendations for essential workers, especially healthcare workers including volunteers, may need to be modified during the course of the epidemic for practical reasons in response to acute workforce shortages in key jobs and sectors.

Readers are advised to refer to current CDC guidance, as this changes frequently (551) (549). It is also advisable for a healthcare employer to consider factors including staffing needs, infection rates,

and individualized assessment of the degree of that person's contact with susceptible patients (especially those with comorbidities). Furthermore, it is advisable that the other CDC guidance be followed (552,553). Depending on those factors, more conservative or more liberal return-to-work timeframes may be advisable to balance the risks of infecting patients with the ability to staff and care for patients.

What to do if an employee tests positive for COVID-19

The sick employee should follow current CDC guidelines in conjunction with local health department guidance, including isolating at home (if able). A symptom-based approach recommends recording temperatures twice daily until at least 24 hours have passed without fever or treatment with any fever-reducing medications. In order to leave isolation, it is advised that a minimum of 5 days have passed since the onset of symptoms, with then at least 1 day of no fever and improvement in other symptoms. A testing-based approach requires two negative PCR (or antigen) viral tests obtained at least 24 hours apart if there is a need for a shorter waiting time.; however as noted above, PCR tests can mislead as viral particles may be detected but they may not be infectious. Otherwise, testing to return to work is not recommended as viral particles (which may not be infectious) can persist for 90 days after acute infection. The areas where the sick employee worked, including conference rooms and common areas, should undergo deep cleaning and decontamination to prevent spread to other employees.

Employees in contact with an infected coworker

This section assumes a transition back to an epidemic/pandemic state with a much more severe variant than delta or omicron.

If a fully vaccinated worker who is between 2 weeks after the second immunization and 90 days after immunization is exposed to a known/suspected case, quarantine is no longer advised by the CDC (554). Otherwise, employees in contact with an infected coworker should continue to undergo medical screening. Close contacts are defined as any individual who was within 6 feet for 15 cumulative minutes over 24 hours starting from 2 days before symptoms onset (555,556). Risk assessment should include the duration of contact with the sick employee, whether they were using any personal protective equipment, and the type of personal protective equipment used (e.g., cloth face covering vs. respirator) (557). The employer should attempt to maintain confidentiality regarding an ill employee's identity. Employers may wish to apply more or less restrictive policies depending on their individual business requirements, organizational characteristics (e.g., closeness and numbers of other workers), and risk tolerances. For higher-risk exposures with greater business considerations (e.g., mission-critical workers), the most conservative approach is to have employees who could be in the incubation stage self-quarantine and work from home for at least 5 days. The reduction to 5 days was primarily driven by the very high rate of disease transmission, relatively low risk after 5 days, and high and incapacitating numbers of individuals projected to be out of work; they may then be released with monitoring of symptoms until day 14 after the possible exposure. If there is an absence of symptoms, another option is to quarantine for 7 days; with a negative test on day 5 or later, the person may be released on day 8 with ongoing monitoring until day 14 (558,559). The CDC has changed their quarantine recommendations for exposed but asymptomatic workers to wearing a mask and testing if possible on day 5.

In certain manpower shortage situations, medical centers, and critical services, COVID-19 exposed workers are being allowed to work while asymptomatic with self-surveillance for symptoms, physical distancing, disinfection of workspaces, and consistent mask-wearing instead of being quarantined (560). This option is becoming less controversial, although it is not without risks because pre-symptomatic spread is believed to be a primary source of epidemic spread. This option should be carefully weighed between the industry sector, criticality of the job, job requirements, and risks of an infectious individual in that particular workplace. This option is likely unduly risky if the workforce or work group is mission critical.

High-risk employee issues

For the purposes of these recommendations, high-risk individuals have any of the following conditions (545,561):

- age 65 years and older
- chronic lung disease, including moderate to severe asthma
- serious heart condition (e.g., history of heart attack or heart failure)
- immunocompromised (e.g., having had bone marrow or organ transplantation, immune deficiencies, poorly controlled HIV or AIDS; using corticosteroids or other immune-modulating medications; undergoing cancer treatment)
- smoking, current or former
- obesity, especially severe (532)
- diabetes mellitus
- chronic kidney disease, especially those undergoing dialysis
- liver disease
- hypertension
- current cancer
- neurological diseases, including stroke and dementia

Generally, the risks of severe illness associated with the above conditions are greater as the severity of the conditions increase. The presence of multiple conditions increases the risk of severe disease (79).

Employers should attempt to reduce exposures to higher-risk situations for workers who self-identify as high-risk, while being cognizant of the implications of the Americans with Disabilities Act and amendments. A full- or part-time medical director and medical department may help to interface between the worker and management to effect these risk assessments and potential risk reductions. Examples of reductions in exposure (beyond electronic questionnaires with or without temperature checks) include the following:

- emphasize distance-based work methods, including telecommuting where feasible.
- place all, but especially high-risk, individuals behind barriers.
- institute physical distancing (10).
- reduce public-facing work.
- provide the affected worker with a fitted N95 or KN95 respirator to protect from exposure.
- consider placing high-risk individuals closer to ventilation that provides fresh air.

Some educational videos have been developed to illustrate the spread of aerosols, as they are now thought to be the primary mechanism of disease spread ([video 1](#), [video 2](#), [video 3](#)).

3.3.2. TRAVEL ISSUES

Travel bans, restrictions, and broad testing are no longer recommended. They are central tools for early control and management of an epidemic. However, these tools are not useful to manage an endemic disease. They are also especially not useful for a variant that moves rapidly and without quick detection to a peak surge in 1-2 months in essentially all countries. Such restrictions may again be needed should a future variant surge have markedly higher mortality rates above that associated with the delta or omicron variants. Still, there are risks, especially for some employees on business trips. Travel risks include those associated with travel to and from a site, as well as business conducted at those sites (6). Risks differ considerably by mode of transportation, geographic locations, current state of the epidemic in any given locale, and vaccination rates. Businesses need to weigh the value of the travel against the risks associated with that travel.

Fully vaccinated employees may reasonably travel. For non-vaccinated employees, especially those not known to have natural or vaccine-related immunity, travel valuations should include costs associated with any potential illness and any post-trip quarantine period. Caution is advised for non-essential travel by non-vaccinated and non-immune employees to locales with outbreaks or community spread in progress (6), which includes the entirety of the United States and most of the world (see [map](#) to help with other risk considerations (7).

International trips continue to be significantly affected as many countries are limiting travel from countries with outbreaks, despite the increasingly dubious value of these restrictions as demonstrated by the universal rapid spread of omicron. Air travel may be safer than some other forms of travel (8), although the primary risks of air travel are more likely to be exposure risks at the destination, which may be challenging or impossible to control.

Employees returning from, or having traveled through, areas known to have COVID-19 infections

For vaccinated employees returning to work after travel, limitations or quarantines are not advised. For non-vaccinated and non-naturally immune employees returning from personal or work-related travel to areas with community-based COVID-19 spread, the safest course of action is to self-quarantine while working from home for 5 days.

3.3.3. PHYSICAL DISTANCING METHODS

In the transition to an endemic disease, physical distancing is increasingly of less use, particularly for what has become an easily-transmitted, aerosol-spread disease. The primary exceptions involve ring isolation to protect those who have major immunosuppressed conditions (e.g., leukemia; undergoing chemotherapy treatment). Otherwise, physical distancing methods are no longer recommended.

Institution of effective physical distancing controls for immunosuppressed people is challenging. With the recognition that the disease is mostly aerosol spread and apparently outdoor spread (e.g., widespread infection among wild deer populations), physical distancing may need to be at least 30 feet, while also avoiding being downstream or crossing “vapor trails” from other people for an as-yet unclear length of time which may well be at least the past 30 minutes, and using an N95/KN95 respirator for added protection. These criteria preclude grocery shopping, walking in malls, going to church, walking on a sidewalk in a large city, and most other common experiences of life. Thus, the practicality of physical distancing, even for immunocompromised individuals, is increasingly dubious.

Physical distancing was believed to be one of the more effective control measures, particularly when the disease was believed to be due to droplets and contact, because it does not rely on training and compliance, and in the initial stages was thought to involve a 6-foot distance (9). The following are some physical distancing options to consider if a considerably more lethal variant arises:

- work from home when feasible.
- consider rotating workers between home and work settings to reduce workplace population densities while facilitating functions that are best performed at work.

The following are examples of commonly instituted physical distancing methods that are likely to be either minimally or wholly unsuccessful with aerosol-spread viruses and are no longer recommended:

- **NOT RECOMMENDED:** improve physical distancing at work (e.g., increase distances between workers and workstations to a minimum of 6 feet, install temporary barriers, mark 6-foot distances on the floor between co-workers).
- **NOT RECOMMENDED:** consider either physical spacing in cafeterias, closing cafeterias and offering individual prepackaged meals, using disposable packaging and utensils to avoid the potential for contamination before cleaning, and/or having workers eat their own food at their workstations.
- **NOT RECOMMENDED:** where there are two options for walking through a workplace, set up one-way walkways.
- **NOT RECOMMENDED:** reorganize shifts to spatially and temporally spread workers.
- **NOT RECOMMENDED:** route shifts of workers to enter through one entrance and exit through a different one.
- **NOT RECOMMENDED:** provide protection for those who interact with the general public (e.g., install temporary barriers to prevent respiratory transmission, install barriers to ensure physical distancing of 6+ feet).
- **NOT RECOMMENDED:** consider discouraging carpooling and mass transit; encourage the use of masks if using either of those options (although a face mask in public places is now a requirement in many cities and states).
- **NOT RECOMMENDED:** minimize reasons for external individuals and the public to enter a workplace (e.g., curbside deliveries, web-based meetings). If there are multiple options for meetings onsite, attempt to limit which rooms are used and have them cleaned after every use.

3.3.4. PERSONAL PROTECTIVE EQUIPMENT

PPE recommendations have changed with the endemic nature of the virus, recognition that the virus is aerosol-spread, vaccination of the population, and the impairments of communication by masking. PPE is no longer recommended for immunocompetent individuals in the general population. As the virus is aerosol spread, and in the potential future event of a variant with much higher morbidity, N95 respirators and other devices (e.g., PAPR) would be a priori recommended. N95/KN95 respirators are selectively recommended for those with immunocompromised states.

PPE measures (respirators, masks, gloves, and eye protection/face shields (10,11,12,13,14,15) were recommended, but they are lower on the list of controls. The speed with which the omicron variant spread and failure to slow or blunt the epidemic curves, irrespective of state or country masking mandates and business openness status highlights the apparent ineffectiveness of masks and other methods at this stage of the pandemic with this degree of aerosol-spread contagion.

Reuse, Extended Use, and Reprocessing of Respirators

The pandemic initially caused demands on all types of PPE far beyond manufacturing capacities, which has been subsequently alleviated. However, a future surge of a variant with high morbidity could easily exhaust supplies of N95 respirators. In such a case, extended use and re-use of respirators will be essential.

3.3.5. VENTILATION ISSUES

Ventilation issues (general and local fresh air supply) have been markedly underutilized as potential COVID controls (16,17,18,19,20). Ventilation may still be of some assistance with management of aerosols, although it may be less effective than for management of a droplet-spread virus. This issue also has potentially major implications for the future reduction in other epidemics, such as influenza or resurgences of COVID-19. Consultation with an HVAC expert may be helpful. Area ventilation may provide a relatively safe zone for workers. The following general ventilation measures can be used to dilute viral concentrations:

- identify the number of air exchanges per hour (ACH) in the room.
- increase ACH in work areas. The number of necessary ACH depends upon occupancy of the area and the purposes for which the area is used (e.g., more ACH in healthcare or crowded areas than in sparsely populated warehouses).
- assure homogeneity of airflow to avoid “dead spots” and short-circuiting from air supply to exhaust.
- run the ventilation system as many hours as possible.
- increase the proportion of fresh (rather than recirculated) air.
- filter and/or disinfect the air.
- use effective filters in the HVAC system. HEPA filters are optimal, but some ventilation systems cannot effectively overcome their added resistance. A minimum filtration efficiency rated at least MERV 13 should be used (21,22).
- air disinfection, such as ultraviolet germicidal irradiation, can be placed within the central HVAC system (19,22). Use portable air cleaners and local exhaust.
- local standalone HEPA filtration in high-risk areas may be potentially helpful for risk mitigation.
- fans and other airflow and/or filtration devices may be used to control the direction of airflow from clean to potentially contaminated areas. Where possible, consider using portable air purification systems for small work areas, generally 8-10L/s.

3.3.6. DISINFECTION PRACTICES AND CONTACT SPREAD MEASURES

Ventilation and other control measures addressing aerosols and microdroplets are far more important than disinfection of surfaces for COVID-19 (23). Disinfection of surfaces may have some limited role in reducing spread. Disinfection of surfaces has not been shown to be effective, distracts and diverts attention from more effective control measures, incurs considerable time and expense, and is no longer recommended for purposes of control of the COVID-19 pandemic.

3.3.7. POLICIES AND PROCEDURES

At this point in the COVID-19 pandemic, it is anticipated that subsequent variant surges may be relatively mild. Regardless of whether the next variant is mild or severe, now is the time for comprehensive after-action reviews (i.e., critical incident stress debriefings). The purpose of such reviews is to define successes and failures, and plan how to better respond in the future. To be successful, these reviews require a focus on why things happened, avoid finding fault, identify the approaches that did or did not meet expectations, and require broad participation from all key stakeholders. These reviews should include public health officials from federal, state, and regional/local levels, healthcare organizations, as well as representatives from large and small employers, with the goal to define effective responses to the pandemic and prepare for the next viral surge or epidemic/pandemic.

The following are potential policies and procedures to consider:

- inform and seek support and authorization for the plan from the organization's leadership.
- develop a plan in conjunction with occupational health and safety professionals, government regulations, and public health authorities (including the CDC).
- ensure affected workers have sufficient paid leave to observe a quarantine period or are able to stay home as indicated.
- continue to monitor sickness absence, but expand sick leave provisions to allow employees to stay at home if ill and to care for sick family members.
- teach workers about the principles of microdroplet/aerosol spread viruses.
- teach workers to use tissues to catch a cough or sneeze, then throw that tissue away and wash their hands.
- in the event of a surge with a high morbidity variant, avoid scheduled aggregate meetings. Encourage use of teleconferences and/or other virtual meeting formats.
- in the event of a surge with a high morbidity variant, consider instituting required daily electronic symptom trackers with an automated management system for all employees to report symptoms of COVID-19 infection, including fever, cough, shortness of breath, myalgias, abdominal discomfort, and diarrhea. Responses should be monitored daily by the medical department or health and safety (24,25,26,27).
- in the event of a surge with a high morbidity variant, and if daily symptom tracking is not instituted, encourage early reporting of any symptoms consistent with COVID-19 to the medical department, designated employer representative, and/or supervisor, following the company's established policies. It is preferable to preclude all symptomatic workers, including those who are mildly symptomatic, from physically entering all workplaces; electronic questionnaires may be useful to facilitate this. Place posters prominently to help remind workers of procedures (e.g., CDC posters).
- in the event of a surge with a high morbidity variant, have employees who develop symptoms stay away from the workplace until clinically evaluated and/or until the symptoms are resolved and any quarantining period has expired.

- in the event of a surge with a high morbidity variant, consider having employees who could be in the incubation stage work from home for at least 10-days after the possible exposure.
- in certain manpower shortage situations, medical centers and critical service workers are being allowed to work while asymptomatic with self-surveillance for symptoms and consistent mask-wearing instead of being quarantined for 5-days. However, this has some residual risks of transmission and may not be compatible with mission-critical operations (e.g., dispatch center, air traffic control tower).
- antibody testing is now widely available, but the sensitivity and specificity vary greatly between kits (see [Diagnostic Testing](#)). With further validation, antibody testing may likely become useful in assessing possible susceptibility to infection versus protective response to prior infection. In the future, COVID-19 serology can determine infection risk in critical and susceptible populations (under medical direction to ensure proper implementation, interpretation, and management). Examples of these critical populations include employees in health care settings, oil drilling platforms, commercial maritime, food preparation, cruise lines, airlines, and assembly lines with workforces working closely together.
- provide proactive assistance to support mental health for the workforce.
- identify and train workplace coordinators who will be responsible for implementing and monitoring the plan.

3.3.8. INDUSTRY-SPECIFIC RECOMMENDATIONS

Below are select industry guidelines, which are in addition to the general guidance above. These guidelines assume lack of herd immunity and/or ongoing community-based spread. Further guidance is available from the CDC (21).

Restaurants

- Menus should ideally be either disposable or laminated and sanitized after each customer contact. Other options are electronic access and use of QR codes.
- Clean and disinfect chairs and tables between use (see [Disinfection Practices](#)).
- Assign high-risk employees with multiple co-morbidities or concerns to low-exposure areas, such as working in non-customer-facing areas as much as possible.
- Encourage drive-through and carryout options

Retail

- Stocking by high-risk individuals should ideally be done when customers are not present.
- Encourage customers with high risk to use N95/KN95 respirators and provide N95/KN95 respirators to such affected customers where feasible.

Gyms

- Gym housekeeping should ideally be performed by low-risk employees.
- Increase local and area ventilation to address increased the increased respiratory rates associated with exercise.

Manufacturing

- Evaluate and improve [ventilation](#).

Food Production Facilities

- Evaluate and improve [ventilation](#).

Schools

Schools have high human population densities. However, extensive data show that children have the lowest risk of symptomatic, severe, and/or fatal COVID-19 disease across the lifespan, with the risks appearing to be lowest in the youngest school-age children (28,29,30). Data to explain these observations are sparse; theories include that children have relative lymphocytosis, superior immunity to coronaviruses, and an ACE2 receptor (to which the virus binds to gain entry) that is inadequately developed in their airways (31,32). Initial reports that children do not become infected appear increasingly dubious (33); however, that they are resilient to symptomatic and/or severe disease is not in question.

Schools in most countries were at least temporarily closed in spring 2020 in response to the pandemic. However, students' learning by distance-based methods has been reportedly suboptimal and sometimes poor. The burden of the inability to educate students using traditional methods also disproportionately falls on the poor and immigrant populations, which have fewer skills and resources to educate and/or guide their children's learning (34,35,36,37,38). For example, increases in computer search intensity for school-centered resources in higher socioeconomic US regions were double those of lower socioeconomic status regions in April 2020 compared with 2015–2020 (36). A 5-month global shutdown of schools has been estimated to have had an adverse worldwide impact, with a loss of \$10 trillion of lifecycle earning for the 1 billion affected students because of lower levels of learning, lost months, or dropping out of school (39). Schools also play important roles in students' social development and mental health (40,41,42).

Restarting of schools was controversial, more so in the US than elsewhere, and widely divergent strategies were deployed. Nearly all reports suggested few problems with most re-openings in Belgium, Denmark, Finland, France, Japan, Norway, Germany, Quebec, Singapore, South Korea, and Sweden; these reports have also included opening without physical distancing, masking, alternate school schedules, or other mitigations (43). The main contrary example is Israel, where school-based transmission to teachers was briefly noted (44,45).

The worldwide omicron surge produced nearly identical epidemic curves, regardless of whether the country or state had in-person mask-less teaching on one extreme to distance-based education on the other extreme of a spectrum. Accordingly, a rationale for masking students and/or not having in-person classes is not apparent.

For those teachers and other staff who have severe immunosuppressed states, there may be modest protection with fitted N95 respirators and use of distance-based teaching methods.

There may be a consideration for institution of distance-based teaching methods with a future surge; however, this would not be recommended due to adverse effects, including marked reductions in learning associated with lockdowns (46), unless a subsequent variant causes markedly higher morbidity and mortality than delta or omicron. If subsequent variants become more transmissible, it is also highly improbable that masks be effective in schools regardless of any potential increases in morbidity. Thus, due to an absence of evidence of efficacy and evidence of harms, including marked reductions in learning associated with lockdowns (46), physical methods such as masking and distancing are not recommended for schools.

3.4. DISABILITY AND RETURN-TO-WORK CONSIDERATIONS

Disability from COVID-19 will be better defined with studies over time. Extrapolation using recovery from other conditions, such as pneumonia and ARDS, may provide some preliminary estimates.

Preliminary reports suggest recovery duration is, unsurprisingly, at least partially correlated with measures of case severity. At least one symptom persisting for at least 60 days has been reported among hospitalized survivors, with the most prevalent symptoms being fatigue, dyspnea, joint pain, chest pain, cough, and anosmia (303). In a cohort consisting primarily of patients treated for COVID in an outpatient setting, 39% reported symptoms at 7-9 months, with fatigue (21%), loss of taste or smell (17%), dyspnea (12%) and headache (10%) being the most common (562). However, persistent symptoms are reported in individuals with mild cases, and long-term symptoms have been reported (563). There also are many cases that require home healthcare after discharge (564).

Permanent disability is determined by the existence of some combination of fixed deficits when a healing plateau has been reached (see the [ACOEM Disability Prevention Guideline](#)). One of the greatest factors facilitating recovery is the interest and ability of the employer to reintegrate the employee into their workforce. Such integration often requires accommodations that hopefully can be reduced as time, recovery, and workarounds progress. While not yet demonstrated for COVID-19, employer support for recovery is critical for many other conditions.

Permanent disability is only appropriate for those with fixed, non-improving chronic impairments (see [Rehabilitation](#)). Some of these cases have obvious permanent deficits from complications such as myocardial infarction and stroke. There is also increasing literature supporting the development of chronic symptoms associated with COVID, which is elsewhere termed “ongoing symptomatic,” “post-COVID syndrome,” and “long COVID” (565). The term “post-acute sequelae of COVID” and “post-acute sequelae of SARS-CoV-2 (PASC)” has also been used by the National Institutes of Health.

Factors contributing to disability beyond fixed but remediable deficits can include a lack of full implementation and utilization of evidence-based treatments, and lack of effort and compliance. Other factors may potentially involve advocagenic, psychological, and other influences.

Return-to-work evaluations should consider the worker’s current status as compared with the physical requirements of the job, mental demands of the job, safety-critical work functions, current treatments, use of impairing medication, residual effects of the virus, requirements for personal protective equipment, potential risk to others if returned too early, and protection of other employees if additional risk is identified. Many of these complex cases will need to be addressed by occupational and environmental medicine physicians.

Currently, for patients without hospitalization, there are no quality data on returning to work, short-term disability, or long-term disability. One random sample (n=292) of affected individuals diagnosed as outpatients reported 65% had returned to normal health at a median of 16 days; no or few comorbidities and age statistically impacted those rates, with 74% among those 18–34 years of age, 68% among those 35–49 years of age, and 53% among those 50 years and older returning to normal health (566). Regarding short-term disability and return to work, recovery from post-infection fatigue is estimated to take approximately 2–3 weeks and appears to correlate with clinical duration and severity. For patients with mild to moderate pneumonia treated with oxygen supplementation, recovery is estimated to require 4–8 weeks after hospitalization or clinical recovery. Severe pneumonia and ARDS have worse prognoses.

The overall trajectory of recovery from COVID-19 remains unclear. Prior experience with diseases that have similar manifestations, such as ARDS, suggest there is significant risk of delayed return to work and long-term disability, as approximately 50% of individuals surviving ARDS have not returned to work after 1 year (567,568). ARDS is also associated with approximately 20% reductions in spirometry and lung volume, which resolve at about 6 months based on prior H7N9 influenza data (569). Lung diffusion abnormalities can take up to 5 years to resolve in ARDS cases (569,570).

Cognitive impairments and psychiatric abnormalities related to ARDS may be projected to occur in 30–55% and 40–60% of patients, respectively; the duration of these impairments is unclear, but other causes of ARDS raise considerable concerns about long-term disability (568,569,570,571,572,573,574). Generalized skeletal muscle deconditioning is expected in patients who are intubated for any extended duration; these patients require graded return to activity and/or exercise programs and possibly rehabilitation, which often results in residual incapacity (568,571,575,576). Cardiac problems are common with COVID-19, with cardiomyopathy, arrhythmia, and direct cardiac muscle injury affecting approximately 30%, 20%, and 10% of patients, respectively (258); they are contributing causes to fatality (258,259,260).

In general, for patients who are intubated and survive, recovery of the cardiorespiratory systems and endurance are estimated to take at least several months. Among recent COVID-ARDS survivors, 78% had evidence of cardiac involvement and 60% had evidence of ongoing myocardial inflammation on MRI (261). It currently appears likely that some hospitalized and severely affected individuals will incur long-term disability with permanent impairments of the cardiac, respiratory, neurological, and/or musculoskeletal systems (568,569,570,571,572,577). There is also the potential for a minority of patients to be permanently totally impaired (572).

Cardiac, respiratory, and neurological disability measures include the following:

- 6-minute walk test and/or sit to stand testing
- metabolic stress echocardiogram (including ECG)
- full pulmonary function testing with impedance booth or washout testing
- high-resolution CT scan of the chest, especially for those with COVID-19 pneumonia
- functional capacity testing (although there are some limits in interpretation)
- neuropsychological testing

For individuals with less symptoms but high exertion requirements, a cardiac evaluation may be indicated.

An approach to evaluating COVID-19 worker's compensation claims has been published (578). There is no specific impairment class for COVID-19 and surrogate diagnoses may be needed and/or used by analogy. Ratings for impairment can be found in the AMA Guides 5th Edition (579) and 6th Edition (580).

4. VACCINES

Vaccine development progressed at record speed on more than 270 COVID-19 vaccine candidates (550,581,582,583). These efforts used at least four types of vaccine classes or approaches against this infection (virus, viral vector, nucleic acid, and protein-based) (582). Although vaccine development was estimated to require 12–18+ months if successful, it was achieved in approximately 9–10 months (584). Several more of these COVID-19 vaccines are in advanced stages of development and have potential for approval. Efficacy data have been published. Safety data suggest the vaccines are largely safe. Reported rates of initial vaccine efficacy range from 62% to 95% (585) (586) (587) (588). After initiating vaccination programs, COVID-19 infections declined markedly (589). However, vaccine efficacy fell with subsequent waves of new variants (590). Accordingly, booster shots which have increasingly been suggested to have retained partial efficacy against the delta and omicron variants have been emphasized (590,591,592).

The [London School of Hygiene & Tropical Medicine's COVID-19 Vaccine Tracker](#) is updated weekly with multiple COVID-19 vaccine databases, including a vaccine pipeline tracker, clinical trials

database, and living review (581,588). The CDC has also provided guidance regarding what is recommended for those who have been vaccinated (593).

The vaccines have very good to excellent rates of efficacy both in randomized trials and in reports from large population-based studies and surveillance systems, which underscores evidentiary support for broad-scale vaccination programs. The following questions require answering going forward, although they should not delay the continued implementation and completion of the vaccination programs:

- duration of vaccine-induced immunity and whether there are differences between the types of immunizations
- whether duration of immunity differs in different subgroups, suggesting the need for (earlier) re-vaccination
- whether immunity is shorter-lived in vaccinated patients or in naturally infected patients
- whether annual immunizations are needed
- utility and/or adverse effects among those who have been infected with COVID-19
- long-term adverse effects
- whether the vaccine is safe in the elderly
- risk/benefit ratios in children, whether all children should be immunized, or whether natural infection is preferable whereby immunity is more durable (594).

4.1. ADVERSE EFFECTS

The vaccines have been associated with a relatively low frequency of adverse effects. Even though vaccine reactions are rare, it is important to address them because they may generate fear, anxiety, and vaccine avoidance that is out of proportion with the actual prevalence of these outcomes. The earliest reports appeared in the January 6, 2021 *MMWR* (47), which described data collected from the December 14–23, 2020 period of vaccine administration of the Pfizer-BioNTech COVID-19 vaccine. Out of 1,893,360 first doses administered, there were 4,393 (0.2%) adverse events reported. After reviewing all cases, only 175 cases were considered to be consistent with a severe allergic reaction; of these, only 21 cases were deemed to represent anaphylaxis, for a rate of 11.1 per million doses administered. Nonallergic adverse events, mostly vasovagal or anxiety-related, were excluded from analyses. The median age of those with anaphylaxis was 40 years, and 90% were women. Typical symptoms included a diffuse erythematous rash, throat closure, hoarseness, swollen lips, difficulty swallowing, wheezing, cough, and nausea. Most (17/21; 81%) had a prior history of allergic reactions to drugs, medical products, foods, and insect stings, and 9.5% (2/21) had prior reactions to a vaccine. Most (19/21; 90.5%) were treated with epinephrine, and no deaths were reported. There was no geographical clustering of cases or associations with any specific vaccine lot. There were 83 cases of non-anaphylactic allergic reactions, with a similar age and sex distribution, and 56 (67%) also had a prior history of allergies or allergic reactions. Almost all reactions occurred in the first 30 minutes after vaccine administration.

A review subsequently published noted that confirmed allergic reactions to vaccines are usually not to the active ingredients, but rather due to reactions to excipients (48). Reactions specifically focus on polyethylene glycol (PEG) and polysorbate, which have been added to multiple other vaccines, injected medications, chemotherapeutic agents, and biologicals to increase water solubility. These

excipients are also found in multiple creams, ointments, lotions, and personal care products. Multiple existing vaccines contain polysorbate 80, including the AstraZeneca and Johnson & Johnson vaccines, and both the Pfizer and Moderna vaccines contain PEG2000. A recent study of the general population found that 5 to 9% of serum samples were positive for anti-PEG IgG (49). Skin tests for polyethylene glycol are available, and other medications containing PEG3350 (methylprednisolone acetate), polysorbate 80 (triamcinolone acetonide, Refresh eye drops, Prevnar) or polysorbate 20 (hepatitis A vaccine, Twinrix) can be used for skin testing to document an allergy to one of these excipients. The authors proposed a risk stratification to determine who should undergo pre-vaccination skin testing or extended observation postvaccine, using the following patient-directed questions:

Do you have a history of a severe allergic reaction to any of the following:'

1. an injectable medication (IV, IM, or SQ)?
2. a prior vaccine?
3. another allergen, such as food, venom, or latex?
4. polyethylene glycol (PEG), a polysorbate, or a paclitaxel-containing injectable or vaccine?

If the patient answers “yes” to question 4, he or she is higher risk and should be referred to an allergist before receiving the vaccine. Questions 1, 2, and 3 represent medium risk; the patient should be observed for 30 minutes after the vaccine. If the patient answers “no” to all four questions, then he or she is lower risk and should be observed for 15 minutes after the vaccine.

Delayed large local reactions to the Moderna vaccine occurring 8 to 12 days after vaccination have been described (50). Of these, 10 were women, 8 had a prior history of allergy or allergic reactions, 9 described itching, 9 described pain, and 7 described fatigue or other systemic symptoms. Most were treated with antihistamines and topical steroids, and two received oral steroids. Reactions resolved by day 14 to 19. All then received the second vaccine dose, with only minor rash or itching reported; none were severe.

In addition, there have been reports of 36 cases of immune thrombocytopenic purpura (ITP) following the vaccination of 31 million people as of February 8, 2021, but no cases were associated with any one vaccine or vaccine lot. The majority of patients received platelet transfusions, IVIG, and/or steroids along with hospital care; there was one reported death. Importantly, ITP has been associated with other vaccines, including the MMR, DTaP, varicella, hepatitis B, and pneumonia vaccines (51), as well as following viral infections. For patients with a pre-existing history of ITP, the American Society of Hematology recommends that platelet counts be checked before receiving the vaccine; however, the presence of ITP is not a contraindication to receiving the vaccine.

A case series has reported ongoing symptoms among those having received one dose of vaccine and who had been previously hospitalized with COVID (52).

Thus far, probably the most important adverse effect appears to be myocarditis among young males. The population-based risk has been estimated at 5.3-fold overall, with highest risk of 13.6-fold among those 16-19 years old males (52).

4.2. VARIANT CONCERNS

The spike protein of the SARS CoV-2 virus is the focus of currently available vaccines. A parade of successive variants have developed and most that become common, have more genetic variation and consequential transmissibility. The omicron has had, by far the most mutations in the spike protein to date.

The spike protein is the primary viral protein responsible for entry into host cells by attaching to the ACE2 cellular receptor present on multiple human tissue cells, including the lungs, heart and blood

vessels, kidney, testis, and brain. The primary antibody response elicited by the virus in natural infections is directed against the spike protein. Hence, as the spike protein appears to be the preferred target of the natural immune response, it was naturally selected as the primary target for the vaccine response.

The first significant variant of the SARS CoV-2 spike protein, D614G, was detected in early March 2020, substituting a glycine for an aspartic acid in the carboxy terminal region of the S1 domain. Not present in any of the viral sequences in January and February 2020, it constituted 26% of viral sequences in March and 70% in May, attributed to enhanced ACE2 binding affinity and infectivity (53,54).

The next set of more transmissible variants, all containing adaptations in the spike protein, were identified in the fall of 2020 and include B.1.1.7 (alpha, UK), B.1.351 (beta, South Africa), and P.1 (Brazil). The B.1.1.7, or beta variant, was first identified on September 20, 2020 in Kent, England. It is thought to have arisen in a patient with an impaired immune system who was treated with antibodies from a recovered patient, and possibly also with remdesivir (55). With this patient's specific scenario, the virus would theoretically have the opportunity to replicate multiple times, increasing the odds of random mutations, and under the pressure of antibodies targeted to the spike protein. Hence, those variants that survived could develop slightly different spike proteins that are less well recognized by existing antibodies. This variant carries a N501Y mutation of asparagine to tyrosine in the S protein that increases its binding strength to the ACE2 cellular receptor, as well as a deletion at positions 69 and 70, which are both hypothesized to increase transmissibility. The deletion causes S-gene target failure in one PCR-based assay, the ThermoFisher TaqPath COVID-19 assay, producing a negative result for the S-gene target and still positive results for the other two targets.

By January 12, 2021, the B.1.1.7 alpha variant had been detected in 12 U.S. states. This became the dominant strain in the United States in 2021 (56).

The beta B.1.351 variant independently emerged in South Africa; it was first detected in the US at the end of January 2021. It carries eight specific mutations in the spike protein, along with the N501Y variant carried in the UK strain.

The vaccines proved effective against the alpha and beta variants. However, one report suggested a poor ability of the ChAdOx1 nCoV-19 vaccine to prevent mild to moderate COVID illness caused by the B.1.351 strain (57).

The P-1 gamma variant of SARS-CoV-2 emerged in Manaus, Brazil, and was detected in the United States at the end of January 2021. This variant carries 20 unique mutations, including three identified in other variants in the receptor binding domain of spike protein (K417T, E484K, and N501Y). A separate study showed that serum samples from subjects immunized with the BNT162b2 (Pfizer) vaccine effectively neutralized engineered CoV-2 viruses carrying all the identified variant spike proteins, most at titers >1:40 (58).

The most important variants to subsequently arise are the Delta (B.1.617.2) and Omicron (B.1.1.529)(59). Delta became the predominant variant by summer 2021 (REF) (60). The Omicron variant, however broke new ground. It both had far more mutations, while far more transmissible, and from detection to becoming the predominant strain only took approximately 6 weeks (61).

All settings of natural and vaccine-induced immunity will exert selection pressures against the virus and drive the emergence of resistance mutations. One study cultured a SARS-CoV-2 recombinant virus in the presence of 18 different neutralizing monoclonal antibodies that were selected for different RBD mutations. In all cases, the antibody selected for the emergence of a resistant variant. This same study also demonstrated that antibodies elicited by either the Moderna (mRNA-1273) or Pfizer BioNTech (BNT162b2) vaccine were nearly identical and were effective against the dominant variant of SARS CoV-2 (D614G), with only a modest decrease in the ability of these antibodies to

neutralize viral variants (62). This likely reflects the polyclonal nature of neutralizing antibodies elicited by the vaccines—that is, that the mRNA carried by these vaccines codes for a number of different proteins with many different antigenic epitopes. Antibody responses will correspond to multiple epitopes, including many sites that remain unchanged in different variants of the virus.

Although the intense scrutiny of the SARS CoV-2 virus has resulted in early identification of viral variants, their emergence should be considered a normal process in a pandemic and will continue as the transition to an endemic disease proceeds. As host susceptibility to infection changes, the virus, under these selection pressures, will change accordingly. More variants will emerge, and in general, it is in the virus' best interest to become more transmissible and less lethal, producing an expectation of generally diminished severity with successive waves. This may, or may not, have an effect on host susceptibility. Thus far, vaccine-elicited antibodies have been shown to remain active against spike protein variants. Most SARS CoV-2 specific CD4+ and CD8+ T-cell responses from both naturally infected and vaccinated subjects are equally effective against variant strains (63). While the vaccines are being altered going forward to address novel variants that have already emerged, as well as those yet to emerge, the speed with which the omicron variant arose raises practical questions about the ability to develop a timely targeted vaccine. Nevertheless, the original vaccine given as a booster shot has shown an ability to reduce severe disease including hospitalizations (64).

4.3. VACCINE RECOMMENDATIONS

Vaccines for the Prevention of COVID-19

Recommended

Vaccination is strongly recommended for the prevention of COVID-19.

Strength of evidence Strongly Recommended, Evidence (A)

Level of confidence High

Indications

Indicated for nearly all adults. Particularly indicated for those with increased risk of severe COVID-19 disease (e.g., elderly, obesity, diabetes mellitus, COPD, cardiovascular disease, renal disease, immunosuppressed states).

Common RCT exclusion criteria include pregnancy, immunodeficiency, immunosuppression, use of glucocorticoids 20+ mg/day in the past 6 months, and prior vaccine allergic reactions. However, studies also now suggest there are no increased adverse risks among pregnant women.

How, and whether to integrate prior COVID-19 infection into the decision to vaccinate and timing of vaccination is somewhat controversial, although evidence supports durability of the antibody responses from natural immunity. Regardless, CDC guidance recommends vaccination irrespective of prior infection(s) (Magnus et al., 2021, Shimabukuro et al., 2021, Zauche et al., 2021, Kharbanda et al., 2021). Regardless, those with immunosuppressed states would be potentially high-impact populations and are recommended to receive early vaccination.

There is evidence from large population-based studies that booster immunizations raise antibody levels and markedly lower risk of mortality or hospitalization (Choi et al., 2021, Intapiboon et al., 2021, Li et al., 2021, Munro et al., 2021)(Thompson et al., 2022)(Schmidt et al., 2022, Accorsi et al., 2022, Arbel et al., 2021, Dhingra et al., 2022, Mattiuzzi et al., 2022).

CDC recommends a single booster immunization at 5+ months for the Pfizer-BioNTech and Moderna vaccines and at least 2 months for the Janssen/Johnson & Johnson vaccine (CDC, 2022)(Mbaeyi et al., 2021).

Benefits

Reduced risk of COVID-19 infection, as well as serious COVID-19 disease. Two 2-shot series of mRNA vaccines (Pfizer and Moderna) have ~95% efficacy, whereas the single shot (Janssen/Johnson & Johnson) has ~67% efficacy (FDA, 2021). Booster immunizations have reportedly resulted in ~90% reductions in risk of hospitalization and death.

Harms

Serious adverse effects shown in a 1.7M population-based study of the Pfizer vaccine were a 3.2-fold risk of cardiomyopathy, 2.4-fold risk of adenopathy and 1.4-fold risks of both herpes zoster and appendicitis (Barda et al., 2021). There are no similar high-quality data yet published for the other vaccines.

Reported rates of adverse effects from a passive but large-scale surveillance system (V-safe) include injection site pain (Pfizer/Moderna; Pfizer dose #2; 73-78% after first dose and 79% after second dose), fatigue (22-25%/25-54%), headache 15-23%/20-43%), myalgia (15-23%/18-47%), chills (6-11%/8-31%), fever (6-11%/8-29%), injection site swelling (6-11%/9-13%), joint pain (5-10%/7-24%), and nausea (4-9%/6-14%) [316, 335, 336]. The population-based risk estimate for cardiomyopathy has been estimated at 5.3-fold overall, with highest risk of 13.6-fold among those 16-19 years old males.

Anaphylactoid reactions are quite rare (4.5 per million doses administered (Gee et al., 2021); those with severe food and/or medicine allergies have been suggested to delay getting the vaccine. Pfizer BioNTech/Fosun Pharma: Grade 3 adverse effects >2% were fatigue 3.8% and headache 2.0% (Soiza et al., 2021).

There is an increased risk of shoulder conditions, particularly among women and elderly recipients (Zheng et al., 2022), which may be related to accidental subacromial injection especially among those with lower deltoid muscle mass.

Frequency/Dose/Duration

The CDC recommends a single booster immunization at 5+ months for the Pfizer-BioNTech and Moderna vaccines and at least 2 months for the Janssen/Johnson & Johnson vaccine (CDC, 2022)(Mbaeyi et al., 2021).

Indications for discontinuation

N/A for single-administration series. A second immunization is not recommended for those with significant and/or serious adverse effects with the first administration of a two-immunization series.

Rationale

Multiple RCTs [337](Polack, 2020, Heath et al., 2021, Mulligan et al., 2020)(Madhi et al., 2021, Goepfert et al., 2021, Keech et al., 2020, Ella et al., 2021, Chappell et al., 2021, Baden et al., 2021, Logunov et al., 2020, Bonelli et al., 2021)(Borobia et al., 2021, Liu et al., 2021, Folegatti et al., 2020, Li et al., 2021, Pan et al., 2021, Al Kaabi et al., 2021, Pu et al., 2021, Ramasamy et al., 2020, Richmond et al., 2021, Sadoff et al., 2021, Stephenson et al., 2021, Zhu et al., 2020, Zhang et al., 2021, Zhang et al., 2021, Wu et al., 2021, Xia et al., 2021, Shu et al., 2021, Tanriover et al., 2021, Walsh et al., 2020, Xia et al., 2020, Shinde et al., 2021, Chu et al., 2021, Che et al., 2020, Yang et al., 2021) and large population-based studies (Thompson et al., 2021, Thompson et al., 2021) have all documented efficacy of the initial vaccinations of ~85-95%. Subsequent large population-based studies all show that booster immunizations raise antibody levels and markedly lower risk of mortality or hospitalization (Arbel et al., 2021)(Mattiuzzi et al., 2022)(Accorsi et al., 2022)(Schmidt et al., 2022)(Dhingra et al., 2022)(Thompson et al., 2022). Evidence also suggests comparable efficacy of prior infection of approximately 90% reductions in hospitalization and death (Abu-Raddad et al., 2021). Current CDC guidance is to not incorporate prior infections in decisions to (re)vaccinate (CDC, 2021), although there are no long term studies to address that question.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: COVID-19 Vaccines, Vaccination, Pfizer, Moderna, Johnson & Johnson, vaccines; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 2,058 articles in PubMed, 95,514 in Scopus, 366 in CINAHL, 124 in Cochrane Library, 36,300 in Google Scholar, and 0 from other sources†. We considered for inclusion 18 from PubMed, 0 from Scopus, 12 from CINAHL, 7 from Cochrane Library, 7 from Google Scholar, and 0 from other sources. Of the 44 articles considered for inclusion, 34 randomized trials and 6 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

5. MASKS AND RESPIRATORS

Mask use was initiated part-way through the COVID-19 pandemic to attempt to control SARS-CoV-2 exposures (65). Masks are easier to use than respirators, and do not require special fitting. Respirators have much higher performance standards, are more challenging to use correctly, and require fit testing to assure protection meets standards. Masks of widely varying quality, including those which are non-commercial and of single-ply material, have been commonly used by the public in attempts to control SARS-CoV-2 exposure. Respirators have been selectively used to attempt to control COVID-19 viral exposures among higher-risk workers or individuals. Masking mandates have been used for control of COVID-19 both in the workplace and in some jurisdictions (e.g., statewide) (66).

Masking for the Prevention of COVID-19 Transmission

Not Recommended

Masking in closed public spaces was used when transmission was theorized to be primarily droplet spread in an attempt to prevent COVID-19 transmission. Masking in some locations has also been maintained irrespective of community transmission rates and changes in understanding of disease transmission. However, masking appears ineffective. In contrast with masks, N95 respirators may be indicated for select populations, such as high-exposure workers and workers with high personal risks (see Respirator recommendation).

Strength of evidence Not Recommended, Insufficient Evidence (I)

Level of confidence Low

Rationale

Most RCTs of masking to protect from respiratory viruses have been negative (see below). Recently in the COVID-19 pandemic, the omicron variant was found and peaked throughout the world in ~2 months and produced nearly identical epidemic curves irrespective of population-based masking requirements. A prior large, cluster-randomized trial found only an 11% reduction in risk of infection, however the trial was 2 SARS-CoV-2 strains ago when the disease was much less transmissible (Abaluck et al., 2022). Since then, omicron has rapidly moved around the world irrespective of masking and/or physical distancing-business openness in countries and/or states. One community-based moderate-quality trial from Denmark found a lack of benefits from mask wearing in addition to other measures in the COVID epidemic (Gupta et al., 2020). One trial of mask use for COVID-19 assessing household transmission failed to find at least 50% reduction in risk and reported that most disease acquisition was thought to be community-based (Bundgaard et al., 2020).

Prior RCTs mostly involve influenza and influenza-like illness (Bartoszek et al., 2020, Li et al., 2020, Ippolito et al., 2020, Coclite et al., 2020, Jefferson et al., 2009, Liang et al., 2020, Long et al., 2020) and show somewhat conflicting results regarding efficacy to reduce risks of infections, particularly with use of respirators; there are more negative (Canini et al., 2010, Simmerman et al., 2011, Suess et al., 2012, MacIntyre et al., 2009, Jacobs et al., 2009) than positive trial results (Aiello et al., 2010, Cowling et al., 2008, Cowling et al., 2009). Equivalency has been reported between surgical mask use and N95 respirators (Radonovich et al., 2019, Loeb et al., 2009), although experimental evidence suggests superiority of respirators to reduce droplet and aerosols (Wilson et al., 2020, Darby et al., 2021, Asadi et al., 2020). Weak evidence suggests masking may be effective and that N95 respirator use may be superior to mask use in healthcare settings (MacIntyre et al., 2020, MacIntyre et al., 2017, MacIntyre et al., 2014, Chou et al., 2020). All of the epidemiological data have the benefits of being real-world data, but weaknesses include unclear compliance and masking techniques (Kolewe et al., 2020). Respirators performed better than masks in simulation studies (Noti et al., 2012); however, a simulation of SARS-CoV-2 found incomplete protection from masks and N95 respirators (Ueki et al., 2020).

Experimental data on filtering were as follows: N95 respirators, 99%; medical masks, 59%; 3-ply cotton, 51% vs. 47%; double-gaiter, 60%; face shield, 2% (Lindsley et al., 2021, Godoy et al., 2020). Surgical and cloth mask efficacies vary widely (Mueller et al., 2020).

Randomized trial data with more remote and less transmissible strains showed only 11% reduction in risk and subsequent data suggest a lack of significant efficacy and inability to either alter the rate of, or shape of epidemic curves. Thus, masking is not recommended. Although there are no

significant quality data for COVID-19, N95 respirator use may be selectively recommended to attempt to protect those with significant immunosuppressed states and the immediate contacts with such individuals (see respirators).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Mask; Scarf; Bandana; reusability, reusable, reusable cloth, clothing, clothes, clothing, textiles, clothed Mask; Standard surgical mask; N-95; face shield; N95Respirators; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 42 articles in PubMed, 175 in Scopus, 172 in CINAHL, 9 in Cochrane Library, 1,523 in Google Scholar, and 0 from other sources†. We considered for inclusion 0 from PubMed, 12 from Scopus, 0 from CINAHL, 1 from Cochrane Library, 20 from Google Scholar, and 0 from other sources. Of the 32 articles considered for inclusion, 1 randomized trial and 30 systematic reviews, and 1 background information, met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

N95 Respirators for the Prevention of COVID-19 Transmission

Sometimes Recommended

Masking in closed public spaces was used when transmission was theorized to be primarily droplet spread in an attempt to prevent COVID-19 transmission. Masking in some locations has also been maintained irrespective of community transmission rates. However, masking appears ineffective (see Masking recommendation). In contrast with masks, N95 respirators may be indicated for select populations (e.g., high-exposure workers, workers with high personal risks).

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Low

Rationale

Most RCTs of masking to protect from respiratory viruses have been negative (see below). Recently in the COVID-19 pandemic, the omicron variant was found and peaked throughout the world in ~2 months and produced nearly identical epidemic curves irrespective of population-based masking requirements. A prior large, cluster-randomized trial found only an 11% reduction in risk of infection, however the trial was 2 SARS-CoV-2 strains ago when the disease was much less transmissible. Since

then, omicron has rapidly moved around the world irrespective of masking in countries and/or states. One community-based moderate-quality trial from Denmark found a lack of benefits from mask wearing in addition to other measures in the COVID epidemic. One trial of mask use for COVID-19 assessing household transmission failed to find at least 50% reduction in risk and reported that most disease acquisition was thought to be community-based (Bundgaard et al., 2020). A low-quality large cluster randomized trial in Bangladesh found an 11% reduction in risk of infection (Abaluck et al., 2022). Prior RCTs mostly involve influenza and influenza-like illness (Li et al., 2020, Ippolito et al., 2020, Coclite et al., 2020, Jefferson et al., 2009, Liang et al., 2020, Long et al., 2020) and show somewhat conflicting results regarding efficacy to reduce risks of infections, particularly with use of respirators; there are more negative (Canini et al., 2010, Simmerman et al., 2011, Suess et al., 2012, MacIntyre et al., 2009, Jacobs et al., 2009) than positive trial results (Aiello et al., 2010, Cowling et al., 2008, Cowling et al., 2009). Equivalency has been reported between surgical mask use and N95 respirators (Radonovich et al., 2019, Loeb et al., 2009), although experimental evidence suggests superiority of respirators to reduce droplet and aerosols (Darby et al., 2021, Asadi et al., 2020). Weak evidence suggests masking may be effective and that N95 respirator use may be superior to mask use in healthcare settings (MacIntyre et al., 2020, MacIntyre et al., 2017, MacIntyre et al., 2014, Chou et al., 2020). All of the epidemiological data have the benefits of being real-world data, but weaknesses include unclear compliance and masking techniques (Kolewe et al., 2020). Respirators performed better than masks in simulation studies (Noti et al., 2012); however, a simulation of SARS-CoV-2 found incomplete protection from masks and N95 respirators (Ueki et al., 2020). Data on filtering were as follows: N95 respirators, 99%; medical masks, 59%; 3-ply cotton, 51% vs. 47%; double-gaiter, 60%; face shield, 2% (Lindsley et al., 2021, Godoy et al., 2020). Surgical and cloth mask efficacies vary widely (Mueller et al., 2020). Randomized trial data with more remote and less transmissible strains showed only 11% reduction in risk and subsequent data suggest a lack of significant efficacy and inability to either alter the rate of or shape of epidemic curves. Thus, masking is not recommended. Although there are no significant quality data for COVID-19, N95 respirator use may be selectively recommended to attempt to protect those with significant immunosuppressed states and the immediate contacts with such individuals.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to February 2020 using the following terms: Mask, bandana, scarf, reusable cloth mask, standard surgical mask, N-95, face shield; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 237 articles in PubMed, 70 in Scopus, 71 in CINAHL, 117 in Cochrane Library, 2882 in Google Scholar, and 3 from other sources[†]. We considered for inclusion 29 from PubMed, 4 from Scopus, 2 from CINAHL, 1 from Cochrane Library, 44 from Google Scholar, and 3 from other sources. Of the 82 articles considered for inclusion, 23 randomized trials and 40 systematic reviews met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

6. LOCKDOWNS AND SHUTDOWNS

Restrictions on businesses, schools, and public gatherings have been used in attempts to control the COVID-19 pandemic, including limitations on travel, large gatherings, in-person schools, restaurants, bars, and non-essential businesses. Even under the strictest shelter-in-place jurisdictions in the United States, the variants spread regardless. Most individuals were permitted to continue to visit grocery stores despite lockdowns, which may have provided a means for continuing community spread despite masking requirements (e.g., walking through others' vapor trails while using inadequate protection for a highly contagious aerosol).

Studies are beginning to be published concerning the efficacy of lockdowns. Most studies have reported some reductions in COVID-19 transmission after the implementation of a lockdown (67,68), although it has been reported that lockdowns were not effective in Europe (67). An ecological study suggested greater spread where restaurant dining was allowed (69). One analysis of multiple countries found non-significant small reductions in COVID-19 case rates in most countries, which was not felt to be outweighed by the costs (67). Reports have questioned the cost-benefit efficacy of lockdowns (70,71)(5). Adverse mental health effects have been reported (72,73,74,75,76). The subject of lockdowns requires considerably greater research, especially as future surges attributed to variants seem likely; the re-implementation of such lockdown policies may necessitate a stronger evidence base.

The overall available evidence does not support lockdown measures as they:

- appear largely ineffective to slow the spread and impact of the SARS-CoV-2 virus and/or any of its variants during this pandemic;
- are likely to have contributed to increased problems with anxiety, depression, other mental health disorders and drug overdoses;
- were associated with job loss and unemployment; and
- were generally accompanied by cessation of in-classroom teaching at schools, which were also ineffective and harmful.

Lockdown measures are not recommended (5).

7. DIAGNOSTIC APPROACH

7.1. LABORATORY TESTS

COVID-19 has a widely varying clinical presentation. Depending on the extent of infection and the organ systems affected, any or all of the following may be found (77,78):

- lymphopenia (a fairly unique and characteristic finding)
- elevated liver enzymes
- elevated lactate dehydrogenase (LDH)
- elevated direct bilirubin
- elevated pancreatic enzymes

- elevated prothrombin time (PT)
- elevated troponin
- elevated creatine phosphokinase (CPK)
- elevated inflammatory markers (e.g., C-reactive protein [CRP], ferritin)
- elevated D-dimer
- elevated fibrinogen
- elevated creatinine
- elevated blood urea nitrogen
- hypoxemia

A risk prediction model has been developed to predict the development of severe disease (79). The 10 variables included in the model are:

- chest radiographic abnormality (odds ratio [OR]: 3.39),
- age (OR: 1.03),
- hemoptysis (OR: 4.53),
- dyspnea (OR: 1.88),
- unconsciousness (OR: 4.71),
- number of comorbidities (OR: 1.60),
- cancer history (OR: 4.07),
- neutrophil-to-lymphocyte ratio (OR: 1.06),
- lactate dehydrogenase (OR: 1.002), and
- direct bilirubin (OR: 1.15).

A free online risk calculator is available (80).

Decreases in creatinine kinase (CK) and LDH have been associated with increased COVID-19 viral clearance in a secondary analysis of hospitalized patients treated with varying antiviral and other medications (IFN- α + lopinavir/ritonavir \pm ribavirin) (81).

7.2. DIAGNOSTIC TESTING

Three main types of diagnostic tests are used for COVID-19:

- polymerase chain reaction (PCR)-based testing (aka, nucleic acid amplification test), typically using swabs (595);
- antigen testing, and

- antibody testing of blood serum.

PCR testing is considered to be diagnostic of the infection because it detects the actual virus or viral particles. Antigen tests have been approved by the U.S. Food and Drug Administration (FDA) and are also considered diagnostic (596). Antibody testing detects prior infection. All types of testing have had limitations in specificity and sensitivity. A difference in performance over time since symptom onset has been reported (597). There are concerns with the accuracy of testing to detect newer variants, especially with false negative tests among those with early infections or mild symptoms and lower viral loads (REF).

Saliva testing for SARS-CoV-2 detection is also available, which is appealing for ease of collection. Pooled saliva testing has been used in employed populations (598). One study detected higher SARS-CoV-2 titers in saliva compared to nasopharyngeal swabs, with less longitudinal variability (599). A meta-analysis found similar sensitivities and lower costs for saliva sampling compared with nasopharyngeal swabs (600). Thus, saliva testing is increasingly replacing swabs to provide near universal sampling coverage for both symptomatic and asymptomatic patients (601). Yet, accuracy for omicron is similarly unclear.

Test results, when accurate, may only indicate the presence or absence of infection at the time of the test; thus, the frequency of testing, and which methods to use, are debatable. In university settings, routine surveillance testing of representative subpopulations of students is recommended, with more frequent testing of higher-risk groups such as athletes. More frequent testing with less sensitive (and often cheaper) tests that are capable of detecting infectious virus (rather than any virus) will shortly become available and are recommended (602).

PCR TESTING

PCR samples and testing techniques amplify viral particles to identify relatively small amounts of virus, with the nucleocapsid antigen test being the most sensitive for detecting early infection (603). Because they also amplify viral fragments, they can show recent infection among those who are still clearing the viral particles, up to weeks after infection; thus, they may not reflect active viral shedding and/or infectiousness. These tests can indicate the RNA debris of coronavirus and may reflect non-viable virus remnants, thus use of PCR for travel restrictions is questionable.

Importantly, the risks of false-negative and false-positive test results change as a pandemic progresses. For example, as disease becomes more common, individuals who present with symptoms but test negative are increasingly likely to represent false-negatives irrespective of testing accuracy. Thus, once an epidemic disease becomes highly pervasive and there is not a common competing cause of similar symptoms, diagnostic testing is often unnecessary for typical cases because it does not materially alter the post-test probability. At an epidemic's peak, the testing of unusual cases is ideally performed with highly accurate tests, as such cases may represent unusual presentations of COVID-19 infection that should be distinguished from non-COVID-19 causes. Because the SARS-CoV-2 virus causes such a wide spectrum of disease, from asymptomatic illness to life-threatening infection, along with the possibility of other co-circulating respiratory viruses at various times (e.g., influenza), the issue of accurate diagnostics for SARS-CoV-2 continues to be important. The ability to widely perform COVID-19 testing is of particular importance during times of anticipated epidemic waves and regarding accuracy in detecting future variants.

Most of the limited evidence suggests that nasopharyngeal and oropharyngeal samples are comparable for the first week, but then the nasopharyngeal sample becomes more sensitive (604,605):

- From days 0–7, oropharyngeal and nasopharyngeal sensitivities are 61/60% and 72/73% for mild/severe disease, respectively.

- On days 8–14, oropharyngeal and nasopharyngeal sensitivities are approximately 30/50% and 54/72% for mild/severe disease, respectively (606).

-

PCR Testing for the Diagnosis of COVID-19

Recommended

PCR testing is recommended for the diagnosis of COVID-19. Testing should be performed either at the time of COVID-19-like symptom onset, or within several days of the onset of symptoms consistent with a COVID-19 infection. Testing without experienced medical judgment is ill-advised given that the risk of false-negative tests are 20–67% (371). Thus, there is a strong indication to presumptively treat cases who test negative, which requires experienced medical judgment. Repeat testing may be indicated for those with a negative test but a high index of suspicion.

PCR testing is also recommended for inpatient and outpatient preoperative assessments. Preoperative tests must be ordered sufficiently ahead of surgery such that the results are received in time to address/respond to the results (generally 72–96 hours before surgery).

Strength of evidence Recommended, Insufficient Evidence (I)

ANTIGEN TESTING

Antigen tests detect viral proteins either on or within the virus. These have been FDA-approved and are considered diagnostic (596). Antigen testing is growing in popularity as its main strength is rapid test results, which are provided in minutes compared with up to several days for PCR tests. One report has suggested lower sensitivity comparing four different rapid antigen tests (biotical, Panbio, Healgen, Roche) with one automated antigen detection test (VITROS) for the diagnosis of COVID-19; yet, the rapid results were felt to still confer an advantage for positive test results to detect infectious patients (607).

Antigen Testing for the Diagnosis of COVID-19

Recommended

Antigen testing is recommended for the diagnosis of COVID-19. Testing should be performed either at the time of COVID-19-like symptom onset, or within several days of the onset of symptoms consistent with a COVID-19 infection. Antigen testing has not been validated for asymptomatic persons. However, the sensitivity among symptomatic persons is estimated to be approximately 80%. Thus, testing without experienced medical judgment is ill-advised (Domeracki et al., 2020), given the risks of false-negative tests. There is a strong indication to presumptively treat cases who test negative, which requires experienced medical judgment. Repeat testing may be indicated for those with a negative test but a high index of suspicion.

Antigen testing is also recommended for inpatient and outpatient preoperative assessments. Preoperative tests must be ordered sufficiently ahead of surgery such that the results are received in time to address/respond to the results (generally 72–96 hours before surgery). Preoperative tests may be needed both for those without any history of symptoms, as well as for those with prior infections, to assure the person is no longer infectious.

ANTIBODY TESTING

Antibody testing detects the body's humoral response to the virus (608,609,610,611,612,613). Most antibody tests detect IgG, although some tests attempt to also detect IgM or IgA. The median IgM seroconversion is 11–13-days (or 5–7-days after symptoms onset), while the median seroconversion for IgG is 14-days (or 8-days after symptoms onset), although IgM may wane after 2- to 3-weeks, and IgG persists for a far longer period of time (614). A positive antibody test does not exclude the potential for the patient being infectious with COVID-19. Antibody tests are in early stages of deployment and reported reliability varies widely (610,611,612). Because there is no reference standard and widespread testing of large populations have not been reported, the determination of test accuracy, sensitivity, and specificity remain problematic. In addition, the timing of the antibody testing is critical to accurate detection: testing too soon after infection onset, or too late after infection resolution, can further increase risks of negative results.

It has been aspirational that immune status testing (IgG, IgM) would eventually be the most important test for population-based risk assessments, such as herd immunity. This still requires considerable research, including large-scale determinations of sensitivity, specificity, reliability, timing, persistence of the immunoglobulins, and whether the immunoglobulin status identified by testing will be associated with true immunity (615). Preliminary evidence includes a large population-based Spanish study suggesting a 87.6–91.8% seroprevalence rate among those who had PCR confirmation of infection; yet, individuals meeting a case definition of anosmia or at least 3 relevant symptoms had a seroprevalence rate of only 15.3–19.3% (616). A large-scale hospital-based study found a sensitivity of 97.6% and 98.8% specificity when performed 14-days or later after symptoms onset; the immunoglobulins levels were correlated with worse disease, and were detectable in those with negative PCR tests but clinical suspicion of infection (617). Others have correlated titers with disease severity (611). An added challenge is that while 1.24% of a community's 5,882 samples showed antibody reactivity to receptor binding domain, 18% of the samples failed to neutralize the SARS-CoV-2 virus (618).

Evidence also suggests immunoglobulins may not be measurable over time (619). Still, other studies suggest laboratory tests assessing T-cell responses remain robust for some time, even among those with no detectable immunoglobulins and/or those who had mild disease (620,621). Hence, a lack of measurable immunoglobulins may not indicate lack of immunity. If these lines of research remain viable, then it is theoretically possible for immunoglobulin testing, perhaps combined with history, to help designate workers who may more safely interact with the public. If proven, antibody testing may be used to assure a workplace that a previously infected worker is safe to return to work (i.e., that they are not actively infected and unlikely to be shedding virus). Unfortunately, the currently available antibody tests have yet to be sufficiently validated on a widespread basis, and inaccuracies are increasingly reported (622,623). Once these problems are addressed, it is anticipated that antibody testing may become widespread in many workplaces and other populations of concern (e.g., nursing homes, mission-critical workers, irreplaceable workers, dispatch centers, C-suite executives).

Immune status determination, if proven, may be of major importance for workplace populations in many, if not all, sectors. It may be complementary with vaccination, particularly if the virus continues to circulate and cause disease. Workforces with the greatest needs for immune status testing include those with isolated populations, increased risk of transmission to vulnerable populations, high worker densities, and/or distance from and lack of access to appropriate healthcare (e.g., oil platform drilling, commercial maritime, cruise lines, overseas workforces, airlines, rail, trucking, mining).

Antibody Testing for Assessing COVID-19 Immune Status

Sometimes Recommended

Antibody testing is selectively recommended for assessing immune status regarding the potential for COVID-19. These tests should be interpreted by experienced medical and/or public health professional(s) who are thoroughly knowledgeable about numerous factors, including the specific test, its reported performance (e.g., sensitivity, specificity), the prevalence of COVID-19 in the specific community, principles of testing, Bayes' theorem, and assessment of pre-test probability and post-test odds. In general and at this point, antibody testing should be limited to only mission-critical workers and special populations. As the experience with these tests improves, the populations assessed may markedly expand. As a general statement, a person who has recovered from COVID-19, has a duration of at least 10-days since first symptoms, and has demonstrated antibodies would not be infectious or capable of transmitting infection and scientifically would no longer have to wear a mask or participate in mitigation procedures.

Specific examples where serology might be helpful include the following:

- patients with symptoms consistent with COVID-19 of more than 1-week in duration, for whom PCR testing has been negative and no alternative diagnosis has been found. For these cases, a positive IgG serology would be diagnostic. A negative serology could be repeated at >2-weeks from symptom onset and repeat negative testing would then effectively rule out COVID-19.
- patients with initial negative PCR and serology at <2-weeks after symptom onset but who remain symptomatic beyond 2-weeks without an alternative diagnosis. Repeat serology testing documenting seroconversion would be diagnostic, whereas failure to seroconvert would help to rule out COVID-19.
- symptomatic, febrile, PCR-positive patients with an unknown time since infection where presence of antibodies might help in choice of therapeutic modalities (e.g., antivirals and/or convalescent serum before antibodies arise).

Strength of evidence Recommended, Insufficient Evidence (I)

7.3. IMAGING

Radiographs for the Diagnosis of COVID-19

Recommended

Radiographs are recommended as part of the diagnostic evaluation of COVID-19. Although radiographs are usually abnormal for individuals with pulmonary involvement, radiography in general should not be used as a stand-alone screening tool for COVID-19. X-ray abnormalities peak at 10–12 days after onset of symptoms (Rodriguez-Morales et al., 2020, Wong et al., 2020). One series reported that chest radiographs most commonly show either consolidation (47%) or ground-glass abnormalities (33%). The same series noted that 41% were peripheral, 50% were lower distribution, and 50% were bilateral (Wong et al., 2020).

Strength of evidence Recommended, Insufficient Evidence (I)

Computed Tomography for the Diagnosis of COVID-19

Recommended

Computed tomography (CT) scans are recommended for the diagnostic evaluation of COVID-19. CT is commonly performed (Udugama et al., 2020, Sun, 2020) and shows patchy infiltrates and ground-glass opacities (Chan et al., 2020, Li et al., 2020, Li et al., 2020, Iwasawa et al., 2020, Liu et al., 2020). One report has suggested comparable sensitivity for screening in the emergency department for CT compared with ultrasound (Lieveld et al., 2020). One series reported 72% of cases with ground-glass appearance, 12% with consolidation, 12% with crazy paving patterns, 37% with interlobular thickening, 56% with adjacent pleural thickening, and 61% with linear opacities (Xu et al., 2020).

Strength of evidence Recommended, Insufficient Evidence (I)

8. TREATMENT RECOMMENDATIONS

8.1. OVERVIEW

Treatment is increasingly guided by randomized controlled trials (RCTs), yet it continues to evolve as data are published. Many additional studies are underway. However, variants are providing a significant challenge as they provide uncertainty regarding the translation of prior RCTs to application for treatment of the recent variant(s). There is a major need for RCT(s) that include comparative trials for therapeutics, especially for the very early phase of infection (24-48 hours after symptom onset) when use of effective treatments may obviate the need for hospitalization and prevent deaths. The design of at least one large, multi-armed RCT, including the available therapeutics with suggested efficacy (see below), is needed now to allow for enrollment at the beginning of the next variant surge. Results will determine which of several options is most effective, and used to calculate comparative efficacy. These trials are especially needed to prepare to treat the immunosuppressed populations, which may incur the worst outcomes in the subsequent variant waves.

There are numerous treatment guidelines available; although these guidelines tend to have similar recommendations, there are some differences regarding individual treatments (624,625,626,627,628,629,630,631)(632)(633). The FDA continues to provide unprecedented flexibility to accelerate the development of new drugs and testing. No treatment is yet unequivocally indicated for asymptomatic cases; there is an NIH recommendation for prophylactic treatment with tixagevimab plus cilgavimab, although it is unclear if this is an effective approach for the latest variant(s) (634).

The four main classes of interventions with evidence of efficacy for more serious infections are antiviral treatments, cytokine storm-reducing and/or immunomodulating agents, anticoagulants, and ventilatory support (both non-invasive and invasive).

Many medications and agents, some with evidence of efficacy and some without, have been used for treatment, including the following: ACE inhibitors, Adalimumab, anticoagulants, bamlanivimab, bebtelovimab, casirivimab/imdevimab, COVID-19 convalescent plasma, famotidine, fluvoxamine, low-molecular-weight heparin, molnupiravir, monoclonal antibodies, azithromycin, baloxavir, baricitinib, chloroquine, colchicine, favipiravir, glucocorticosteroids, hydroxychloroquine, immunoglobulin, interferons, ivermectin, lopinavir/ritonavir, nitric oxide, paxlovid, ribavirin, remdesivir, sarilumab, siltuximab, statins, Sotrovimab, thrombolytics, tocilizumab, zinc

(635,636,637,638), vitamin C (639), and vitamin D (640,641,642,643). Many of these treatments have no quality evidence of efficacy, and some have evidence of a lack of efficacy. There is no clear evidence of lower risk of mortality with statin use (644). Vitamin D levels have been strongly correlated with COVID-19 disease severity (640,642,643); for example, individuals with low vitamin D levels were reported to have an approximate 8-fold greater risk of a severe outcome and 20-fold greater risk of a critical outcome (640). Still, while some studies are suggestive, strong evidence that vitamin D supplementation affects mortality risk has not been published. In general, considering public health, having a vitamin D level of at least 20-30 ng/mL is reasonable (238) (237)(645))

Glucocorticosteroids have been shown in multiple quality trials to reduce mortality (646,647,648). Data also suggest that low-molecular-weight heparin likely reduces mortality and shortens ICU stays (649).

If individuals develop more severe symptoms or have complications (e.g., ARDS or respiratory failure), they are primarily treated with non-invasive ventilatory support measures, glucocorticosteroids, anti-cytokine storm agents, mechanical ventilation (including prone positioning), other respiratory support measures, and prophylaxis for deep vein thrombosis, including low-molecular-weight heparins (650,651,652). Evaluations should include exclusion of other causes (e.g., influenza). The efficacy of glucocorticoids appears to be related to the stage of the COVID-19 infection. Glucocorticosteroids used early in the time course of infection do not appear to improve outcomes, and in theory could potentially allow viral replication to increase and foster the development of other infections.

Multiple agents have been studied to attempt to suppress the purported cytokine storm; most of the trials are centered around interleukin-6 (IL-6) (653). Yet, most quality data on IL-6 receptor antagonists have been negative. There is ongoing controversy regarding a cytokine storm in relation to ARDS caused by COVID-19 [457]. There are many cytokines believed to be involved in the cytokine release syndrome (IL-2, IL-7, G-CSF, IFN- γ , inducible protein 10, MIP 1- β , TNF- α).

Antiviral medications may have minimal to no role in advanced pneumonia or ARDS (654), particularly as viral replication appears to peak at or about the time of symptoms onset. However, antiviral therapies are showing increasing promise to lessen the severity of the disease among outpatients who are treated early in the disease. Two therapies targeting this window were previously approved by FDA under emergency use authorization: bamlanivimab and casirivimab/imdevimab; however, the omicron variant appears to evade these therapies. Data on hydroxychloroquine (HCQ) suggest efficacy early in the symptomatic phase, while there is also clear evidence of inefficacy for later stage use (655). Unfortunately, there are few studies and a high need for assessing the efficacy of antiviral medications within the first 1–2 days of symptom onset (656); there appear to be parallels with influenza medications used within 48 hours of symptoms onset.

Projecting this pandemic forward through expected waves of variant surges, it is expected that monoclonal antibodies will potentially become rapidly partially or wholly ineffective, especially those targeting the spike protein when a new variant arises to thwart that antibody. Antiviral medications that target the replication processes of the virus may also become ineffective. However, based on other viruses (e.g., influenza), the rate of such resistance is anticipated to occur at a far slower rate.

Potential hierarchical approaches for the treatment of COVID-19 are as follows:

- **Outpatient**

Mild: No treatment unless high risk for progression to severe disease

Moderate/severe:

1. Sotrovimab
2. Molnupiravir
3. Paxlovid (Nirmatrelvir/Ritonovir)
4. Fluvoxamine
5. Ivermectin
6. Hydroxychloroquine for 5 days
7. Vitamin D, Zinc

- **Inpatient moderate**

1. Glucocorticosteroids
2. Low-molecular-weight heparin/unfractionated heparin
3. Remdesivir
4. Oxygen supplementation
5. Vitamin D

- **Inpatient severe/critical**

1. Glucocorticosteroids
2. Low-molecular-weight heparin/unfractionated heparin
3. Remdesivir
4. Oxygen supplementation
5. Prone positioning (due to shunting) and/or non-invasive ventilation (NIV)
6. Mechanical ventilation, prone
7. Extracorporeal membrane oxygenation (ECMO)
8. Vitamin D

Additional treatments are being studied, including antiviral mouthwashes and nasal sprays (657,658)(659,660),(661,662)(663) melatonin (664,665,666,667,668,669,670), and others.

Mental health issues are increasingly recognized as problematic, both among those infected as well as those otherwise impacted by the epidemic but not infected. Several references are available that include evidence of an epidemic of depression (50% increased), suicidal ideation, anxiety, post-traumatic stress disorder (PTSD), substance use, divorce (30% increased), and violence (77,671,672,673,674,675,676,677,678). An association between adverse mental health and financial concerns has been noted (679). Recent data from a small autopsy study suggest findings similar to Alzheimer disease in the brains of those deceased from COVID (680).

8.2. HYDROXYCHLOROQUINE OR CHLOROQUINE

Hydroxychloroquine has been used for the treatment of COVID-19 (635,638,654,681,682,683,684,685,686,687,688,689,690,691,692,693,694,695,696,697,698,699,700,701,702,703,704,705,706,707,708,709,710,711,712,713,714,715,716,717,718,719,97). There also are many in vitro studies suggesting antiviral activity (720,721,722,723,724,725,726,727,728). Chloroquine has been used for the treatment of COVID-19 (729). Hydroxychloroquine has been used for prophylaxis for COVID-19, most typically among healthcare workers (717,730).

Hydroxychloroquine for Treatment of COVID-19 - Use in First 3 Days of Symptoms

Recommended

Hydroxychloroquine (HCQ) is recommended for use in the first 3 days of symptom onset.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Indicated for early symptom onset, ideally in the first 1–3 days during the COVID-19 phase with viral replication. Not indicated for late symptoms, especially days 5 or later. Generally for moderate to severely affected patients with COVID-19 and would include zinc supplementation. Use in mild cases could be justified, especially for a patient with multiple comorbidities (e.g., pre-diabetes, diabetes, cardiovascular disease, COPD) and thus risk of progression.

Benefits

Meta-analysis evidence of a 24% reduction in composite risk of COVID-19 infection, hospitalization, and death (Ladapo et al., 2020). Earlier clearance of pneumonia on CT scan (CHEN et al., 2020).

Harms

Negligible for most patients undergoing short-course use (Lofgren et al., 2020). Gastrointestinal symptoms occur above rates of placebo. Prior concerns about prolonged corrected QT intervals, and thus arrhythmias (Magagnoli et al., 2020, Chorin et al., 2020), have been largely resolved among previously healthy patients without risks for arrhythmias who are given HCQ at typical doses. ECG monitoring may be indicated for patients with underlying cardiovascular disease, history of prolonged QT, unexplained syncope, family history of premature sudden cardiac death, electrolyte

abnormalities, renal insufficiency, and use of other drugs reported to prolong QT intervals, including when there is planned adjunctive use with azithromycin. Renal insufficiency also may increase toxicity risks. Retinopathy appears highly unlikely with these short courses, as it has been reported at levels of >100-fold greater cumulative doses (Marmor et al., 2016).

Frequency/Dose/Duration

Multiple regimens have been used. There is both a mechanistic rationale for the concomitant use of zinc to inhibit viral replication and pre-post interventional clinical evidence of efficacy for the adjunctive use of zinc (Carlucci et al., 2020). The following are the most common regimens, the first of which was used in the one quality RCT:

- Hydroxychloroquine 400mg BID x 1 day, then 200mg BID for 4 days (Yao et al., 2020).
- Hydroxychloroquine 400mg BID x 1 day, then 400mg QD for 4 day.
- Hydroxychloroquine 200mg BID x 5 days (CHEN et al., 2020)
- Hydroxychloroquine 200mg TID x 10 days (Gautret et al., 2020)
- Hydroxychloroquine 200mg TID x 10 days plus azithromycin 500mg x 1 day then 250mg QD x 4 days (Gautret et al., 2020)
- Hydroxychloroquine 600mg BID x 1 day, then 400mg QD for 4 days

Because the half-life of these medications is long, a loading dose for the first day or two may be preferable.

Rationale

Late use of HCQ (>3 days) has been assessed in many quality RCTs among hospitalized and/or ICU patients. These trials consistently show late use of HCQ does not improve clinical outcomes, including mortality (Cavalcanti et al., 2020, Ulrich et al., 2020, Lyngbakken et al., 2020) (Self et al., 2020, Schwartz et al., 2021) (Johnston et al., 2021)(Reis et al., 2021, Axfors et al., 2021)(Beltran Gonzalez et al., 2022) or viral clearance (Barratt-Due et al., 2021). One trial of HCQ with and without AZT found no efficacy although there was a trend towards faster viral clearance with HCQ (Johnston et al., 2021). Another comparative trial found comparable (in)efficacy of HCQ and AZT (Brown et al., 2021). One trial with mostly late treatment patients found lack of benefit with Favipiravir with and without HCQ (Bosaeed et al., 2021). Two open-label trials reported worse outcomes with either HCQ or CQ (Réa-Neto et al., 2021) or HCQ (Arabi et al., 2021). There are observational studies suggesting more than doubling survival with combined HCQ/AZT among hospitalized patients on ventilators (Smith et al., 2021). Because there is reasonably consistent high- and moderate-quality evidence that HCQ is ineffective as a solitary intervention in COVID-19 patients treated late after the viral replication phase has largely ceased, the use of HCQ in that timeframe is not recommended. Early use of HCQ has been assessed in multiple studies that range from pre-diagnosis to within a few days of symptom onset (Ladapo et al., 2020). These trials are naturally individually underpowered for severe outcomes such as mortality as they tend to include younger, healthier patients. A meta-analysis of 5 RCTs that analyzed 5,577 patients found that all studies trended towards efficacy and the combined data showed a statistically significant 24% reduction in composite risk of infection, hospitalization, and death (Ladapo et al., 2020). One other RCT trended towards reduced hospitalization and time to symptom resolution (Mitjà et al., 2021) and another showed earlier resolution of cough (Rodrigues et al., 2021). A nationwide cohort study in the Netherlands found evidence of efficacy of hydroxychloroquine for reducing risk of transfer to an ICU by 53% compared

with no treatment, but there was no similar effect for chloroquine (Lammers et al., 2020). A study of 1,274 outpatients in a propensity-matched cohort from New Jersey found a 31.2% reduced risk of hospitalization (Ip et al., 2021). An observational study found an 84% ($p<0.001$) reduction in risk of hospitalization and risk of death was also 80% ($p=0.12$) reduced among those treated early with HCQ, zinc, and AZT (Derwand et al., 2020).

One early-use trial found non-significant reductions, with 20% being symptomatic at 14 days and a 60% reduced risk of death (Skipper et al., 2020). Another trial of HCQ used within 4 days of high-risk exposure found a 17% reduced risk of subsequent infection (Boulware et al., 2020). Another trial of once-weekly or twice-weekly HCQ as pre-exposure prophylaxis among HCWs found a non-significant 26–28% reduced risk of infection (Rajasingham et al., 2020). Because there is quality evidence of efficacy for the early use of HCQ, it is recommended for these select patients.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2020 to October 2021 using the following terms:

Hydroxychloroquine; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 153 articles in PubMed, 25,203 in Scopus, 1,779 in CINAHL, 18 in Cochrane Library, 10,300 in Google Scholar, and 0 from other sources†. We considered for inclusion 11 from PubMed, 1 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 17 from Google Scholar, and 0 from other sources. Of the 74 articles considered for inclusion, 7 randomized trials, 2 non-randomized trials, 5 case series, 11 retrospective studies, and 5 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens, then the remaining articles are not reviewed due to a lack of relevancy.

Hydroxychloroquine for Treatment of COVID-19 - Use Beyond First 3 Days of Symptoms

Not Recommended

Hydroxychloroquine (HCQ) is not recommended for the treatment of patients with COVID-19 after the first 3 days of symptoms (Oxford, 2020).

Strength of evidence Moderately Not Recommended, Evidence (B)

Level of confidence Moderate

Indications

Indicated for early symptom onset, ideally in the first 1–3 days during the COVID-19 phase with viral replication. Not indicated for late symptoms, especially days 5 or later. Generally for moderate to

severely affected patients with COVID-19 and would include zinc supplementation. Use in mild cases could be justified, especially for a patient with multiple comorbidities (e.g., pre-diabetes, diabetes, cardiovascular disease, COPD) and thus risk of progression.

Benefits

Meta-analysis evidence of a 24% reduction in composite risk of COVID-19 infection, hospitalization, and death (Ladapo et al., 2020). Earlier clearance of pneumonia on CT scan (CHEN et al., 2020).

Harms

Negligible for most patients undergoing short-course use (Lofgren et al., 2020). Gastrointestinal symptoms occur above rates of placebo. Prior concerns about prolonged corrected QT intervals, and thus arrhythmias (Magagnoli et al., 2020, Chorin et al., 2020), have been largely resolved among previously healthy patients without risks for arrhythmias who are given HCQ at typical doses. ECG monitoring may be indicated for patients with underlying cardiovascular disease, history of prolonged QT, unexplained syncope, family history of premature sudden cardiac death, electrolyte abnormalities, renal insufficiency, and use of other drugs reported to prolong QT intervals, including when there is planned adjunctive use with azithromycin. Renal insufficiency also may increase toxicity risks. Retinopathy appears highly unlikely with these short courses, as it has been reported at levels of >100-fold greater cumulative doses (Marmor et al., 2016).

Frequency/Dose/Duration

Multiple regimens have been used. There is both a mechanistic rationale for the concomitant use of zinc to inhibit viral replication and pre-post interventional clinical evidence of efficacy for the adjunctive use of zinc (Carlucci et al., 2020). The following are the most common regimens, the first of which was used in the one quality RCT:

- Hydroxychloroquine 400mg BID x 1 day, then 200mg BID for 4 days (Yao et al., 2020).
- Hydroxychloroquine 400mg BID x 1 day, then 400mg QD for 4 days.
- Hydroxychloroquine 200mg BID x 5 days (CHEN et al., 2020)
- Hydroxychloroquine 200mg TID x 10 days (Gautret et al., 2020)
- Hydroxychloroquine 200mg TID x 10 days plus azithromycin 500mg x 1 day then 250mg QD x 4 days (Gautret et al., 2020)
- Hydroxychloroquine 600mg BID x 1 day, then 400mg QD for 4 days

Because the half-life of these medications is long, a loading dose for the first day or two may be preferable.

Rationale

Late use of HCQ (>3 days) has been assessed in many quality RCTs among hospitalized and/or ICU patients. These trials consistently show late use of HCQ does not improve clinical outcomes, including mortality (Oxford, 2020, Cavalcanti et al., 2020, Ulrich et al., 2020, Lyngbakken et al., 2020, Abd-Elsalam et al., 2020, Horby et al., 2020) (Self et al., 2020, Schwartz et al., 2021) (Johnston et al., 2021)(Axfors et al., 2021, Reis et al., 2021)(Beltran Gonzalez et al., 2022) or viral clearance (Barratt-Due et al., 2021). One trial of HCQ with and without AZT found no efficacy although there was a

trend towards faster viral clearance with HCQ (Johnston et al., 2021). Another comparative trial found comparable (in)efficacy of HCQ and AZT (Brown et al., 2021). One trial with mostly late treatment patients found lack of benefit with Favipiravir with and without HCQ (Bosaeed et al., 2021). Two open-label trials reported worse outcomes with either HCQ or CQ (Réa-Neto et al., 2021) or HCQ (Arabi et al., 2021). There are observational studies suggesting more than doubling survival with combined HCQ/AZT among hospitalized patients on ventilators (Smith et al., 2021). Because there is reasonably consistent high- and moderate-quality evidence that HCQ is ineffective as a solitary intervention in COVID-19 patients treated late after the viral replication phase has largely ceased, the use of HCQ in that timeframe is not recommended.

Early use of HCQ has been assessed in multiple studies that range from pre-diagnosis to within a few days of symptom onset (Ladapo et al., 2020). These trials are naturally individually underpowered for severe outcomes such as mortality as they tend to include younger, healthier patients. A meta-analysis of 5 RCTs that analyzed 5,577 patients found that all studies trended towards efficacy and the combined data showed a statistically significant 24% reduction in composite risk of infection, hospitalization, and death (Ladapo et al., 2020). One other RCT trended towards reduced hospitalization and time to symptom resolution (Mitjà et al., 2021) and another showed earlier resolution of cough (Rodrigues et al., 2021). A nationwide cohort study in the Netherlands found evidence of efficacy of hydroxychloroquine for reducing risk of transfer to an ICU by 53% compared with no treatment, but there was no similar effect for chloroquine (Lammers et al., 2020). A study of 1,274 outpatients in a propensity-matched cohort from New Jersey found a 31.2% reduced risk of hospitalization (Ip et al., 2021). An observational study found an 84% ($p < 0.001$) reduction in risk of hospitalization and risk of death was also 80% ($p = 0.12$) reduced among those treated early with HCQ, zinc, and AZT (Derwand et al., 2020).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to November 2020 using the following terms:

Hydroxychloroquine; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 528 articles in PubMed, 741 in Scopus, 137 in CINAHL, 425 in Cochrane Library, 9,380 in Google Scholar, and 38 from other sources[†]. We considered for inclusion 24 from PubMed, 6 from Scopus, 1 from CINAHL, 2 from Cochrane Library, 6 from Google Scholar, and 35 from other sources. Of the 74 articles considered for inclusion, 7 randomized trials, 2 non-randomized trials, 5 case series, 11 retrospective studies, and 5 systematic reviews met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens, then the remaining articles are not reviewed due to a lack of relevancy.

Chloroquine for Treatment of COVID-19 - Use in First 3 Days of Symptoms

No Recommendation

There is no recommendation for or against the use of chloroquine in the first 3 days of symptoms.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Rationale

Chloroquine is a closely related compound to hydroxychloroquine. There is no RCT-level evidence that chloroquine has different efficacy. There are sparse trials of chloroquine, especially compared with the evidence base for hydroxychloroquine. One population-based cohort study found evidence of efficacy of hydroxychloroquine but not chloroquine (Lammers et al., 2020). Thus, by analogy to hydroxychloroquine, chloroquine is not recommended for treatment of hospitalized COVID-19 patients. See the Hydroxychloroquine Rationale for Recommendation for details.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: **Chloroquine**; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 766 articles in PubMed, 18,961 in Scopus, 33 in CINAHL, 1,279 in Cochrane Library, 9,500 in Google Scholar, and 0 from other sources†. We considered for inclusion 0 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 0 articles considered for inclusion, 0 randomized trials and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Chloroquine for Treatment of COVID-19 - Use Beyond First 3 Days of Symptoms

Not Recommended

Chloroquine is not recommended for the treatment of patients with COVID-19 after the first 3 days of symptoms (Oxford, 2020).

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Rationale

Chloroquine is a closely related compound to hydroxychloroquine. There is no RCT-level evidence that chloroquine has different efficacy. There are sparse trials of chloroquine, especially compared with the evidence base for hydroxychloroquine. One population-based cohort study found evidence of efficacy of hydroxychloroquine but not chloroquine (Lammers et al., 2020). Thus, by analogy to hydroxychloroquine, chloroquine is not recommended for treatment of hospitalized COVID-19 patients. See the Hydroxychloroquine Rationale for Recommendation for details.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Chloroquine, Prophylaxis; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 766 articles in PubMed, 18,961 in Scopus, 33 in CINAHL, 1,279 in Cochrane Library, 9,500 in Google Scholar, and 0 from other sources†. We considered for inclusion 0 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 0 articles considered for inclusion, 0 randomized trials and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

†The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Hydroxychloroquine or Chloroquine for Widespread Prophylaxis Against COVID-19

No Recommendation

There is no recommendation for or against the use of hydroxychloroquine and chloroquine for widespread prophylaxis against COVID-19.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Rationale

One high-quality trial of hydroxychloroquine (without zinc) for postexposure prophylaxis suggested no statistically significant benefit (11.8% vs. 14.3%, 17.5% reduction, $p=0.35$), although there was a

17% reduction of risk (Boulware et al., 2020); thus, underpowering is possible. One large RCT found a 21% reduction in risk of COVID-19 (Seet et al., 2021). A cluster-randomized trial found a nonsignificant 8.1% reduction in PCR-confirmed COVID (Mitja, 2020). Some RCTs found lack of efficacy for prophylaxis among healthcare workers (Abella et al., 2020) (Syed et al., 2021), whereas two others trended towards modest efficacy (Rojas-Serrano et al., 2021, Naggie et al., 2021). Other RCTs found lack of efficacy (Barnabas et al., 2021)(Mitjà et al., 2021). A meta-analysis was performed with multiple RCTs that included early use of HCQ, ranging from pre-diagnosis to within a few days of symptoms onset (Ladapo et al., 2020). These trials are naturally individually underpowered for severe outcomes such as mortality as they tend to include younger, healthier patients. This meta-analysis of 5 RCTs that analyzed 5,577 patients found that all studies trended towards efficacy and the combined data showed a statistically significant 24% reduction in composite risk of infection, hospitalization, and death (Ladapo et al., 2020). A systematic review found weak and conflicting evidence (Hernandez et al., 2020). As evidence for widespread prophylactic use is weak and somewhat conflicting, there is no recommendation.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to November 2020 using the following terms:

Hydroxychloroquine, Prophylaxis; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 73 articles in PubMed, 180 in Scopus, 25 in CINAHL, 41 in Cochrane Library, 8,280 in Google Scholar, and 0 from other sources†. We considered for inclusion 2 from PubMed, 4 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 3 from other sources. Of the 12 articles considered for inclusion, 3 randomized trials and 1 systematic review met the inclusion criteria. There were no exclusion criteria.

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to November 2020 using the following terms: Chloroquine Prophylaxis; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 73 articles in PubMed, 18 in Scopus, 4 in CINAHL, 44 in Cochrane Library, 9560 in Google Scholar, and 0 from other sources†. We considered for inclusion 1 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 0 from other sources. Of the 2 articles considered for inclusion, 0 randomized trials, 0 non-randomized trial, and 2 systematic review met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.3. AZITHROMYCIN

Azithromycin has been suggested to inhibit the growth of both the Zika and Ebola viruses, as well as prevent severe lower respiratory tract infections (82,83,84,85). Azithromycin has been used for treatment of COVID-19, as both stand-alone and combined therapy (86,87,88).

Azithromycin for Treatment of COVID-19 - Use in First 3 Days of Symptoms

No Recommendation

There is no recommendation for or against the use of azithromycin in the first 3 days of COVID-19 symptoms.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

A moderate-quality RCT found the addition of azithromycin (AZT) to standard care that included HCQ produced no apparent benefit among hospitalized patients with severe COVID-19 (Lau et al., 2020). A moderate-quality RCT found benefits with shortened hospital stay, improved oxygenation, and reduced respiratory rates associated with the addition of AZT to a combination of HCQ and lopinavir/ritonavir (Sekhavati et al., 2020).

There are no quality RCTs regarding early treatment. Adjunctive use with hydroxychloroquine in severely affected patients with COVID-19. For severely affected patients, AZT has been added (Gautret et al., 2020), but ECG monitoring should be particularly considered when adjunctive therapy with agents prolonging the QT interval is considered, including azithromycin plus HCQ/CQ (see Harms). Low-quality evidence suggests better efficacy if administered earlier in the clinical course when viral replication is occurring. There is no quality evidence of efficacy after ARDS is established (CHEN et al., 2020).

Benefits

Theoretical reduced need for a ventilator or ICU stay.

Harms

Negligible for most patients undergoing short-course use. There are concerns about the potential for prolonged corrected QT intervals when used in combination therapy, and thus arrhythmias. ECG monitoring is particularly indicated in those undergoing adjunctive treatment with HCQ/CQ with underlying cardiovascular disease, history of prolonged QT, unexplained syncope, family history of premature sudden cardiac death, electrolyte abnormalities, renal insufficiency, and use of other drugs reported to prolong QT intervals, including when there is planned adjunctive use with hydroxychloroquine/chloroquine.

Frequency/Dose/Duration

The regimen used for treatment of COVID is azithromycin 500mg on day 1 and then 250 mg/day for 4 days (Gautret et al., 2020, Gautret et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

Late administration of AZT has been assessed in multiple RCTs and been found to be mostly ineffective (Sivapalan et al., 2021, Horby, 2021, Johnston et al., 2021, Hinks et al., 2021, Butler et al., 2021). One trial of HCQ/AZT and another of AZT alone that included patients who were mostly beyond the early stage found a lack of efficacy (Rodrigues et al., 2021, Oldenburg et al., 2021). Another comparative trial found comparable (in)efficacy of HCQ and AZT (Brown et al., 2021). One RCT has suggested no difference between AZT, HCQ, and the combination for treatment of hospitalized patients (Cavalcanti et al., 2020). Thus, as the data are mostly consistent, AZQ is not recommended for late treatment of COVID-19.

One trial of uncertain symptom duration among patients with mild COVID-19 suggested faster resolution of symptoms with AZT (Rashad et al., 2021).

Most non-randomized but controlled studies have suggested some evidence of efficacy, particularly for early adjunctive use when combined with HCQ (Gautret et al., 2020, Lagier et al., 2020, Davido et al., 2020, Arshad et al., 2020, Guérin et al., 2020, Gautret et al., 2020), although some other studies have suggested a lack of efficacy (Rosenberg et al., 2020, Sbidian et al., 2020). Thus, there is no recommendation for use of AZT in the early phase of COVID-19.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to November 2020 using the following terms: Azithromycin ; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 164 articles in PubMed, 1161 in Scopus, 40 in CINAHL, 77 in Cochrane Library, 5170 in Google Scholar, and 16 from other sources†. We considered for inclusion 19 from PubMed, 9 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 16 from other sources. Of the 45 articles considered for inclusion, 2 randomized trials, 2 non-randomized trials, 4 case series, 9 retrospective studies, and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Azithromycin for Treatment of COVID-19 - Use Beyond First 3 Days of Symptoms

Not Recommended

Azithromycin is not recommended for the adjunctive treatment of selected patients with more severe COVID-19.

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Indications

A moderate-quality RCT found the addition of azithromycin (AZT) to standard care that included HCQ produced no apparent benefit among hospitalized patients with severe COVID-19 (Lau et al., 2020). A moderate-quality RCT found benefits with shortened hospital stay, improved oxygenation, and reduced respiratory rates associated with the addition of AZT to a combination of HCQ and lopinavir/ritonavir (Sekhavati et al., 2020).

There are no quality RCTs regarding early treatment. Adjunctive use with hydroxychloroquine in severely affected patients with COVID-19. For severely affected patients, AZT has been added (Gautret et al., 2020), but ECG monitoring should be particularly considered when adjunctive therapy with agents prolonging the QT interval is considered, including azithromycin plus HCQ/CQ (see Harms). Low-quality evidence suggests better efficacy if administered earlier in the clinical course when viral replication is occurring. There is no quality evidence of efficacy after ARDS is established (CHEN et al., 2020).

Benefits

Theoretical reduced need for a ventilator or ICU stay.

Harms

Negligible for most patients undergoing short-course use. There are concerns about the potential for prolonged corrected QT intervals when used in combination therapy, and thus arrhythmias. ECG monitoring is particularly indicated in those undergoing adjunctive treatment with HCQ/CQ with underlying cardiovascular disease, history of prolonged QT, unexplained syncope, family history of premature sudden cardiac death, electrolyte abnormalities, renal insufficiency, and use of other drugs reported to prolong QT intervals, including when there is planned adjunctive use with hydroxychloroquine/chloroquine.

Frequency/Dose/Duration

The regimen used for treatment of COVID is azithromycin 500mg on day 1 and then 250 mg/day for 4 days (Gautret et al., 2020, Gautret et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

Late administration of AZT has been assessed in multiple RCTs and been found to be mostly ineffective (Sivapalan et al., 2021, Johnston et al., 2021, Hinks et al., 2021, Butler et al., 2021)(Horby, 2021). One trial of HCQ/AZT and another of AZT alone that included patients who were mostly beyond the early stage found a lack of efficacy (Rodrigues et al., 2021, Oldenburg et al., 2021). Another comparative trial found comparable (in)efficacy of HCQ and AZT (Brown et al., 2021). One RCT has suggested no difference between AZT, HCQ, and the combination for treatment of hospitalized patients (Cavalcanti et al., 2020). Thus, as the data are mostly consistent, AZQ is not recommended for late treatment of COVID-19.

One trial of uncertain symptom duration among patients with mild COVID-19 suggested faster resolution of symptoms with AZT (Rashad et al., 2021).

Most non-randomized but controlled studies have suggested some evidence of efficacy, particularly for early adjunctive use when combined with HCQ (Lagier et al., 2020, Gautret et al., 2020, Davido et al., 2020, Arshad et al., 2020, Guérin et al., 2020, Gautret et al., 2020), although some other studies have suggested a lack of efficacy (Rosenberg et al., 2020, Sbidian et al., 2020). Thus, there is no recommendation for use of AZT in the early phase of COVID-19.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to November 2020 using the following terms: Azithromycin ; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 164 articles in PubMed, 1161 in Scopus, 40 in CINAHL, 77 in Cochrane Library, 5170 in Google Scholar, and 16 from other sources†. We considered for inclusion 19 from PubMed, 9 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 16 from other sources. Of the 45 articles considered for inclusion, 2 randomized trials, 2 non-randomized trials, 4 case series, 9 retrospective studies, and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.4. FAVIPIRAVIR

Favipiravir, a guanine analogue to inhibit RNA-dependent RNA polymerase, has been used to treat influenza. Favipiravir has now been used to treat mostly severely affected COVID-19 patients (89,90,91,92,93,94,95,96).

Favipiravir for the Treatment of COVID-19

No Recommendation

There is no recommendation for or against favipiravir the treatment of COVID-19.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Rationale

RCTs among those who were treated late and/or who initiated treatment while hospitalized are mostly negative. One trial with mostly late treatment patients found lack of benefit with favipiravir with and without HCQ (Bosaeed et al., 2021). A moderate-quality RCT found a lack of efficacy for combined favipiravir with interferon beta-1b compared with HCQ for moderate to severe COVID-19 pneumonia patients (Khamis et al., 2020). A comparative trial found no differences between favipiravir and lopinavir/ritonavir (Solaymani-Dodaran et al., 2021). A moderate-quality RCT found no evidence of benefit of favipiravir for viral clearance, although there was faster defervescence (Doi et al., 2020), while another low-quality study found faster viral clearance (Zhao et al., 2021). Another trial of many combinations found accelerated viral clearance but more deaths with favipiravir plus lopinavir-ritonavir (Atipornwanich et al., 2021).

There are two RCTs that appear to have included both early and late patients, with one reporting Favipiravir was associated with shortening of symptoms by 3 days compared with placebo (Shinkai et al., 2021), while another reported a trend towards earlier viral clearance (Udwadia et al., 2021).

One RCT comparing favipiravir with arbidol found no significant differences in the main clinical outcome measure, although fever and cough resolved more quickly in the favipiravir group (Chen et al., 2020). A low-quality RCT of baloxavir, marboxil, and favipiravir found no evidence that favipiravir accelerated viral clearance (Lou et al., 2020). There is one non-randomized controlled trial suggesting acceleration of viral clearance compared with lopinavir-ritonavir (Cai et al., 2020). One comparative trial possibly of early use among those hospitalized found favipiravir associated with a trend in modestly lower hospitalization days (Dabbous et al., 2021).

There is a dearth of evidence of early treatment and conflicting evidence for the late use of favipiravir. Thus, there is no recommendation for either early or late use.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Favipiravir, Amides, Pyrazines, T-705 cpd, 6-fluoro-3-hydroxy-2-pyrazinecarboxamide, Avigan; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 105 articles in PubMed, 4023 in Scopus, 28 in CINAHL, 118 in Cochrane Library, 13190 in Google Scholar, and 0 from other sources†. We considered for inclusion 23 from PubMed, 5 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 10 from Google Scholar, and 0 from other sources. Of the 37 articles considered for inclusion, 10 randomized trials and 3 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.5. LOPINAVIR/RITONAVIR

Lopinavir-ritonavir has been used for the treatment of COVID-19 (97,98,99,100,101,102,103,104,105,106).

Lopinavir/Ritonavir for the Treatment of COVID-19 - Combination Therapy

Recommended

Lopinavir-ritonavir is recommended in combination therapy (Hung et al., 2020), but is not recommended as a stand-alone treatment for COVID-19.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Most evidence has suggested a lack of efficacy of Lopinavir-Ritonavir and adjunctive use with ribavirin and interferon beta-1b in moderately and severely affected patients with COVID-19 (Hung et al., 2020). Some evidence has suggested better efficacy if administered within 7 days of symptom onset; after 7 days, data suggest no differences between this combination therapy and lopinavir-ritonavir (Hung et al., 2020).

Benefits

Faster symptom resolution, viral clearance, and hospital discharge. Reduced need for a ventilator or ICU stay.

Harms

Nausea, diarrhea, hepatitis.

Frequency/Dose/Duration

The regimen used for the treatment of COVID-19 is lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days (Hung et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

One open-label RCT found combination therapy of lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days to be superior to lopinavir-ritonavir (Hung et al., 2020). However, another trial found comparable faster clinical improvement (9 vs 11 days), fewer adverse events, and ~67% reduction in mortality (6.1 vs. 18.2%) when comparing treatment with interferon beta-1-b with treatment with the control group (lopinavir-ritonavir/HCQ or atazanavir/ritonavir/HCQ) (Rahmani et al., 2020), which could suggest that the only medication effective in the triple therapy is the interferon beta-1b. Another RCT found lack of efficacy and trends towards worse outcomes with Lopinavir-ritonavir (Arabi et al., 2021).

Lopinavir-ritonavir as a stand-alone antiviral treatment has been trialed in five RCTs, all of which showed a lack of efficacy compared with standard care (WHO, 2020, Cao et al., 2020, Li et al., 2020, Horby et al., 2020) (Reis et al., 2021). Another double-blind RCT also suggested lack of efficacy, although it may have been underpowered (Li et al., 2020). One RCT treated severe patients and the other treated mild/moderately severe patients at an average of 4–5 days duration. It is unclear if lopinavir-ritonavir would be effective if provided earlier in the clinical course. One trial with unclear symptom durations and baseline outcomes differences suggesting possible randomization failure nevertheless suggested potentially faster PCR negative results in 3 days with various combinations of drugs (either lopinavir/ritonavir-doxycycline; lopinavir/ritonavir-azithromycin; or azithromycin-HCQ) (Purwati et al., 2021). Lopinavir-ritonavir has also been suggested to be inferior to favipiravir in a non-randomized comparative trial [(Cai et al., 2020). Another comparative trial found no significant difference between lopinavir/ritonavir and umifenovir (Darazam et al., 2021).

Most studies suggest lopinavir is not associated with improved outcomes. Thus, it is not recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Lopinavir-Ritonavir; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 256 articles in PubMed, 0 in Scopus, 0 in CINAHL, 213 in Cochrane Library, 19300 in Google Scholar, and 0 from other sources†. We considered for inclusion 23 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 25 articles considered for inclusion, 5 randomized trials and 17 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Lopinavir/Ritonavir for the Treatment of COVID-19 - Stand-alone Treatment

Not Recommended

Lopinavir-ritonavir is not recommended as a stand-alone treatment for COVID-19.

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Indications

Most evidence has suggested a lack of efficacy. Adjunctive use with ribavirin and interferon beta-1b in moderately and severely affected patients with COVID-19 (Hung et al., 2020). Some evidence has suggested better efficacy if administered within 7 days of symptom onset; after 7 days, data suggest no differences between this combination therapy and lopinavir-ritonavir (Hung et al., 2020).

Benefits

Faster symptom resolution, viral clearance, and hospital discharge. Reduced need for a ventilator or ICU stay.

Harms

Nausea, diarrhea, hepatitis.

Frequency/Dose/Duration

The regimen used for the treatment of COVID-19 is lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days (Hung et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

One open-label RCT found combination therapy of lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days to be superior to lopinavir-ritonavir (Hung et al., 2020). However, another trial found comparable faster clinical improvement (9 vs 11 days), fewer adverse events, and ~67% reduction in mortality (6.1 vs. 18.2%) when comparing treatment with interferon beta-1b with treatment with the control group (lopinavir-ritonavir/HCQ or atazanavir/ritonavir/HCQ) (Rahmani et al., 2020), which could suggest that the only medication effective in the triple therapy is the interferon beta-1b. Another RCT found lack of efficacy and trends towards worse outcomes with Lopinavir-ritonavir (Arabi et al., 2021). Lopinavir-ritonavir as a stand-alone antiviral treatment has been trialed in five RCTs, all of which showed a lack of efficacy compared with standard care (Cao et al., 2020, Li et al., 2020, Horby et al., 2020) (Reis et al., 2021). Another double-blind RCT also suggested lack of efficacy, although it may have been underpowered (Li et al., 2020). One RCT treated severe patients and the other treated mild/moderately severe patients at an average of 4–5 days duration. It is unclear if lopinavir-

ritonavir would be effective if provided earlier in the clinical course. One trial with unclear symptom durations and baseline outcomes differences suggesting possible randomization failure nevertheless suggested potentially faster PCR negative results in 3 days with various combinations of drugs (either lopinavir/ritonavir-doxycycline; lopinavir/ritonavir-azithromycin; or azithromycin-HCQ) (Purwati et al., 2021). Lopinavir-ritonavir has also been suggested to be inferior to favipiravir in a non-randomized comparative trial [(Cai et al., 2020)]. Another comparative trial found no significant difference between lopinavir/retinovir and umifenovir (Darazam et al., 2021). Most studies suggest lopinavir is not associated with improved outcomes. Thus, it is not recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Lopinavir-Ritonavir; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 256 articles in PubMed, 0 in Scopus, 0 in CINAHL, 213 in Cochrane Library, 19300 in Google Scholar, and 0 from other sources†. We considered for inclusion 23 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 25 articles considered for inclusion, 5 randomized trials and 17 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.6. REMDESIVIR

Remdesivir has been used to treat COVID-19 (107,108,109,110,111,112,113,114) and is recommended by other systematic reviews (115).

Remdesivir for the Treatment of COVID-19 - Use in First 3 Days of Symptoms

No Recommendation

There is no recommendation for or against the use of remdesivir during the first 3 days of COVID-19 symptoms.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

Severe COVID-19 patients, with <94% O₂ saturation or need for O₂ supplementation, mechanical ventilation, or extracorporeal membrane oxygenation (Hinton, 2020). Patients included in trials had creatinine clearance >30 mL/min; ALT and AST <5 times upper limit of normal.

Benefits

Shortened ICU stay, but minimal to no impact on survival

Harms

Increased hepatic enzymes, diarrhea, rash, renal impairment, hypotension. However, the largest RCT did not report significantly increased adverse events in any category (Beigel et al., 2020).

Frequency/Dose/Duration

Remdesivir 200 mg IV on day 1, then 100 mg QD for 9 additional days (Beigel et al., 2020, Grein et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect.

Rationale

There is one high-quality RCT of remdesivir suggesting a lack of clinical efficacy, although it also suggests non-significant trends toward earlier clinical improvements (Wang et al., 2020). A larger, moderate-quality NIH trial showed modest efficacy, including 31% shorter ICU stays and earlier clinical improvements, although the finding was discovered after the endpoint was changed. A RCT comparing remdesivir with standard care found a trend towards better results with a 5-day course of remdesivir (Spinner et al., 2020). However, other RCTs found a lack of efficacy (WHO, 2020) (Ader et al., 2021, Mahajan et al., 2021) and another found lack of efficacy to either speed viral clearance and/or improve mortality among hospitalized patients (Barratt-Due et al., 2021). One trial suggested reduced hospital stay but other outcomes were negative (Abd-El salam et al., 2021). None of the RCTs was able to show statistically improved survival, although the NIH trial trended toward improved survival (Beigel et al., 2020). There is one case series suggesting a fairly low death rate (13%) (Grein et al., 2020) and another non-randomized study suggesting potential efficacy (Antinori et al., 2020). A low-quality RCT found no difference between 5 and 10 days of treatment (Goldman et al., 2020). There is evidence that remdesivir inhibits viral replication in vitro studies (Wang et al., 2020). It is possible that remdesivir is more effective if administered in the viral replication stage (Cubeddu et al., 2021, Glaus et al., 2020).

Remdesivir is invasive (IV), has minimal adverse effects, is high cost, has conflicting evidence of modest efficacy (particularly for the treatment of hospitalized patients requiring oxygen), and thus is selectively recommended. There are other treatments with stronger efficacy at reducing mortality (e.g., glucocorticosteroids, low-molecular-weight heparin).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Remdesivir; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV;

controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 471 articles in PubMed, 9083 in Scopus, 112 in CINAHL, 230 in Cochrane Library, 28800 in Google Scholar, and 0 from other sources†. We considered for inclusion 41 from PubMed, 3 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 47 articles considered for inclusion, 5 randomized trials and 21 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Remdesivir for the Treatment of COVID-19 - Use Beyond First 3 Days of Symptoms

Sometimes Recommended

Remdesivir is selectively recommended for the treatment of patients with COVID-19.

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Low

Indications

Severe COVID-19 patients, with <94% O₂ saturation or need for O₂ supplementation, mechanical ventilation, or extracorporeal membrane oxygenation (Hinton, 2020). Patients included in trials had creatinine clearance >30 mL/min; ALT and AST <5 times upper limit of normal.

Benefits

Shortened ICU stay, but minimal to no impact on survival

Harms

Increased hepatic enzymes, diarrhea, rash, renal impairment, hypotension. However, the largest RCT did not report significantly increased adverse events in any category (Beigel et al., 2020).

Frequency/Dose/Duration

Remdesivir 200 mg IV on day 1, then 100 mg QD for 9 additional days (Grein et al., 2020, Beigel et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect.

Rationale

There is one high-quality RCT of remdesivir suggesting a lack of clinical efficacy, although it also suggests non-significant trends toward earlier clinical improvements (Wang et al., 2020). A larger, moderate-quality NIH trial showed modest efficacy, including 31% shorter ICU stays and earlier clinical improvements, although the finding was discovered after the endpoint was changed. A RCT comparing remdesivir with standard care found a trend towards better results with a 5-day course of remdesivir (Spinner et al., 2020). However, other RCTs found a lack of efficacy (Ader et al., 2021, Mahajan et al., 2021) and another found lack of efficacy to either speed viral clearance and/or improve mortality among hospitalized patients (Barratt-Due et al., 2021). One trial suggested reduced hospital stay but other outcomes were negative (Abd-El Salam et al., 2021). None of the RCTs was able to show statistically improved survival, although the NIH trial trended toward improved survival (Beigel et al., 2020). There is one case series suggesting a fairly low death rate (13%) (Grein et al., 2020) and another non-randomized study suggesting potential efficacy (Antinori et al., 2020). A low-quality RCT found no difference between 5 and 10 days of treatment (Goldman et al., 2020). There is evidence that remdesivir inhibits viral replication in vitro studies (Wang et al., 2020). It is possible that remdesivir is more effective if administered in the viral replication stage (Cubeddu et al., 2021, Glaus et al., 2020). Remdesivir is invasive (IV), has minimal adverse effects, is high cost, has conflicting evidence of modest efficacy (particularly for the treatment of hospitalized patients requiring oxygen), and thus is selectively recommended. There are other treatments with stronger efficacy at reducing mortality (e.g., glucocorticosteroids, low-molecular-weight heparin).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Remdesivir; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 471 articles in PubMed, 9083 in Scopus, 112 in CINAHL, 230 in Cochrane Library, 28800 in Google Scholar, and 0 from other sources†. We considered for inclusion 41 from PubMed, 3 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 47 articles considered for inclusion, 5 randomized trials and 21 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.7. LOW-MOLECULAR-WEIGHT HEPARIN

Low-molecular-weight heparin has been used for the treatment of hospitalized, severely affected patients with COVID-19; the degree of coagulopathy has been associated with worsened survival (116) (117) (118) (119) (120) (121) (122) (123) (124) (125) (126) (127)(128). Fondaparinux and unfractionated heparin have also been recommended in the *Chest* guidelines (129).

Thrombectomies and other procedures have been performed in COVID-19 patients with known venous thromboembolism (129,130).

Low-Molecular-Weight Heparin for the Treatment of COVID-19

Recommended

Low-molecular-weight heparin is recommended for the treatment of select patients with COVID-19 (Di Perri, 2020, van Haren et al., 2020, Ayerbe et al., 2020, Hippensteel et al., 2020, Moores et al., 2020, Canoglu et al., 2020, D'Ardes et al., 2020, Falcone et al., 2020, Jose Ramon Gonzalez-Porras, 2020, Ma et al., 2020, Mattioli et al., 2020, Paolisso et al., 2020, Pavoni et al., 2020, Shi et al., 2020, Stattin et al., 2020, Vergori et al., 2020, White et al., 2020, Zhang et al., 2020, Gonzalez-Ochoa et al., 2020, Nadeem et al., 2021).

Strength of evidence Recommended, Evidence (C)

Level of confidence Moderate

Indications

Severely affected COVID-19 patients, especially those with known evidence or suspicion of having coagulopathy (e.g., small-vessel thromboses, large-vessel arterial and/or venous thromboses [e.g., infarcts, DVTs, pulmonary emboli], thrombocytopenia, increased D-dimer, increased fibrin degradation products, prolonged coagulation times). May also be indicated for those who are hospitalized and either (i) sedentary, as there is some evidence of post-mortem coagulopathy in those without pre-morbid suspicions of coagulopathy and/or (ii) on a worsening clinical trajectory that suggests trending towards critical status and/or cytokine storm (Nadkarni et al., 2020).

Benefits

Possible improved survival, improved oxygenation, reduced time on ventilator (Lemos et al., 2020), reduced risks of DVT, pulmonary emboli, myocardial infarction, cerebrovascular thromboembolic disease.

Harms

Usual risks of heparin, particularly bleeding complications. In rare cases, acute COVID-19 infection is associated with the development of anti-heparin- platelet factor 4 autoantibodies that bind heparin and platelets, and may be associated with thrombocytopenia and increased risk of thrombosis (Cai et al., 2020, Dragonetti et al., 2020). In that case, alternate methods of anticoagulation, such as with apixaban or others, should be instituted.

Frequency/Dose/Duration

Per manufacturer's recommendations. A stepped approach with more intensive prophylaxis for more severely affected patients has been reportedly successful (Atallah et al., 2020). Unfractionated heparin is another therapeutic option.

Indications for discontinuation

Recovery from COVID-19 and resolution of findings of coagulopathy with regaining of normal ambulation. Also discontinue for significant adverse effects. May be continued after hospital discharge for a period of time during recovery and while still not as active and ambulatory as pre-morbid.

Rationale

One RCT reported efficacy of enoxaparin over standard anticoagulation (unfractionated heparin, generally 5,000U TID) to significantly increase gas exchange and reduce need for ventilatory support (Lemos et al., 2020). A trial of sulodexide found reduced need for hospitalization and oxygen therapy (Gonzalez-Ochoa et al., 2020). Reductions in mortality have been reported in non-randomized studies (Ayerbe et al., 2020, Albani et al., 2020, Hsu et al., 2020, Paranjpe et al., 2020, Ionescu et al., 2020, Hanif et al., 2020), including an estimated 47–50% reduced risk of mortality among those on therapeutic anticoagulation among 4,389 in a hospital system (Nadkarni et al., 2020). Another cohort of patients on mechanical ventilation was found to have a 54% reduction in mortality (Paranjpe et al., 2020, Dobesh et al., 2020).

An early escalating thromboprophylactic approach has been suggested as preventive among hospitalized patients with less severe disease (Daughety et al., 2020).

Low-molecular-weight heparins are minimally invasive, have potentially significant adverse effects, are moderately costly, and have evidence suggesting associations with lower mortality rates and fewer complications among severely affected COVID-19 patients; thus, they are selectively recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to December 2020 using the following terms: Low Molecular Weight Heparin; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 60 articles in PubMed, 837 in Scopus, 11 in CINAHL, 22 in Cochrane Library, 4,410 in Google Scholar, and 0 from other sources[†]. We considered for inclusion 16 from PubMed, 21 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 13 from Google Scholar, and 0 from other sources. Of the 51 articles considered for inclusion, 2 randomized trials and 13 systematic reviews met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.8. FLUVOXAMINE

Fluvoxamine has been used for the treatment of COVID-19 patients (131)(132)(133). Observational evidence suggests that two SSRI antidepressants, fluvoxamine and fluoxetine (but not other SSRIs), reduce mortality by 10% (134).

Fluvoxamine for the Treatment of COVID-19

Recommended

Fluvoxamine is recommended for the treatment of patients with COVID-19, particularly within 7 days of symptoms onset.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Outpatients with moderate to severe symptoms who are within 7 days of onset of symptomatic disease (Lenze et al., 2020, Reis et al., 2022). Indicated for those with mild symptoms and risks of adverse outcomes or those who are trending towards worsening of symptoms.

Benefits

Reduces probability of clinical deterioration, including hospitalization (Lenze et al., 2020, Reis et al., 2022).

Harms

Headache, nausea, diarrhea, dry mouth, dizziness, increased sweating, feeling nervous, restlessness, fatigue, insomnia.

Frequency/Dose/Duration

Fluvoxamine 100mg TID for 15 days (Lenze et al., 2020).

Indications for discontinuation

Completion of a course of treatment, intolerance of adverse effects, recovery

Rationale

Two moderate-quality RCTs both suggest efficacy of fluvoxamine used within 7 days of symptoms onset to prevent clinical deterioration including hospitalization (Lenze et al., 2020, Reis et al., 2022). One trial found a 32% reduction in risk of hospitalization and/or going to the emergency room (Reis et al., 2022) and the other suggested fluvoxamine was associated with less clinical deterioration at 15 days compared with placebo (0% vs. 8.7%, $p=0.009$), although there also was no difference in the most severe symptom changes between the two groups (Lenze et al., 2020). Both trials suggest

efficacy to prevent clinical worsening. Fluvoxamine is non-invasive, has low adverse effects, is low cost, and is recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Fluvoxamine, Fluvoxamine Maleate; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 6 articles in PubMed, 203 in Scopus, 9 in CINAHL, 7 in Cochrane Library, 1367 in Google Scholar, and 0 from other sources†. We considered for inclusion 3 from PubMed, 4 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 7 articles considered for inclusion, 2 randomized trials and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.9. PAXLOVID

Paxlovid (nirmatrelvir/ritonavir) has been approved by the U.S. FDA through an [emergency use authorization](#) for the treatment of COVID-19 patients (135).

Paxlovid (Nirmatrelvir/Ritonovir) for Treatment of COVID-19

Recommended

Paxlovid (nirmatrelvir/ritonavir) is recommended for the treatment of patients with COVID-19, particularly within 3 days of symptom onset.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Outpatients with moderate to severe symptoms who are within 5 days, but ideally within 3 days of onset of symptoms (Mahase, 2021). Also indicated for those with mild symptoms and risks of adverse outcomes or those who are trending towards worsening of symptoms. The FDA EUA has not authorized use for treatment initiation among patients requiring hospitalization for severe or critical COVID-19 (Administration, 2022).

Benefits

89% reduction in the probability of hospitalization and/or death (Mahase, 2021).

Harms

Adverse effects include diarrhea, altered taste/small, hypertension, myalgia, elevated hepatic transaminases, clinical hepatitis, jaundice. There are significant adverse drug-drug Interactions especially including those dependent on CYP3A for clearance. Drugs recommended to be not taken with paxlovid include alfuzosin, pethidine, piroxicam, propoxyphene, ranolazine, amiodarone, dronedarone, flecainide, propafenone, quinidine, colchicine, lurasidone, pimozide, clozapine, dihydroergotamine, ergotamine, methylergonovine, lovastatin, simvastatin, sildenafil for pulmonary arterial hypertension (PAH), triazolam, oral midazolam, apalutamide, carbamazepine, phenobarbital, phenytoin, rifampin, and St. John's Wort (Drugs.com, 2021).

Frequency/Dose/Duration

Paxlovid (Nirmatrelvir/Ritonovir) 300mg nirmatrelvir (2-150mg tablets)/100mg (1-100mg tablet) BID for 5 days (Administration, 2022).

Indications for discontinuation

Completion of a course of treatment, intolerance of adverse effects.

Rationale

Interim analyses from one moderate-quality RCT suggests 89% reduction in risk of hospitalization and/or death associated with Paxlovid (Nirmatrelvir/Ritonovir), especially when used within 3 days of symptoms onset (Mahase, 2021). There were 10 deaths in the placebo group vs. 0 in the Paxlovid group over 28 days (Mahase, 2021). As interim analyses from one RCT suggest major efficacy to prevent hospitalization and/or death, Paxlovid (Nirmatrelvir/Ritonovir) is non-invasive, has considerable adverse effects, is high cost, and is recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Paxlovid, Nirmatrelvir, Nirmatrelvir and Ritonavir Drug Combination, Nirmatrelvir and Ritonavir; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 0 articles in PubMed, 0 in Scopus, 0 in CINAHL, 0 in Cochrane Library, 122 in Google Scholar, and 0 from other sources†. We considered for inclusion 0 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 2 articles considered for inclusion, 0 randomized trials and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant

literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.10. MOLNUPIRAVIR

Molnupiravir has been approved by the U.S. FDA through an emergency use authorization for the treatment of COVID-19 patients (136,137,138,139)(140).

Molnupiravir for the Treatment of COVID-19

Recommended

Molnupiravir is recommended for the treatment of patients with COVID-19, particularly within 3 days of symptoms onset.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Outpatients with moderate to severe symptoms who are within 5 days, but ideally within 3 days, of onset of symptoms (Fischer et al., 2021, Painter et al., 2021, Khoo et al., 2021)(Jayk Bernal et al., 2022). Also indicated for those with mild symptoms and risks of adverse outcomes or those who are trending towards worsening of symptoms. The FDA EUA has not authorized use for treatment initiation among patients requiring hospitalization for severe or critical COVID-19 (Administration et al., 2022).

Data suggest lack of efficacy among those with known positive SARS-CoV-2 nucleocapsid antibody levels at baseline, with data favoring placebo (Jayk Bernal et al., 2022).

Benefits

48% reduction in the probability of hospitalization and/or death (Jayk Bernal et al., 2022).

Harms

Adverse effects include diarrhea, nausea, dizziness. Not recommended for use during pregnancy, in children under 18 years of age due to potential altered bone/cartilage growth, and among women who are breastfeeding (Administration, 2022).

Frequency/Dose/Duration

Molnupiravir 800mg (4-200mg tablets) BID for 5 days (Administration, 2022).

Indications for discontinuation

Completion of a course of treatment, intolerance of adverse effects.

Rationale

One moderate quality RCT reported a 48% reduction in risk of hospitalization and/or death; there also was 1 death with molnupiravir vs. 9 in the placebo group (Jayk Bernal et al., 2022). Another RCT reported marked reductions in viral detection at 3 days of treatment with a dose of 800mg (2 vs 17%, $p=0.02$), while slower but superior clearance to placebo was shown with 400mg (Fischer et al., 2021). Two early-phase trials in combination with the clinical trials suggest low toxicity and a dose of 800mg (Khoo et al., 2021, Painter et al., 2021, Fischer et al., 2021) (Jayk Bernal et al., 2022). The RCTs suggest efficacy of molnupiravir to prevent hospitalization and/or death. It is non-invasive, has low adverse effects, is high cost, and is recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Molnupiravir, Lagevrio, EIDD-2801, MK-4482; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 10 articles in PubMed, 141 in Scopus, 1 in CINAHL, 24 in Cochrane Library, 574 in Google Scholar, and 0 from other sources†. We considered for inclusion 4 from PubMed, 1 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 5 articles considered for inclusion, 3 randomized trials and 1 systematic review met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.11. IL-6 RECEPTOR ANTAGONISTS

Various interleukin-6 receptor antagonists have been used for the treatment of hospitalized patients with COVID-19 (141,142,143,144,145,146,147,148,149,150,151,152,153,154,155,156,157,158,159, 160,161,162,163,164,165,166,167,168,169,170,171,172,173,174,175).

IL-6 Receptor Antagonists (Tocilizumab, Sarilumab, and Siltuximab) for the Treatment of COVID-19

Not Recommended

Interleukin-6 inhibitors (sarilumab, siltuximab, and tocilizumab) are not recommended for the treatment of selected patients with COVID-19.

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Rationale

One moderate-quality trial suggested a reduced need for mechanical ventilation but no improved survival (Salama et al., 2020). One trial found improved organ-support free days and survival with both tocilizumab and sarilumab (Gordon et al., 2021), and another suggested modest efficacy (Horby et al., 2021). However, six other moderate-quality RCTs found a lack of efficacy of tocilizumab (Stone et al., 2020, Salvarani et al., 2020, Rosas et al., 2021) and sarilumab (Lescure et al., 2021, Veiga et al., 2021, Wang et al., 2020, Soin et al., 2021). Two suggested a trend towards efficacy (Sivapalasingam et al., 2021, Lescure et al., 2021). One moderate-quality RCT found trends towards reduced mortality by 2 weeks but not 4 weeks associated with tocilizumab (Hermine et al., 2020). One controlled study suggested increased adjusted survival rates among the group of patients treated with tocilizumab, although there were baseline differences likely favoring survival among the treated (Somers et al., 2020). Another controlled but non-randomized study of tocilizumab added to a standard-care regimen of HCQ, lopinavir, plus ritonavir suggested efficacy if administered earlier in the hospital course (Capra et al., 2020). One retrospective study found no benefit of tocilizumab (Campochiaro et al., 2020). One case series suggested significant survival and oxygenation benefits (Xu et al., 2020).

As there is now evidence that mostly suggests of a lack of efficacy of the IL-6 receptor antagonists, they are not recommended. There also are currently other treatments with demonstrated efficacy.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: interleukin-6, tocilizumab, sarilumab, siltuximab; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 11,008 articles in PubMed, 28,542 in Scopus, 142 in CINAHL, 68 in Cochrane Library, 30,600 in Google Scholar, and 10 from other sources[†]. We considered for inclusion 25 from PubMed, 0 from Scopus, 0 from CINAHL, 1 from Cochrane Library, 2 from Google Scholar, and 10 from other sources. Of the 38 articles considered for inclusion, 12 randomized trials and 24 systematic reviews met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.12. ADALIMUMAB

Adalimumab for the Treatment of COVID-19

Not Recommended

Adalimumab is not recommended for the treatment of patients with COVID-19.

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Rationale

One moderate-quality trial suggested lack of efficacy (Fakharian et al., 2021). Thus, adalimumab is not recommended.

8.13. BARICITINIB

Baricitinib is an orally bioavailable reversible inhibitor of Janus kinases 1 and 2 (JAK 1/2) typically used to rheumatoid arthritis. It has anti-inflammatory, immunomodulating, and antineoplastic activities, and has an FDA emergency use authorization (EUA) for use in COVID-19 infection due to its antiviral effects. Baricitinib has been used for the treatment of patients with COVID-19 (176) (177) (178) (179) (180).

Baricitinib for the Treatment of COVID-19

No Recommendation

There is no recommendation regarding baricitinib for the treatment of COVID-19 (Kalil et al., 2020).

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

The U.S. FDA issued an emergency use authorization for use in combination with remdesivir (Administration, 2020, Administration, 2020). Severely affected patients with COVID-19 with cytokine storm manifestations, including ARDS. Also indicated for those requiring supplemental oxygen and/or mechanical ventilation. Other treatments may be combined (e.g., glucocorticosteroids). The U.S. FDA issued an Emergency Use Authorization for use in combination with remdesivir (Administration, 2020, Administration, 2020).

Benefits

Data conflict regarding efficacy for prior variants.

Harms

Fever, chills, tiredness, muscle pain, increased urination, stomach pain, diarrhea, weight loss, cough, dyspnea.

Frequency/Dose/Duration

Doses used have included 4mg loading and 2mg/day for or 4mg/day (Rodriguez-Garcia et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effects.

Rationale

RCTs address variants prior to omicron. Regardless, one RCT found lack of efficacy (Marconi et al., 2021). Another found that adding baricitinib to remdesivir compared with remdesivir alone resulted in one less day of ICU stay. The evidence was stronger in the non-mechanical ventilated group with a 44% reduction in recovery time, and there was a trend in a 35% reduction in 28-day mortality (Kalil et al., 2020, Administration, 2020, Administration, 2020).

There are multiple non-randomized studies suggesting efficacy at mitigating the cytokine storm. A non-randomized trial found that the addition of baricitinib to glucocorticosteroids was associated with improved clinical outcomes, including an 82% reduced need for supplemental oxygen at discharge (Rodriguez-Garcia et al., 2020). A comparative consecutive case series suggested significant benefits, such as eliminating ICU transfers and 58% vs. 8% discharge at 2 weeks (Cantini et al., 2020).

Baricitinib is invasive, has some adverse effects, is costly, has conflicting evidence of modest efficacy, and has some evidence suggesting efficacy, although that evidence was developed prior to the omicron variant; thus, there is no recommendation.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to November 2020 using the following terms: Baricitinib, Olumiant; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 15 articles in PubMed, 1,177 in Scopus, 2 in CINAHL, 14 in Cochrane Library, 2,670 in Google Scholar, and 1 from other sources†. We considered for inclusion 1 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 8 from Google Scholar, and 1 from other sources. Of the 10 articles considered for inclusion, 1 randomized trial and 2 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.14. CASIRIVIMAB PLUS IMDEVIMAB (REGENERON)

Casirivimab plus Imdevimab are recombinant human monoclonal antibodies that bind to nonoverlapping epitopes of the SARS-CoV-2 spike protein receptor-binding domain and have been used to treat COVID-19. These have been approved for use by [FDA under the emergency use authorization provision](#) (181).

Casirivimab plus Imdevimab (Regeneron) for the Treatment of COVID-19

No Recommendation

There is no recommendation regarding the combination of casirivimab plus imdevimab (Regeneron) for the treatment of patients with mild to moderate COVID-19 at risk of severe disease. The National Institutes of Health has cautioned that the combination of casirivimab plus imdevimab is not likely to have significant efficacy against the omicron variant ((NIH), 2022).

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

Generally only for outpatient treatment of mild to moderate COVID-19 cases and at high risk of disease progression. Criteria for adults included BMI 35+, chronic renal disease, diabetes mellitus, immunocompromising condition, current receipt of immunosuppressive treatment, age 65+, age 55+ with comorbidity (cardiovascular disease, hypertension, COPD). Oxygen therapy is an exclusion.

Prophylactic use is for those not fully vaccinated or not expected to mount an adequate immune response and either exposed to an infected close contact or who are at high risk from an individual in the same institutional setting (e.g., nursing home, prison). However, as the combination of Casirivimab plus Imdevimab targets the receptor binding site, it is believed to have lower efficacy against the omicron variant, NIH has cautioned that efficacy is not anticipated ((NIH), 2022); thus, sotrovimab is preferably recommended if available.

Benefits

Data conflict regarding efficacy for prior variants.

Harms

Allergic reactions which may be severe; other than localized complications from parenteral administration.

Frequency/Dose/Duration

N/A

Indications for discontinuation

Completion of a course, intolerance, adverse effect.

Rationale

The RCTs addressed prior variants before omicron. A moderate-quality RCT found significant reductions in viral load among those infected, with greater reductions including among those with higher viral loads (Weinreich et al., 2021). Data provided to the FDA suggest a reduction of 67% in the risk of hospitalization (9% vs. 3%) (NIH, 2020). A prophylaxis trial of subcutaneous administration found development of symptomatic infection in 1.5% vs. 7.8% among controls (O'Brien et al., 2021). An open label trial among those hospitalized found a 20% reduction in risk of mortality at 28 days (Horby et al., 2021).

The combination of Casirivimab/imdevimab is invasive, has some adverse effects, is high cost, has past evidence of efficacy, is not anticipated to have significant efficacy against omicron ((NIH), 2022), and thus there is no recommendation. Sotrovimab is preferably recommended if available. Casirivimab/imdevimab may be a reasonable choice when other options are exhausted.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Casirivimab, Imdevimab; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 27 articles in PubMed, 393 in Scopus, 4 in CINAHL, 15 in Cochrane Library, 1,682 in Google Scholar, and 0 from other sources†. We considered for inclusion 0 from PubMed, 1 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 0 from other sources. Of the 3 articles considered for inclusion, 3 randomized trials and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.15. BAMLANIVIMAB PLUS ETESEVIMAB

Bamlanivimab is a neutralizing monoclonal IgG1 antibody that targets the receptor-binding domain of the spike protein of SARS-CoV-2 and has been used to treat COVID-19, often with etesevimab. These have been approved for use by FDA under the emergency use authorization provision (182) (183). The National Institutes of Health has cautioned that the combination of bamlanivimab with etesevimab is not likely to have significant efficacy against the omicron variant (184).

Bamlanivimab plus Etesevimab for the Treatment of COVID-19 - Within 5 Days of Symptom Onset

No Recommendation

There is no recommendation for or against bamlanivimab plus etesevimab are not recommended for the treatment of patients within 5 days of mild to moderate COVID-19 (IDSA, 2021). Sotrovimab is predicted to have preserved efficacy against the omicron variant and thus is recommended as preferable when available ((NIH), 2022).

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

Generally only for outpatient treatment of patients with mild to moderate COVID-19 cases and those at high risk of disease progression and when sotrovimab is unavailable. Ideally within 3-5 days of symptoms onset. FDA criteria for adults include BMI 35+, chronic renal disease, diabetes mellitus, immunocompromising condition, current receipt of immunosuppressive treatment, age 65+, age 55+ with comorbidity (cardiovascular disease, hypertension, COPD). Oxygen therapy is an exclusion.

As the receptor binding site is targeted by these antibodies, and while not proven, this is believed to result in less efficacy than with e.g., Sotrovimab when used to treat subsequent variants including omicron ((NIH), 2022).

Benefits

Reduced risk of hospitalization and death.

Harms

Unclear. Reported reactions include anaphylaxis and a serious infusion-related reaction.

Frequency/Dose/Duration

One IV infusion of 2,800mg bamlanivimab. May also be given in combination with 2,800mg IV etesevimab.

Indications for discontinuation

Completion of a course, intolerance, adverse effect.

Rationale

RCTs involve variants prior to omicron. One moderate-quality trial found marked reductions in the need for hospitalization or need for emergency room visits compared with placebo, while also reporting reduced viral loads (Gottlieb et al., 2021). Another moderate-quality RCT that also combined treatment with etesevimab found 67% reduced risk of hospitalization and 0 vs 10 deaths in the placebo group (Dugan et al., 2021). Data provided to the FDA suggest a reduction of 68–84%

in the risk of combined 28-day hospitalization, emergency department visit, or death (NIH, 2020). Another study suggested a 72% reduction in the risk of hospitalization among those at high risk (NIH, 2020).

An RCT found no difference in recovery time from a combination of bamlanivimab and remdesivir among hospitalized patients (Lundgren et al., 2022). Another small hospital-based dose-ranging RCT did not report statistical outcomes data (Chen et al., 2021). Thus, bamlanivimab is not recommended among hospitalized or late cases.

One moderate-quality RCT suggested 44% reduced risk of skilled nursing home residents and staff contracting COVID-19 within 8 weeks of randomization; there were 5 vs. 0 deaths with all deaths in the placebo group (Cohen et al., 2021).

The combination of bamlanivimab/etesevimab is invasive, has some adverse effects, is high cost, has past evidence of efficacy, is not anticipated to have significant efficacy against omicron ((NIH), 2022) and thus there is no recommendation. Sotrovimab is preferably recommended if available ((NIH), 2022). Bamlanivimab/etesevimab may be a reasonable choice when other options are exhausted.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Bamlanivimab; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 41 articles in PubMed, 345 in Scopus, 10 in CINAHL, 0 in Cochrane Library, 1520 in Google Scholar, and 0 from other sources†. We considered for inclusion 11 from PubMed, 1 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 0 from other sources. Of the 14 articles considered for inclusion, 10 randomized trials and 4 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Bamlanivimab plus Etesevimab for the Treatment of COVID-19 - Hospitalized or Late Cases

Not Recommended

Bamlanivimab plus etesevimab is not recommended for the treatment of patients hospitalized with COVID-19 (IDSA, 2021).

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Indications

Generally only for outpatient treatment of patients with mild to moderate COVID-19 cases and those at high risk of disease progression and when sotrovimab is unavailable. Ideally within 3-5 days of symptoms onset. FDA criteria for adults include BMI 35+, chronic renal disease, diabetes mellitus, immunocompromising condition, current receipt of immunosuppressive treatment, age 65+, age 55+ with comorbidity (cardiovascular disease, hypertension, COPD). Oxygen therapy is an exclusion.

As the receptor binding site is targeted by these antibodies, and while not proven, this is believed to result in less efficacy than with e.g., Sotrovimab when used to treat subsequent variants including omicron ((NIH), 2022).

Benefits

Reduced risk of hospitalization and death.

Harms

Unclear. Reported reactions include anaphylaxis and a serious infusion-related reaction.

Frequency/Dose/Duration

One IV infusion of 2,800mg bamlanivimab. May also be given in combination with 2,800mg IV etesevimab.

Indications for discontinuation

Completion of a course, intolerance, adverse effect.

Rationale

RCTs involve variants prior to omicron. One moderate-quality trial found marked reductions in the need for hospitalization or need for emergency room visits compared with placebo, while also reporting reduced viral loads (Gottlieb et al., 2021). Another moderate-quality RCT that also combined treatment with etesevimab found 67% reduced risk of hospitalization and 0 vs 10 deaths in the placebo group (Dogan et al., 2021). Data provided to the FDA suggest a reduction of 68–84% in the risk of combined 28-day hospitalization, emergency department visit, or death. Another study suggested a 72% reduction in the risk of hospitalization among those at high risk. An RCT found no difference in recovery time from a combination of bamlanivimab and remdesivir among hospitalized patients (Lundgren et al., 2022). Another small hospital-based dose-ranging RCT did not report statistical outcomes data (Chen et al., 2021). Thus, bamlanivimab is not recommended among hospitalized or late cases. One moderate-quality RCT suggested 44% reduced risk of skilled nursing home residents and staff contracting COVID-19 within 8 weeks of randomization; there were 5 vs. 0 deaths with all deaths in the placebo group (Cohen et al., 2021). The combination of bamlanivimab/etesevimab is invasive, has some adverse effects, is high cost, has past evidence of efficacy, is not anticipated to have significant efficacy against omicron ((NIH), 2022) and thus there is no recommendation. Sotrovimab is preferably recommended if available ((NIH), 2022). Bamlanivimab/etesevimab may be a reasonable choice when other options are exhausted.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Bamlanivimab;

coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 41 articles in PubMed, 345 in Scopus, 10 in CINAHL, 0 in Cochrane Library, 1520 in Google Scholar, and 0 from other sources†. We considered for inclusion 11 from PubMed, 1 from Scopus, 1 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 0 from other sources. Of the 14 articles considered for inclusion, 10 randomized trials and 4 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.16. SOTROVIMAB

Sotrovimab is a neutralizing monoclonal IgG antibody that targets an evolutionarily-conserved epitope that is not on the rapidly evolving receptor-binding domain of the spike protein of SARS-CoV-2 and has been used to treat COVID-19. It has been approved for use by the U.S. FDA under the emergency use authorization provision (185)(186)(187).

Sotrovimab for the Treatment of COVID-19 - Within 5 Days of Symptom Onset

Recommended

Sotrovimab is moderately recommended for the treatment of patients with mild to moderate COVID-19 (Administration, 2021, Administration et al., 2022).

Strength of evidence Moderately Recommended, Evidence (B)

Level of confidence Low

Indications

Generally only for outpatient treatment of patients with mild to moderate COVID-19 cases and those at high risk of disease progression. Ideally within 3-5 days of symptoms onset. FDA criteria for adults include age 65+, BMI 25+, pregnancy, chronic renal disease, diabetes mellitus, immunocompromising condition, current receipt of immunosuppressive treatment, cardiovascular disease, hypertension, chronic lung disease, sickle cell disease, neurodevelopmental disorders, and having medical appliance dependence (e.g., tracheostomy, gastrostomy, positive pressure ventilation). Oxygen therapy, increased oxygenation requirements compared with baseline, and hospitalization status are exclusions.

Benefits

Reduced risk of hospitalization and death.

Harms

Hypotension, dizziness, arrhythmias, chest pain, dyspnea, fever, chills, nausea, headache, confusion, sweating, muscle pain.

Frequency/Dose/Duration

One IV infusion of 500mg sotrovimab.

Indications for discontinuation

N/A

Rationale

The RCTs preceded the omicron variant. However, sotrovimab is projected by NIH to have retained efficacy against this variant unlike most other monoclonal antibodies ((NIH), 2022). In one high-quality RCT of early use of sotrovimab, disease progression with hospitalization or death occurred among 7% on placebo vs. 1% on sotrovimab (Gupta et al., 2021). One RCT of use among hospitalized patients found a lack of efficacy (Self et al., 2021).

Sotrovimab is invasive, has some adverse effects, is high cost, has evidence of efficacy for early use but not late use, and is anticipated to have significant efficacy against omicron ((NIH), 2022). Thus, it is recommended for early, but not late, use.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Sotrovimab; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 8 articles in PubMed, 90 in Scopus, 1 in CINAHL, 8 in Cochrane Library, 377 in Google Scholar, and 0 from other sources[†]. We considered for inclusion 3 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 0 from other sources. Of the 4 articles considered for inclusion, 2 randomized trials and 1 systematic review met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Sotrovimab for the Treatment of COVID-19 - Hospitalized or Late Cases

Not Recommended

Sotrovimab is not recommended for the treatment of patients with late or hospitalized cases of COVID-19 (Administration, 2021, Administration et al., 2022).

Strength of evidence Not Recommended, Evidence (C)

Level of confidence Low

Indications

Generally only for outpatient treatment of patients with mild to moderate COVID-19 cases and those at high risk of disease progression. Ideally within 3-5 days of symptoms onset. FDA criteria for adults include age 65+, BMI 25+, pregnancy, chronic renal disease, diabetes mellitus, immunocompromising condition, current receipt of immunosuppressive treatment, cardiovascular disease, hypertension, chronic lung disease, sickle cell disease, neurodevelopmental disorders, and having medical appliance dependence (e.g., tracheostomy, gastrostomy, positive pressure ventilation). Oxygen therapy, increased oxygenation requirements compared with baseline, and hospitalization status are exclusions.

Benefits

Reduced risk of hospitalization and death.

Harms

Hypotension, dizziness, arrhythmias, chest pain, dyspnea, fever, chills, nausea, headache, confusion, sweating, muscle pain.

Frequency/Dose/Duration

One IV infusion of 500mg sotrovimab.

Indications for discontinuation

N/A

Rationale

The RCTs preceded the omicron variant. However, sotrovimab is projected by NIH to have retained efficacy against this variant unlike most other monoclonal antibodies ((NIH), 2022). In one high-quality RCT of early use of sotrovimab, disease progression with hospitalization or death occurred among 7% on placebo vs. 1% on sotrovimab (Gupta et al., 2021). One RCT of use among hospitalized patients found a lack of efficacy (Self et al., 2021). Sotrovimab is invasive, has some adverse effects, is high cost, has evidence of efficacy for early use but not late use, and is anticipated to have significant efficacy against omicron ((NIH), 2022). Thus, it is recommended for early, but not late, use.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Sotrovimab; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 8 articles in PubMed, 90 in Scopus, 1 in CINAHL, 8 in Cochrane Library, 377 in Google Scholar, and 0 from other sources†. We considered for inclusion 3 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 1 from Google Scholar, and 0 from other sources. Of the 4 articles considered for inclusion, 2 randomized trials and 1 systematic review met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.17. BEBTELOVIMAB

Bebtelovimab for the Treatment of COVID-19

Recommended

Bebtelovimab is recommended for treatment of COVID-19 under the FDA Emergency Use Authorization process (February 11, 2022).

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Low

Indications

Adults and teenagers 12 years of age and older with an early case of confirmed COVID-19 within 7 days of symptoms onset, who are at high risk of progression to severe disease. Not indicated for those hospitalized and/or on oxygen therapy. FDA EUA does not authorize use for severe disease (Administration, 2022).

Benefits

Purported improvement in the clinical case and reduction in risk of progression.

Harms

Infusion-related symptoms/signs, pruritis, and rash. Limited data are not yet available (Administration, 2022).

Frequency/Dose/Duration

Bebtelovimab 175mg IV, one administration

Indications for discontinuation

N/A

Rationale

Bebtelovimab has no phase 3 trials published in the peer-reviewed literature. There are preliminary data published (Administration, 2022)(Administration, 2022) on which the FDA issued its Emergency Use Authorization. Bebtelovimab is minimally invasive, has low adverse effects, is high cost, has preliminary evidence of efficacy, and is thus recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: bebtelovimab, LY-CoV1404; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 1 article in PubMed, 0 in Scopus, 0 in CINAHL, 0 in Cochrane Library, 8 in Google Scholar, and 0 from other sources†. We considered for inclusion 1 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 1 article considered for inclusion, 0 randomized trials and 0 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.18. IVERMECTIN

Ivermectin has been used for the treatment of COVID-19 (188)(189) (190) (191) (192) (193) (194) (195)(196). There is experimental evidence of its ability to inhibit viral replication (197).

Ivermectin for the Treatment of COVID-19 - Late Onset of Symptoms**Not Recommended**

Ivermectin is not recommended for late treatment of COVID-19 (Podder et al., 2020, Chaccour et al., 2020, Krolewiecki et al., 2020, Niaee et al., 2020, Chowdhury et al., 2020, Hashim et al., 2020, Elgazzar et al., 2020, Ahmed et al., 2020, Alam et al., 2020, Behera et al., 2020, Cadegiani et al., 2020, Camprubí et al., 2020, Gorial et al., 2020, Heidary et al., 2020, Ortiz-Muñoz et al., 2020, Padhy et al., 2020, Rajter et al., 2020, Soto-Becerra et al., 2020).

Strength of evidence Not Recommended, Insufficient Evidence (I)

Level of confidence Low

Indications

Outpatients with COVID-19 and less than 5 days of symptoms.

Benefits

Reduced risk of progression, shorter recovery, reduced hospitalization, and faster viral clearance.

Harms

Dizziness, loss of appetite, nausea, vomiting, stomach pain, diarrhea, constipation, weakness, headache, myalgias, rash.

Frequency/Dose/Duration

Single dose of ivermectin 12 mg and doxycycline 100mg BID for 5 days (Mahmud et al., 2021).

Indications for discontinuation

Completion of course, adverse effects, intolerance

Rationale

There are many RCTs assessing ivermectin. Trials used different dose regimens (including single dose), in patients of varying symptoms duration. One metanalysis of controlled studies reported 61% reductions in mortality, time to recovery, and time to viral clearance (Marik et al., 2021).

There are several trials assessing use of ivermectin for early treatment (e.g., within 3-5 days). One high-quality trial of a combination of a single dose of ivermectin 12 mg and doxycycline 100mg BID for 5 days found 51% reduced risk of progression to serious disease and recovery averaged 2 fewer days (Mahmud et al., 2021). Another high-quality trial found a non-significant 33% reduced risk of hospitalization among those treated with weight-dependent dosing of Ivermectin (e.g., among 80-110kg, 18mg and another 18mg at 24 hrs) at 3-6 days of symptoms (Vallejos et al., 2021). Another RCT among asymptomatic patients found reduced risk of symptom development (e.g., 91% less fever development), faster viral clearance and reduced hospitalization (0 vs. 6%) (Samaha et al., 2021). Three RCTs of treatment within 5 days of symptoms onset showed an association of ivermectin with subsequently lower viral loads (Chaccour et al., 2020, Krolewiecki et al., 2020)(Biber et al., 2021), although one trial of a single dose of Ivermectin 12mg or 24mg failed to accelerate viral clearance (Mohan et al., 2021). Thus, most studies of early use of Ivermectin found efficacy. As Ivermectin is non-invasive, has low adverse effects, is low cost and there is evidence of efficacy, it is recommended for treatment of early COVID-19 cases.

Late treatment (>5 days) with Ivermectin has been assessed in multiple RCTs). There is one moderate-quality RCT comparing usual care to usual care plus ivermectin, which found no benefits (Podder et al., 2020). Another with mostly late treatment found lack of efficacy (Reis 2022). Another found a lack of benefit when started an average 5 days after symptom onset (López-Medina et al., 2021). Another trial found lack of benefits (Beltran-Gonzalez et al., 2021). However, one trial found improvements in duration of symptoms and reduced hospitalization (Shahbaznejad et al., 2021). Another RCT reported faster recovery and reduced mortality when Ivermectin was added to

HCQ/AZT/Favipiravir (Okumuş et al., 2021). Another trial found reduced symptoms and viral loads (Pott-Junior et al., 2021). One trial found lack of statistical significance for viral clearance, although there were 0 deaths in the ivermectin vs. 4 in the placebo groups; there were 100% discharges in the Ivermectin vs. 93% in the placebo groups, $p=0.045$ (Ravikirti et al., 2021). As data somewhat conflict, there is no recommendation regarding Ivermectin for treatment of late COVID-19 disease.

One low-quality study with patients of unclear symptoms duration found ivermectin to be comparable to HCQ/AZT/Ivermectin combination therapy, although there appear to have been baseline differences in oxygen saturation (Babalola et al., 2021).

There are a few studies of the use of ivermectin as a prophylactic for those close contacts exposed to COVID cases. One trial found 7.4% developed COVID compared with 58.4% of controls (Shoumann et al., 2021). Another trial assessed a combination of high-dose ivermectin (12mg/day for 7 days) and inhaled iota-carrageenan, finding 3.4% developed COVID vs. 21.4% of controls; those with moderate and severe symptoms were solely in the control group (Chahla et al., 2021). There are insufficient studies for a recommendation for prophylactic use of ivermectin, although the current studies suggest a potential for benefit and more studies are needed.

There are some RCTs treating patients of unclear symptoms duration. A comparative trial of HCQ, CQ and Ivermectin among hospitalized, severely-affected patients of unclear symptoms duration found lack of differences (Galan et al., 2021). One trial suggested no differences in duration of hospitalization (Abd-El salam et al., 2021). One trial suggested accelerated viral clearance (Babalola et al., 2021). A trial among patients of unclear severity (severely affected per abstract, mildly affected per Table) found lack of efficacy (Saxena et al., 2021).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2020 to October 2021 using the following terms: Ivermectin; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 72 articles in PubMed, 1 in Scopus, 22 in CINAHL, 711 in Cochrane Library, 7730 in Google Scholar, and 26 from other sources†. We considered for inclusion 13 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 14 from Google Scholar, and 0 from other sources. Of the 28 articles considered for inclusion, 13 randomized trials and 14 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Ivermectin for the Treatment of COVID-19 - Within 3 Days of Symptom Onset

Recommended

Ivermectin is recommended for early treatment of COVID-19, within 3 days of symptom onset. However, it is not recommended for late treatment of COVID-19 (Podder et al., 2020, Chaccour et

al., 2020, Krolewiecki et al., 2020, Niaee et al., 2020, Chowdhury et al., 2020, Hashim et al., 2020, Elgazzar et al., 2020, Ahmed et al., 2020, Alam et al., 2020, Behera et al., 2020, Cadegiani et al., 2020, Camprubí et al., 2020, Gorial et al., 2020, Heidary et al., 2020, Ortiz-Muñoz et al., 2020, Padhy et al., 2020, Rajter et al., 2020, Soto-Becerra et al., 2020).

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Outpatients with COVID-19 and less than 3 days of symptoms.

Benefits

Reduced risk of progression, shorter recovery, reduced hospitalization and faster viral clearance.

Harms

Dizziness, loss of appetite, nausea, vomiting, stomach pain, diarrhea, constipation, weakness, headache, myalgias, rash.

Frequency/Dose/Duration

Single dose of ivermectin 12 mg and doxycycline 100mg BID for 5 days (Mahmud et al., 2021).

Indications for discontinuation

Completion of course, adverse effects, intolerance

Rationale

There are many RCTs assessing ivermectin. Trials used different dose regimens (including single dose), in patients of varying symptoms duration. One metanalysis of controlled studies reported 61% reductions in mortality, time to recovery, and time to viral clearance (Marik et al., 2021).

There are several trials assessing use of ivermectin for early treatment (e.g., within 3-5 days). One high-quality trial of a combination of a single dose of ivermectin 12 mg and doxycycline 100mg BID for 5 days found 51% reduced risk of progression to serious disease and recovery averaged 2 fewer days (Mahmud et al., 2021). Another high-quality trial found a non-significant 33% reduced risk of hospitalization among those treated with weight-dependent dosing of Ivermectin (e.g., among 80-110kg, 18mg and another 18mg at 24 hrs) at 3-6 days of symptoms (Vallejos et al., 2021). Another RCT among asymptomatic patients found reduced risk of symptom development (e.g., 91% less fever development), faster viral clearance and reduced hospitalization (0 vs. 6%) (Samaha et al., 2021). Three RCTs of treatment within 5 days of symptoms onset showed an association of ivermectin with subsequently lower viral loads (Chaccour et al., 2020, Krolewiecki et al., 2020)(Administration, 2022), although one trial of a single dose of Ivermectin 12mg or 24mg failed to accelerate viral clearance (Mohan et al., 2021). Thus, most studies of early use of Ivermectin found efficacy. As Ivermectin is non-invasive, has low adverse effects, is low cost and there is evidence of efficacy, it is recommended for treatment of early COVID-19 cases. Late treatment (>3 days) with Ivermectin has been assessed in multiple RCTs). There is one moderate-quality RCT comparing usual care to usual

care plus ivermectin, which found no benefits (Podder et al., 2020). Another found a lack of benefit when started an average 5 days after symptom onset (López-Medina et al., 2021). Another trial found lack of benefits (Beltran-Gonzalez et al., 2021). However, one trial found improvements in duration of symptoms and reduced hospitalization (Shahbaznejad et al., 2021). Another RCT reported faster recovery and reduced mortality when Ivermectin was added to HCQ/AZT/Favipiravir (Okumuş et al., 2021). Another trial found reduced symptoms and viral loads (Pott-Junior et al., 2021). One trial found lack of statistical significance for viral clearance, although there were 0 deaths in the ivermectin vs. 4 in the placebo groups; there were 100% discharges in the Ivermectin vs. 93% in the placebo groups, $p=0.045$ (Ravikirti et al., 2021). As data somewhat conflict, there is no recommendation regarding Ivermectin for treatment of late COVID-19 disease. One low-quality study with patients of unclear symptoms duration found ivermectin to be comparable to HCQ/AZT/Ivermectin combination therapy, although there appear to have been baseline differences in oxygen saturation. There are a few studies of the use of ivermectin as a prophylactic for those close contacts exposed to COVID cases. One trial found 7.4% developed COVID compared with 58.4% of controls (Shoumann et al., 2021). Another trial assessed a combination of high-dose ivermectin (12mg/day for 7 days) and inhaled iota-carrageenan, finding 3.4% developed COVID vs. 21.4% of controls; those with moderate and severe symptoms were solely in the control group (Chahla et al., 2021). There are insufficient studies for a recommendation for prophylactic use of ivermectin, although the current studies suggest a potential for benefit and more studies are needed. There are some RCTs treating patients of unclear symptoms duration. A comparative trial of HCQ, CQ and Ivermectin among hospitalized, severely-affected patients of unclear symptoms duration found lack of differences. One trial suggested no differences in duration of hospitalization (Marques et al., 2022). One trial suggested accelerated viral clearance. A trial among patients of unclear severity (severely affected per abstract, mildly affected per Table) found lack of efficacy (Saxena et al., 2021).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2020 to October 2021 using the following terms: Ivermectin; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 72 articles in PubMed, 1 in Scopus, 22 in CINAHL, 711 in Cochrane Library, 7730 in Google Scholar, and 26 from other sources[†]. We considered for inclusion 13 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 14 from Google Scholar, and 0 from other sources. Of the 28 articles considered for inclusion, 13 randomized trials and 14 systematic reviews met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Ivermectin for the Treatment of COVID-19 - Prophylaxis

No Recommendation

There is no recommendation for ivermectin prophylaxis.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

Outpatients with COVID-19 and less than 5 days of symptoms.

Benefits

Reduced risk of progression, shorter recovery, reduced hospitalization and faster viral clearance.

Harms

Dizziness, loss of appetite, nausea, vomiting, stomach pain, diarrhea, constipation, weakness, headache, myalgias, rash.

Frequency/Dose/Duration

Single dose of ivermectin 12 mg and doxycycline 100mg BID for 5 days (Mahmud et al., 2021).

Indications for discontinuation

Completion of course, adverse effects, intolerance

Rationale

There are many RCTs assessing ivermectin. Trials used different dose regimens (including single dose), in patients of varying symptoms duration. One metanalysis of controlled studies reported 61% reductions in mortality, time to recovery, and time to viral clearance (Marik et al., 2021).

There are several trials assessing use of ivermectin for early treatment (e.g., within 3-5 days). One high-quality trial of a combination of a single dose of ivermectin 12 mg and doxycycline 100mg BID for 5 days found 51% reduced risk of progression to serious disease and recovery averaged 2 fewer days (Mahmud et al., 2021). Another high-quality trial found a non-significant 33% reduced risk of hospitalization among those treated with weight-dependent dosing of Ivermectin (e.g., among 80-110kg, 18mg and another 18mg at 24 hrs) at 3-6 days of symptoms (Vallejos et al., 2021). Another RCT among asymptomatic patients found reduced risk of symptom development (e.g., 91% less fever development), faster viral clearance and reduced hospitalization (0 vs. 6%) (Samaha et al., 2021). Three RCTs of treatment within 5 days of symptoms onset showed an association of ivermectin with subsequently lower viral loads (Chaccour et al., 2020, Krolewiecki et al., 2020)(Administration, 2022), although one trial of a single dose of Ivermectin 12mg or 24mg failed to accelerate viral clearance (Mohan et al., 2021). Thus, most studies of early use of Ivermectin found efficacy. As Ivermectin is non-invasive, has low adverse effects, is low cost and there is evidence of efficacy, it is recommended for treatment of early COVID-19 cases.

Late treatment (>5 days) with Ivermectin has been assessed in multiple RCTs). There is one moderate-quality RCT comparing usual care to usual care plus ivermectin, which found no benefits (Podder et al., 2020). Another found a lack of benefit when started an average 5 days after symptom onset (López-Medina et al., 2021). Another trial found lack of benefits (Beltran-Gonzalez et al., 2021). However, one trial found improvements in duration of symptoms and reduced hospitalization (Shahbaznejad et al., 2021). Another RCT reported faster recovery and reduced mortality when Ivermectin was added to HCQ/AZT/Favipiravir (Okumuş et al., 2021). Another trial found reduced symptoms and viral loads (Pott-Junior et al., 2021). One trial found lack of statistical significance for viral clearance, although there were 0 deaths in the ivermectin vs. 4 in the placebo groups; there were 100% discharges in the Ivermectin vs. 93% in the placebo groups, $p=0.045$ (Ravikirti et al., 2021). As data somewhat conflict, there is no recommendation regarding Ivermectin for treatment of late COVID-19 disease.

One low-quality study with patients of unclear symptoms duration found ivermectin to be comparable to HCQ/AZT/Ivermectin combination therapy, although there appear to have been baseline differences in oxygen saturation .

There are a few studies of the use of ivermectin as a prophylactic for those close contacts exposed to COVID cases. One trial found 7.4% developed COVID compared with 58.4% of controls (Shoumann et al., 2021). Another trial assessed a combination of high-dose ivermectin (12mg/day for 7 days) and inhaled iota-carrageenan, finding 3.4% developed COVID vs. 21.4% of controls; those with moderate and severe symptoms were solely in the control group (Chahla et al., 2021). There are insufficient studies for a recommendation for prophylactic use of ivermectin, although the current studies suggest a potential for benefit and more studies are needed.

There are some RCTs treating patients of unclear symptom duration. A comparative trial of HCQ, CQ and Ivermectin among hospitalized, severely-affected patients of unclear symptoms duration found lack of differences . One trial suggested no differences in duration of hospitalization (Marques et al., 2022). One trial suggested accelerated viral clearance . A trial among patients of unclear severity (severely affected per abstract, mildly affected per Table) found lack of efficacy (Saxena et al., 2021).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2020 to October 2021 using the following terms: Ivermectin; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 72 articles in PubMed, 1 in Scopus, 22 in CINAHL, 711 in Cochrane Library, 7730 in Google Scholar, and 26 from other sources†. We considered for inclusion 13 from PubMed, 0 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 14 from Google Scholar, and 0 from other sources. Of the 28 articles considered for inclusion, 13 randomized trials and 14 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.19. CONVALESCENT COVID-19 ANTIBODIES

Convalescent COVID-19 antibodies have been used to treat COVID-19 (175) (198) (199) (200) (201) (202) (203) (204) (205) (206) (207) (208) (209) (210) (211) (212) (213) (214) (215) (216).

Convalescent COVID-19 Antibodies for Treatment of COVID-19

Not Recommended

The use of convalescent antibodies is not recommended for the treatment of patients with COVID-19.

Strength of evidence Moderately Not Recommended, Evidence (B)

Level of confidence Moderate

Rationale

There are many quality RCTs and nearly all reported lack of efficacy (Li et al., 2020, Simonovich et al., 2020, AlQahtani et al., 2020, Agarwal et al., 2020)(O'Donnell et al., 2021, Sekine et al., 2021, Devos et al., 2021, Horby et al., 2021, Gharbharan et al., 2020, Bennett-Guerrero et al., 2021, Balcells et al., 2021)(Bégin et al., 2021)(Körper et al., 2021)(Avendaño-Solà et al., 2020)(Duan et al., 2020)(Chen et al., 2021)(Gharbharan et al., 2021)(Tabarsi et al., 2021). One moderate-quality trial suggested potential reduction in the need for mechanical ventilation (Avendaño-Solà et al., 2020). A pilot study suggested efficacy (Bajpai et al., 2021).

Convalescent antibodies are invasive, have adverse effects, and are costly. Nearly all quality data suggest they are ineffective for treatment of COVID-19 and thus they are not recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Convalescent Antibodies; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 279 articles in PubMed, 10,212 in Scopus, 18 in CINAHL, 45 in Cochrane Library, 23,060 in Google Scholar, and 0 from other sources†. We considered for inclusion 17 from PubMed, 0 from Scopus, 0 from CINAHL, 1 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 20 articles considered for inclusion, 10 randomized trials and 10 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.20. GLUCOCORTICOSTEROIDS

Glucocorticosteroids for the Treatment of COVID-19

Recommended

Glucocorticosteroids are recommended for the treatment of COVID-19. There are other indications for use that may occur in the context of treatment of COVID-19 (e.g., asthma, COPD).

Strength of evidence Moderately Recommended, Evidence (B)

Level of confidence Moderate

Indications

Hospitalized patients with moderate or severe COVID-19. Especially effective reportedly for those critically ill on ventilators, requiring supplemental oxygen and/or cardiovascular support. Use is also indicated in outpatients at risk of hospitalization which includes the use of inhaled steroids.

Benefits

A meta-analysis estimated a 36% reduction in mortality with dexamethasone, 31% reduction with hydrocortisone, and 9% reduction with methylprednisolone (Sterne et al., 2020). One trial estimated a reduced mortality by 20% if requiring supplemental oxygen and 35% if ventilated. A reduced number of ventilator days has also been reported. Inhaled steroid reportedly reduce illness duration by 3 days among those at higher risk (Yu et al., 2021).

Harms

Hyperglycemia, risk of secondary infection, higher blood pressure.

Frequency/Dose/Duration

Different treatments have been used. There are no comparative trials and optimal dosing is somewhat unclear. Medications and doses used have included:

- Dexamethasone 6 mg PO or IV QD x 10 days or until discharge (or equivalent dose)s.
- Hydrocortisone 50mg or 100mg every 6 hours (Angus et al., 2020)

Two trials have suggested efficacy of budesonide 2 puffs BID, with use including outpatients (Ramakrishnan et al., 2021, Yu et al., 2021).

Indications for discontinuation

Completion of a course, intolerance, adverse effect.

Rationale

There are multiple RCTs, with all larger sized studies suggesting efficacy (Angus et al., 2020, Tomazini et al., 2020, Horby et al., 2020, Edalatifard et al., 2020, Jeronimo et al., 2020), while the negative studies were small sample sizes and appear tended to trend towards efficacy (Abd-El Salam et al., 2020, Dequin et al., 2020)(Tang et al., 2021, Jamaati et al., 2021, Ghanei et al., 2021). A meta-analysis estimated a 36% reduction in mortality with dexamethasone, 31% reduction with hydrocortisone, and 9% reduction with methylprednisolone (Sterne et al., 2020). A comparative trial suggested superiority of methylprednisolone 2mg/kg/day to dexamethasone 6mg/kg/day, which is a lower steroid dose of the methylprednisolone (Ranjbar et al., 2021). Two studies of inhaled budesonide suggest modest efficacy (Yu et al., 2021, Ramakrishnan et al., 2021), with one suggesting shortening of recovery by ~3 days (Yu et al., 2021). A comparative trial found dexamethasone superior to tocilizumab (Rashad et al., 2021).

A large RCT found mortality reductions with dexamethasone (Horby et al., 2020, Oxford, 2020, Ledford, 2020). An RCT found a 65% increase in ventilator-free days from 4.0 to 6.6 days over a 28-day period, although there was no difference in mortality (Tomazini et al., 2020). Another RCT found superiority of glucocorticosteroid (Angus et al., 2020). Two RCTs of modest size found no significant benefits, but appear underpowered (Dequin et al., 2020, Abd-El Salam et al., 2020). Another negative study used a low dose of hydrocortisone (Dequin et al., 2020). Because glucocorticosteroids have moderate adverse effects, low costs, and have significant efficacy in reducing mortality based on meta-analyses, they are moderately recommended for treatment of COVID-19.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Glucocorticoids, Glucocorticoid Steroid, Prednisone, Dexamethasone, Hydrocortisone; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 364 articles in PubMed, 939 in Scopus, 32 in CINAHL, 20 in Cochrane Library, 11980 in Google Scholar, and 0 from other sources[†]. We considered for inclusion 30 from PubMed, 4 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 5 from Google Scholar, and 0 from other sources. Of the 39 articles considered for inclusion, 12 randomized trials and 8 systematic reviews met the inclusion criteria. There were no exclusion criteria.

[†] The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.21. INTERFERON BETA-1B

Interferon beta-1b has been used both as sole therapy and combination therapy for the treatment of patients with COVID-19 (217) (118).

Interferon Beta-1b for the Treatment of COVID-19 - Combination Therapy

Recommended

Adjunctive use of interferon beta-1b is recommended for the treatment of selected patients with COVID-19.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Adjunctive use with lopinavir-ritonavir and ribavirin in moderately and severely affected patients with COVID-19 (Hung et al., 2020). Evidence suggests better efficacy if administered within 7 days of symptom onset; after 7 days, data suggest no differences between this combination therapy and lopinavir-ritonavir (Hung et al., 2020).

Benefits

Faster symptom resolution, viral clearance, and hospital discharge. Reduced need for a ventilator or ICU stay.

Harms

Nausea, diarrhea, hepatitis.

Frequency/Dose/Duration

Two successful trials utilized sole therapy with interferon beta-1b 250ug SQ QOD for 2 weeks (Rahmani et al., 2020). The combination regimen used successfully for the treatment of COVID-19 is lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days (Hung et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

Two successful moderate-quality trials utilized sole therapy with interferon beta-1b and one study found accelerated clinical improvement and a non-statistically significant reduction in death by 67% at 1-month (Rahmani et al., 2020). The second trial found comparable results to the other RCT with

faster clinical improvement (9 vs 11 days), fewer adverse events, and ~67% reduction in mortality (6.1 vs. 18.2%) when compared with treatment with the control group (lopinavir-ritonavir/HCQ or atazanavir/ritonavir/HCQ) (Rahmani et al., 2020). One open-label RCT found combination therapy of lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days to be superior to lopinavir-ritonavir (Hung et al., 2020). Another three-arm trial found faster clinical recovery and non-significant trends in mortality compared to control (mortality 20% for beta1a vs. 30% for beta1b vs. 45% controls) (Darazam, 2021). However, one RCT found a lack of efficacy (WHO, 2020). Based on the two moderate-quality RCTs showing considerable evidence of efficacy, stand-alone treatment with interferon beta-1b is moderately recommended.

A moderate-quality RCT found a lack of efficacy for combined favipavir with interferon beta-1b compared with HCQ for moderate to severe COVID-19 pneumonia patients (Khamis et al., 2021).

Based on one trial with demonstrated efficacy, the regimen of triple-combination therapy using lopinavir, ritonavir, ribavirin, and interferon beta-1b is recommended (Hung et al., 2020), although it should be noted that it is possible that the only medication effective in the combination therapy is interferon beta-1b.

Other interferons are being investigated. One successful trial used a different interferon in a Phase 2 trial that was nebulized interferon-1a (SNG001) (Monk et al., 2021). A trial with interferon beta-1a when added to (lopinavir-ritonavir/HCQ or atazanavir/ritonavir/HCQ) found earlier 14-day hospital discharge rates (67% vs. 44%) (Davoudi-Monfared et al., 2020). A trial on interferon-kappa plus TFF2 and including many potentially active cointerventions found reduced viral RNA (Fu et al., 2020).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Interferon beta-1b; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 44 articles in PubMed, 2,504 in Scopus, 8 in CINAHL, 5 in Cochrane Library, 3,280 in Google Scholar, and 0 from other sources†. We considered for inclusion 1 from PubMed, 13 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 16 articles considered for inclusion, 1 randomized trial and 15 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Interferon Beta-1b for the Treatment of COVID-19 - Stand-alone Treatment

Recommended

Adjunctive use of interferon beta-1b is recommended for the treatment of selected patients with COVID-19.

Strength of evidence Moderately Recommended, Evidence (B)

Level of confidence Low

Indications

Adjunctive use with lopinavir-ritonavir and ribavirin in moderately and severely affected patients with COVID-19 (Hung et al., 2020). Evidence suggests better efficacy if administered within 7 days of symptom onset; after 7 days, data suggest no differences between this combination therapy and lopinavir-ritonavir (Hung et al., 2020).

Benefits

Faster symptom resolution, viral clearance, and hospital discharge. Reduced need for a ventilator or ICU stay.

Harms

Nausea, diarrhea, hepatitis.

Frequency/Dose/Duration

Two successful trials utilized sole therapy with interferon beta-1b 250ug SQ QOD for 2 weeks (Rahmani et al., 2020). The combination regimen used successfully for the treatment of COVID-19 is lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days (Hung et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

Two successful moderate-quality trials utilized sole therapy with interferon beta-1b and one study found accelerated clinical improvement and a non-statistically significant reduction in death by 67% at 1-month (Rahmani et al., 2020). The second trial found comparable results to the other RCT with faster clinical improvement (9 vs 11 days), fewer adverse events, and ~67% reduction in mortality (6.1 vs. 18.2%) when compared with treatment with the control group (lopinavir-ritonavir/HCQ or atazanavir/ritonavir/HCQ) (Rahmani et al., 2020). One open-label RCT found combination therapy of lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days to be superior to lopinavir-ritonavir (Hung et al., 2020). Another three-arm trial found faster clinical recovery and non-significant trends in mortality compared to control (mortality 20% for beta1a vs. 30% for beta1b vs. 45% controls) (Darazam, 2021). However, one RCT found a lack of efficacy (WHO, 2020). Based on the two moderate-quality RCTs showing considerable evidence of efficacy, stand-alone treatment with interferon beta-1b is moderately recommended.

A moderate-quality RCT found a lack of efficacy for combined favipavir with interferon beta-1b compared with HCQ for moderate to severe COVID-19 pneumonia patients (Khamis et al., 2021).

Based on one trial with demonstrated efficacy, the regimen of triple-combination therapy using lopinavir, ritonavir, ribavirin, and interferon beta-1b is recommended (Hung et al., 2020), although it should be noted that it is possible that the only medication effective in the combination therapy is interferon beta-1b.

Other interferons are being investigated. One successful trial used a different interferon in a Phase 2 trial that was nebulized interferon-1a (SNG001) (Monk et al., 2021). A trial with interferon beta-1a when added to (lopinavir-ritonavir/HCQ or atazanavir/ritonavir/HCQ) found earlier 14-day hospital discharge rates (67% vs. 44%) (Davoudi-Monfared et al., 2020). A trial on interferon-kappa plus TFF2 and including many potentially active cointerventions found reduced viral RNA (Fu et al., 2020).

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Interferon beta-1b; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 44 articles in PubMed, 2,504 in Scopus, 8 in CINAHL, 5 in Cochrane Library, 3,280 in Google Scholar, and 0 from other sources†. We considered for inclusion 1 from PubMed, 13 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 16 articles considered for inclusion, 1 randomized trial and 15 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.22. RIBAVIRIN

Ribavirin has been used to treat patients with COVID-19 (218) (219) (220) (221).

Ribavirin for the Treatment of COVID-19 - Combination Therapy

Recommended

Adjunctive use of ribavirin is recommended for the treatment of selected patients with COVID-19.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Adjunctive use with lopinavir-ritonavir and interferon beta-1b in moderately and severely affected patients with COVID-19 (Hung et al., 2020). Evidence suggests better efficacy if administered within 7 days of symptom onset; after 7 days, data suggest no differences between this combination therapy and lopinavir-ritonavir (Hung et al., 2020).

Benefits

Faster symptom resolution, viral clearance, and hospital discharge. Reduced need for a ventilator or ICU stay.

Harms

Nausea, diarrhea, hepatitis.

Frequency/Dose/Duration

The regimen used for the treatment of COVID-19 is lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days (Hung et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

One open-label RCT found combination therapy of lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days to be superior to lopinavir-ritonavir (Hung et al., 2020). Two other RCTs were underpowered for meaningful clinical differences (Huang et al., 2020, Abbaspour Kasgari et al., 2020). One trial of adjunctive use also did not define the timing of treatment vs. symptoms onset (Panda et al., 2021).

Based on the one moderate-quality RCT showing evidence of efficacy, the regimen of triple-combination therapy using lopinavir, ritonavir, ribavirin, and interferon beta-1b is recommended (Hung et al., 2020). However, there is no quality evidence demonstrating efficacy and thus no recommendation for stand-alone treatment with ribavirin.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Ribavirin; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 103 articles in PubMed, 7,064 in Scopus, 10 in CINAHL, 977 in Cochrane Library, 16,570 in Google Scholar, and 0 from other sources†. We considered for inclusion 7 from PubMed, 1 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 8 articles considered for inclusion, 1 randomized trial and 7 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

Ribavirin for the Treatment of COVID-19 - Stand-alone Treatment

No Recommendation

Adjunctive use of ribavirin is recommended for the treatment of selected patients with COVID-19.

Strength of evidence No Recommendation, Insufficient Evidence (I)

Level of confidence Low

Indications

Adjunctive use with lopinavir-ritonavir and interferon beta-1b in moderately and severely affected patients with COVID-19 (Hung et al., 2020). Evidence suggests better efficacy if administered within 7 days of symptom onset; after 7 days, data suggest no differences between this combination therapy and lopinavir-ritonavir (Hung et al., 2020).

Benefits

Faster symptom resolution, viral clearance, and hospital discharge. Reduced need for a ventilator or ICU stay.

Harms

Nausea, diarrhea, hepatitis.

Frequency/Dose/Duration

The regimen used for the treatment of COVID-19 is lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days (Hung et al., 2020).

Indications for discontinuation

Completion of a course, intolerance, adverse effect, prolongation of QT interval.

Rationale

One open-label RCT found combination therapy of lopinavir 400mg, ritonavir 100mg every 12hrs, ribavirin 400mg every 12hrs, plus 3 doses of 8M IU interferon beta-1b on alternate days to be superior to lopinavir-ritonavir (Hung et al., 2020). Two other RCTs were underpowered for

meaningful clinical differences (Huang et al., 2020, Abbaspour Kasgari et al., 2020). One trial of adjunctive use also did not define the timing of treatment vs. symptoms onset (Panda et al., 2021).

Based on the one moderate-quality RCT showing evidence of efficacy, the regimen of triple-combination therapy using lopinavir, ritonavir, ribavirin, and interferon beta-1b is recommended (Hung et al., 2020). However, there is no quality evidence demonstrating efficacy and thus no recommendation for stand-alone treatment with ribavirin.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Ribavirin; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 103 articles in PubMed, 7,064 in Scopus, 10 in CINAHL, 977 in Cochrane Library, 16,570 in Google Scholar, and 0 from other sources†. We considered for inclusion 7 from PubMed, 1 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 8 articles considered for inclusion, 1 randomized trial and 7 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.23. ZINC

Zinc serum levels have been found to be low in those with more severe COVID-19 disease (222) (223) (224). Zinc supplementation has been used typically as adjunctive treatment to reduce severity of COVID-19 (225) (226).

Zinc for the Treatment of COVID-19

Recommended

Zinc is recommended for potential prevention of more severe disease as well as for the treatment of patients with COVID-19.

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Low

Indications

Ongoing use during the epidemic, as well as for mild, moderate, and severe COVID-19 disease. Also especially recommended for those with zinc deficiency.

Benefits

Potential to reduce disease severity.

Harms

Negligible.

Frequency/Dose/Duration

10-15 mg/day (>100% Recommended Daily Allowance).

Indications for discontinuation

After cessation of the epidemic.

Rationale

There are multiple RCTs, but few test the value of zinc alone (Derwand et al., 2020, Skalny et al., 2020, Finzi, 2020, Carlucci et al., 2020). One pilot RCT suggested a non-significant inverse correlation between zinc levels and oxygenation requirements (Patel et al., 2021). There is one low-quality study suggesting lack of efficacy of zinc added to HCQ (Abd-Elsalam et al., 2020) and another low-quality trial found lack of efficacy of high-dose zinc and ascorbic acid added to usual care (Thomas et al., 2021). However, one study of HCQ, AZT, and zinc suggested earlier treatment resulted in 84% lower risk of hospitalization and lower risk of death among patients treated by ~day 4 (Derwand et al., 2020). A large-scale pre/post intervention study showed that adjunctive use of zinc to hydroxychloroquine was associated with a 44–49% decreased need for ventilation, admission to the ICU, mortality, or transfer to hospice, and increased the frequency of being discharged home (Carlucci et al., 2020). This is supported by evidence that hydroxy/chloroquine are zinc ionophores, which increase intracellular zinc and reduce or prevent viral replication in laboratory studies (te Velthuis et al., 2010, Xue et al., 2014).

Zinc supplementation has negligible adverse effects and has been associated with improved outcomes in non-randomized studies; thus, it is recommended with insufficient evidence.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Zinc, Zinc Compounds; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 17 articles in PubMed, 20 in Scopus, 6 in CINAHL, 114 in Cochrane Library, 17000 in Google Scholar, and 0 from other sources†. We considered for inclusion 2 from PubMed, 1 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 6 articles considered for inclusion, 1

randomized trial and 3 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

8.24. VITAMIN D

Vitamin D levels have been low in those with more severe COVID-19 disease and supplementation has been used for the treatment of patients with COVID-19 (227) (228) (229) (230) (231) (232) (233) (234) (235) (236) (237) (238) (239) (240) (241) (242) (243) (244). It has also been used in patients with COVID-19 to maintain bone health.

Vitamin D for the Treatment of COVID-19

Recommended

Vitamin D is recommended for potential prevention of more severe disease as well as for the treatment of patients with COVID-19.

Strength of evidence Recommended, Evidence (C)

Level of confidence Low

Indications

Ongoing use during the epidemic, as well as for mild, moderate, and severe COVID-19 disease. High-dose use may be considered for those with onset of COVID-19 disease. Also recommended for those with vitamin D deficiency and/or risks for deficiency.

Benefits

Potential to reduce disease severity

Harms

Negligible

Frequency/Dose/Duration

A moderate-quality trial utilized calcifediol 0.532mg on day 1, 0.266mg days 3 and 7 and weekly in addition to HCQ+AZT until hospital discharge (Entrenas Castillo et al., 2020). Other daily dosing used

among healthy individuals at risk include 600 IU/day for up to 70 years of age and 800 IU/day for those over 70 years of age (>100% Recommended Daily Allowance).

Indications for discontinuation

After cessation of the epidemic.

Rationale

A moderate-quality RCT used calcifediol compared with no calcifediol in addition to HCQ+AZT until hospital discharge and found a 96% reduction in risk of needing an ICU stay (Entrenas Castillo et al., 2020). Another RCT for treatment of asymptomatic or mildly symptomatic but vitamin D deficient individuals treated with vitamin D supplementation cleared virus sooner and with reduced fibrinogen levels (Rastogi et al., 2020). Another RCT found a 32% reduction in symptoms duration associated with vitamin D 5000 IU vs. 1000 IU for 2 weeks (Sabico et al., 2021). One RCT found lack of efficacy using only one administration of 200,000 IU, although the risk of mechanical ventilation trended towards reduction by 51% ($p=0.09$) (Murai et al., 2020). Vitamin D levels have been strongly correlated with COVID-19 disease severity (Alipio, 2020, Lau et al., 2020, D'Avolio et al., 2020), with a reported ~8-fold risk of a severe outcome and ~20-fold risk of a critical outcome among those with low vitamin D levels (Alipio, 2020).

Vitamin D supplementation has negligible adverse effects, especially over shorter periods of time, and low vitamin D levels have been strongly associated with worse outcomes in non-randomized studies. Vitamin D levels also fall with illness status affecting bone health. Thus, vitamin D supplementation is recommended.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to October 2021 using the following terms: Vitamin D, Vitamin D Supplementation; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 185 articles in PubMed, 14,869 in Scopus, 68 in CINAHL, 7 in Cochrane Library, 13,400 in Google Scholar, and 0 from other sources†. We considered for inclusion 2 from PubMed, 3 from Scopus, 0 from CINAHL, 2 from Cochrane Library, 2 from Google Scholar, and 0 from other sources. Of the 43,843 articles considered for inclusion, 2 randomized trials and 7 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

9. REHABILITATION

9.1. OVERVIEW

Although most patients with COVID-19 completely recover, some cases may experience a multitude of disorders (731). It is beyond the scope of this guideline to address every possible presentation, combination, and permutation (190). Indeed, it is arguably impossible to do so. Instead, this guideline addresses what currently appear to be the most common conditions needing rehabilitative services after COVID-19. This also may suggest a framework for approaching treatment of less common presentations.

For simplicity, clarity, and consistency with other diagnoses and the general medical literature, this review defines symptoms lasting less than 1 month as *acute*, from 1–3 months as *subacute*, and more than 3 months as *chronic*. Some of the alternate terms for these conditions include “ongoing symptomatic,” “post-COVID syndrome” (565), “post-acute sequelae of COVID,” and “long COVID.”

The severity of the COVID-19 infection has been associated with the risk of long-term symptoms and impairments (247). For example, approximately two-thirds of outpatients diagnosed with COVID-19 return to normal health by the fourth week (732). In contrast, of those who were evaluated in an emergency department (66% hospitalized), 50.9% developed chronic COVID-19 symptoms (733). Yet, a mild case does not preclude development of chronic COVID-19 symptoms. The comparatively large numbers of mildly affected patients likely mean that most patients with chronic symptoms will be found in this group, despite the higher risk among those who are more severely affected.

Evidence also suggests that symptoms improve over time. Overall, 5–51% of patients have symptoms persisting up to 12 weeks (566,734,735,736), whereas 2–15% have symptoms beyond 12 weeks after onset (733,734,736,737,738). Long-term symptoms have wide-ranging estimates of prevalence and include fatigue (17–98%) (566,733,734,735,736,739,740,741), dyspnea (17–93%) (566,733,734,735,736,739,740,741), cough (29–43%), chest pain (44–88%) (566,733,734,735,740), back pain, muscle pain, and headache (38–91%) (733). Cognitive changes, such as impaired memory, concentration, and multitasking ability, are also reportedly common. Risk factors for chronic COVID-19 beyond severity of the initial disease appear to include increased age, having more comorbidities, and psychological disorders (736,742).

Acute mental health disorders are common and reportedly affect 55% of those having visited an emergency department (75% were hospitalized) in the first month (743). New-onset psychiatric illness was reported in 5.8% [801]. Of these, 4.7% were anxiety disorders and 2% were depression (744). One report noted persistence of post-traumatic stress among survivors (745). Another reported PTSD symptoms related to illness at 4–8 weeks after discharge among 46.9% of ICU survivors and 23.5% of ward survivors (247).

Some rehabilitation protocols are heavily multidisciplinary, reportedly including pulmonologists, physiatrists, neurologists, cardiologists, physical therapists, occupational therapists, psychologists, neuropsychologists, psychiatrists, speech therapists, and nutritionists (746,747)(748). Telemedicine has been used for rehabilitation of COVID-19 patients (747,749). There are no quality trials to assess the various disciplines on rehabilitation teams, comparative trials of different treatment regimens, and/or efficacy of telemedicine approaches.

9.2. PULMONARY REHABILITATION

Dyspnea is typically the presenting complaint for emergency and hospitalized treatment. However, dyspnea has been shown to persist into many chronic COVID-19 case histories (245,246) (247)(248). The most common spirometric abnormalities after initial recovery are reduced diffusion capacity and

restrictive ventilatory defects (249,250). Risk and severity of spirometric abnormalities are correlated with COVID-19 severity (250).

Pulmonary rehabilitation is used for COVID-19 (251) and has been shown to be successful for functional improvements in individuals with non-COVID-related pulmonary deficits (252,253), including those from pneumonia (254), interstitial lung disease (255), and SARS (256). It commonly includes behavioral components (257). Consensus guidelines have also been produced (248).

Pulmonary Rehabilitation for Treatment of Pulmonary Problems Related to COVID-19

Recommended

Pulmonary rehabilitation is recommended for the treatment of pulmonary problems related to COVID-19.

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Moderate

Indications

Indicated for COVID-19 affected patients with pulmonary dysfunction and/or dyspnea, especially when combined with activity reductions or exercise intolerances attributed to the infection's pulmonary complications. Important targets are gaps between current function and job demands (esp. objectively measured). Earlier institution of exercises, as tolerated, is advised to counter the debility associated with the disease (Spruit et al., 2013). Careful cardiac evaluation (e.g., 24-hr ECG, echocardiogram, exercise testing, MRI (Barker-Davies et al., 2020)) is also indicated due to the probability of cardiac abnormalities.

Benefits

Improved pulmonary function, maximum ventilation, health-related quality of life, emotional involvement in everyday life, activity levels, 6-minute walk distance, peak workload, activity tolerance and stamina/endurance.

Harms

As fatalities in the recovery period have been thought to be due to arrhythmias, a careful assessment of cardiac involvement is advised to help guide the onset and progression of exercises. Those with evidence of thrombotic tendencies and/or multi-system involvement may have greater risk of harm from aggressive exercise regimens.

Frequency/Dose/Duration

An individualized but interdisciplinary treatment regimen is usually formulated based on a comprehensive baseline assessment (Dowman et al., 2021, Spruit et al., 2020). Careful cardiac evaluation (e.g., 24-hr ECG, echocardiogram, exercise testing, MRI (Barker-Davies et al., 2020)) is also indicated due to the probability of cardiac abnormalities, which may result in a recommendation to delay onset of exercises and/or slow the rate of progression. While program components include education, exercise training, and behavior change to "promote the long-term adherence to health-enhancing behaviours" (Spruit et al., 2020), exercise training is the central

component, and is usually either walking or cycling. One consensus statement recommended beginning at not more than 3 METS, especially when supplemental oxygen is needed (Barker-Davies et al., 2020). Another review suggested an exercise regimen of 18-60 min at 55–80% of VO₂Max or 60–80% of heart rate maximum, 1–3 times per week (Alawna et al., 2020). Program duration is typically at least 4 weeks.

Indications for discontinuation

Completion of a treatment course, noncompliance, reaching a plateau in recovery.

Rationale

There is one low-quality pilot study suggesting efficacy for treatment of COVID-19 patients (Liu et al., 2020), but no quality trials. There are many trials documenting efficacy for other pulmonary conditions (McCarthy et al., 2015, Hill, 2006, Cheng et al., 2018, Dowman et al., 2021, Lau et al., 2005). Pulmonary rehabilitation has negligible adverse effects, is moderate to high cost depending on number of treatments and durations required, and is recommended for patients meeting indications.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to February 2021 using the following terms: rehabilitation; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 112 articles in PubMed, 4,699 in Scopus, 3 in CINAHL, 13 in Cochrane Library, 34,300 in Google Scholar, and 0 from other sources†. We considered for inclusion 14 from PubMed, 4 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 18 articles considered for inclusion, 1 randomized trial and 8 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

9.3. CARDIAC REHABILITATION

Cardiomyopathy, cardiac muscle damage, and arrhythmias have been reported to affect 30–78% of patients (258), and cardiac problems contribute to COVID-19 fatalities (258,259,260,261). Vascular inflammation, hypotension, and direct muscle damage are all potential mechanisms (262,263). The probability of cardiac problems is correlated with the severity of the COVID-19 infection, including cardiac biomarkers (e.g., troponin) and numbers of comorbidities (262,263), although ongoing, subclinical cardiac problems have been detected among recovered patients (261,264).

Cardiac rehabilitation is used for COVID-19 (265) and has been shown to be successful for functional improvements in individuals with non-COVID-related cardiac deficits (266,267,268,269,270), including those from myocardial infarction (262), as well as quality-of-life measures.

Cardiac Rehabilitation for Treatment of Cardiac Problems Related to COVID-19

Recommended

Cardiac rehabilitation is recommended for the treatment of cardiac problems related to COVID-19.

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Moderate

Indications

Indicated for COVID-19 affected patients with cardiac dysfunction and/or dyspnea, especially when combined with activity reductions or exercise intolerances attributed to the infection's cardiac complications. A consensus statement advises waiting 2–3 weeks after cessation of COVID-related symptoms to start exercise (Barker-Davies et al., 2020), although there is no quality evidence to support the expert consensus. Important targets are gaps between current function and job demands (esp. objectively measured).

Benefits

Improved cardiac function, health-related quality of life, 6-minute walk test, time to perform 10 sit-to-stands, emotional involvement in everyday life, activity levels, activity tolerance, and stamina/endurance (Piquet et al., 2021).

Harms

As fatalities in the recovery period have been thought to be due to arrhythmias, a careful assessment of cardiac involvement is advised to help guide the onset and progression of exercises. Those with evidence of thrombotic tendencies, and/or multi-system involvement may have greater risk of harm from aggressive exercise regimens.

Frequency/Dose/Duration

An individualized but multidisciplinary treatment regimen is usually formulated based on a comprehensive baseline assessment (Hevey et al., 2003, Price et al., 2016). Careful cardiac evaluation (e.g., 24-hr ECG, echocardiogram, exercise testing, MRI (Barker-Davies et al., 2020)) is indicated due to the probability of cardiac abnormalities, which may result in a recommendation to delay onset of exercises and/or slow the rate of progression. Program components typically include education, aerobic exercise training, strength/resistance training, and psychological factors (Linden, 2000). Exercise training is the central component. Aerobic exercise is usually either walking or cycling. Strength training is another component thought to be important in cardiac rehabilitation (Price et al., 2016). A slowed and cautious progression may be indicated in COVID patients due to the underlying cardiac disease, and tailoring regarding arrhythmias and monitoring for exercise-induced arrhythmias has been recommended (Price et al., 2016). Program duration is typically at least 4 weeks.

High-demand occupations may be analogized to sports, where a consensus recommendation is for resumption of sports if: (1) left ventricular systolic function is normal, (2) serum biomarkers of cardiac injury are normal, (3) absence of relevant cardiac arrhythmias on 24-hr monitoring, and (4) absence of relevant cardiac arrhythmias on 24-hr monitoring on exercise testing (Barker-Davies et al., 2020).

Indications for discontinuation

Completion of a treatment course, noncompliance, reaching a plateau in recovery.

Rationale

There is one low-quality pilot study suggesting efficacy for treatment of COVID-19 patients (Liu et al., 2020), but no quality trials. There are many trials documenting efficacy for other pulmonary conditions. Cardiac rehabilitation has negligible adverse effects, is moderate to high cost depending on numbers of treatments and durations required, and is recommended for patients meeting indications.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to February 2021 using the following terms: rehabilitation; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 112 articles in PubMed, 4,699 in Scopus, 3 in CINAHL, 13 in Cochrane Library, 34,300 in Google Scholar, and 0 from other sources†. We considered for inclusion 14 from PubMed, 4 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 18 articles considered for inclusion, 1 randomized trial and 8 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

9.4. EXERCISE THERAPY

Research has supported rehabilitation for hospital-associated deconditioning prior to the COVID pandemic (271,272,273). Early mobilization of COVID-19 patients has been encouraged, and yet others suggest delaying until after the acute COVID-related symptoms have been resolved for 2-3 weeks (274)(275). Early therapy has also been used in the ICU and pre-discharge for COVID patients (276,277,278). A review of physical therapy suggests that there will eventually be efficacy, but currently the available literature is sparse and mostly low quality (279)(280).

For those with fibromyalgia, please refer to the [ACOEM Chronic Pain Guideline](#). Also consider chronic fatigue syndrome and myalgic encephalomyelitis (281).

Exercise Therapy for Physical Debility and/or Chronic Fatigue Associated with COVID-19

Recommended

A titrated return to physical activity/exercise therapy is recommended for the treatment of physical debility and/or chronic fatigue associated with COVID-19 (Vickory et al., 2021).

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Moderate

Indications

Indicated for COVID-19 affected patients with debility and/or chronic fatigue attributed to the COVID-19. Important targets are gaps between current function and job demands (esp. objectively measured). Baseline testing should indicate the area(s) of deficits (e.g., 6-min walk test; sit to stand; leg strength; grip strength). Rehabilitation should target, measure, and track progress for those specific areas.

Benefits

Improved distance walked, strength, functional gains, ability to perform ADLs independently, return to work.

Harms

As fatalities in the recovery period have been thought to be due to arrhythmias, a careful assessment of cardiac involvement is advised to help guide the onset and progression of exercises. Those with evidence of thrombotic tendencies and/or multi-system involvement may have greater risk of harm from aggressive exercise regimens.

Frequency/Dose/Duration

A multidisciplinary approach may be beneficial (e.g., physical therapy, occupational therapy, medical, psychology). Generally, sets of appointments are ordered (e.g., 6-8). Two to three appointments per week plus a home exercise program are normally prescribed. Those with marked deficits may benefit from more intensive regimens (e.g., 5 times/week). Aerobic and strengthening exercises are normally prescribed. Some exercises are ideally repeated exertions that directly target specific deficits (e.g., sit to stand or walking endurance) and should be tracked and easily increased as indicated. When there is a lack of further improvement, the course of treatment should be discontinued. Web-based programs are also possible.

Indications for discontinuation

Completion of course of treatment, noncompliance, reaching a plateau in recovery.

Rationale

There are no quality trials of outpatient-based exercise therapy for the treatment of physical debility and/or chronic fatigue attributed to COVID. Titrated return to activity and exercise programs have

negligible adverse effects, is moderate to high cost depending on numbers of treatments required, and is recommended for patients meeting indications.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to February 2021 using the following terms: rehabilitation; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 112 articles in PubMed, 4,699 in Scopus, 3 in CINAHL, 13 in Cochrane Library, 34,300 in Google Scholar, and 0 from other sources†. We considered for inclusion 14 from PubMed, 4 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from Google Scholar, and 0 from other sources. Of the 18 articles considered for inclusion, 1 randomized trial and 8 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

9.5. MEMORY AND COGNITION

Memory issues are also potentially problematic for some workers who have had COVID (282,283,284,285,286,287,288,289,290,291,292,293,294). One report noted new or worsened short-term memory problems at 4–8 weeks after discharge among 18.8% of ICU patients and 17.6% of ward patients (247), yet that same study found strong relationships for some other data such as COVID severity for breathlessness and any PTSD symptoms related to illness. It is recommended that these problems be evaluated and treated (295,296,297)(298,299). Cognitive rehabilitation has been successfully used for various infectious disease complications, especially for HIV (300) and severe malaria (301). One guideline recommended cognitive screening, assessment of contributing other conditions, imaging, testing, and consensus statements on treatment options (302).

Cognitive Rehabilitation for Treatment of Cognitive Problems Related to COVID-19

Recommended

Cognitive rehabilitation is recommended for the treatment of cognitive problems related to COVID-19.

Strength of evidence Recommended, Insufficient Evidence (I)

Level of confidence Moderate

Indications

Indicated for COVID-19 affected patients with evidence of ongoing cognitive dysfunction attributed to the infection without trending towards rapid resolution. Important targets are gaps between current function and job demands (esp. objectively measured). Screening for cognitive function should be performed. Testing should indicate the area(s) of deficits and the rehabilitation should target, measure, and track progress for those specific areas.

Benefits

Improved memory and executive functions.

Harms

Negligible

Frequency/Dose/Duration

Generally, sets of appointments are ordered (e.g., 6-8), most commonly with psychology, neuropsychology and potentially speech pathology. Depending on the severity, more intensive regimens may be indicated, e.g., in acute inpatient stroke patients, daily regimens of 30min/day for 4 weeks have been used, but likely would only be indicated for the most severely affected COVID patients. Objective improvement should be tracked. When there is a lack of further improvement, the course of treatment should be discontinued and/or re-evaluated and changed to a more effective approach (e.g., addressing a different aspect of cognitive function). Web-based programs and virtual reality (Faria et al., 2016, Bunketorp-Käll et al., 2017, Cho et al., 2019) are also possible. There is some evidence in stroke patients that combining cognitive rehabilitation with aerobic exercise results in superior outcomes (Yeh et al., 2019).

Indications for discontinuation

Completion of course of treatment, noncompliance, reaching a plateau in recovery.

Rationale

There are no quality trials of cognitive rehabilitation for the treatment of memory and executive problems attributed to COVID. Cognitive rehabilitation has negligible adverse effects, is moderate to high cost depending on numbers of treatments required, and is recommended for patients meeting indications.

Evidence

A comprehensive literature search was conducted using PubMed, Scopus, CINAHL, Cochrane Library, and Google Scholar from January 2019 to February 2021 using the following terms: rehabilitation; coronavirus infections, coronavirus, COVID-19, novel coronavirus, SARS-CoV-2, 2019-nCoV; controlled clinical trial, controlled trials, randomized controlled trial, randomized controlled trials, random allocation, random*, randomized, randomization, randomly; systematic, systematic review, retrospective, prospective studies; clinical study; observational study clinical trial; non-randomized controlled trials as topics. We found and reviewed 112 articles in PubMed, 4,699 in Scopus, 3 in CINAHL, 13 in Cochrane Library, 34,300 in Google Scholar, and 0 from other sources†. We considered for inclusion 14 from PubMed, 4 from Scopus, 0 from CINAHL, 0 from Cochrane Library, 0 from

Google Scholar, and 0 from other sources. Of the 18 articles considered for inclusion, 1 randomized trial and 8 systematic reviews met the inclusion criteria. There were no exclusion criteria.

† The results for databases are sorted by relevancy based on customized search term algorithms. Algorithms for each database determine relevancy. The first 100 articles are reviewed in each search, and if relevant literature appears in the first 100 articles, we review an additional 100 articles. If relevant articles appear in these additional 100 articles, we then review another 100. We continue this pattern of review until we review a batch of 100 articles that contains no relevant literature. When this happens then the remaining articles are not reviewed due to a lack of relevancy.

9.6. JOINT PAIN

Joint pain is common in subacute and chronic COVID (303,304,305), with 27.3% reporting joint pain at 2 months after COVID onset (303). Detailed guidance is available by body part in other [ACOEM guidelines](#) (see, e.g., [Ankle and Foot Disorders](#), [Elbow Disorders](#), [Chronic Pain](#), [Hand/Wrist/Forearm Disorders](#), [Hip and Groin Disorders](#), [Knee Disorders](#), [Low Back Disorders](#), [Neck Disorders](#), [Shoulder Disorders](#)).

9.7. MENTAL HEALTH

Treatments for mental health disorders that result from COVID-19 have not undergone rigorous trials for efficacy. Only low-quality trials have thus far been reported. One combined anxiety, depression, and stress, reporting that cognitive behavioral therapy (CBT) was effective (306). Another trial found progressive muscle relaxation helpful for anxiety and sleep quality (307). Another trial found an internet-based intervention on depression and anxiety to be effective (308).

In the absence of quality evidence specific to COVID-19, analogy to existing quality evidence and evidence-based guidance is recommended for screening, diagnosis, and treatments. These are addressed in detail in guidelines on [Anxiety Disorders](#), [Depressive Disorders](#), and [Posttraumatic Stress Disorder](#).

Utilizing evidence for generalized anxiety disorder, anxiety related to COVID-19 is recommended to be best initially treated with education (I), CBT (B,C), aerobic exercise (C), and strengthening exercise (I). Due to strong addictive potential, benzodiazepines are not recommended for routine use (C). Other potential early treatments include insight-oriented therapies (I), distractive methods (C), exposure therapy/prolonged exposure therapy (I), virtual reality exposure therapy (I) and mindfulness therapy (I). Other medications with evidence of efficacy include buspirone (C), quetiapine (B), beta-blockers (B), pregabalin (B), and hydroxyzine (C). Details are in the [Anxiety Disorders Guideline](#).

Utilizing evidence for major depressive disorder, depression related to COVID is best treated initially by reducing or eliminating sedating medication (I), education (I), antidepressant medication (SSRI, SNRI, TCA, MAOI) (B), cognitive behavioral therapy (B), aerobic exercise (C), and strengthening exercise (I). Benzodiazepine medication is not recommended. Other recommended medications include antipsychotics, olanzapine/fluoxetine, agomelatine, eszopiclone, nefazodone, zolpidem for sleep disorders (C). Weight loss may be selectively indicated in patients with obesity (B). Transcranial magnetic stimulation (C), repetitive transcranial magnetic stimulation (C), low-field magnetic stimulation (B), and light therapy (C) are also potential treatments. Severe cases may be treated with electroconvulsive therapy (B). See [Depressive Disorders Guideline](#).

Utilizing evidence for posttraumatic stress disorder, PTSD related to COVID is best treated initially with aerobic exercise (B), strengthening exercise (B), cognitive behavioral therapy (B), exposure

therapy (B), prolonged exposure therapy (B), virtual reality (B), imagery rehearsal training (B), and narrative exposure therapy (C). Medications with evidence of efficacy include sertraline (B), paroxetine (B), fluoxetine (I), escitalopram (I), citalopram (C), venlafaxine (B), mirtazapine (B), phenelzine (C), nefazodone (C), quetiapine (I), olanzapine (C), and prazosin (I). Other treatments potentially indicated include guided imagery (I), deep breathing exercises (I), meditation (I), and mindfulness (I). See the [Posttraumatic Stress Disorder Guideline](#).

10. CONTRIBUTORS

Editor-in-Chief

Kurt T. Hegmann, MD, MPH, FACOEM, FACP

Evidence-based Practice COVID-19 Panel Members

Clayton T. Cowl, MD, MS, FACOEM

Philip Harber, MD, MPH, FACOEM, FCCP, ATSF

Mark H. Hyman, MD, FACP, FIAIME

Karin A. Pacheco, MD, MSPH, FAAAAI

Thomas Winters, MD, FACOEM, FACPM, FACP

Eric M. Wood, MD, MPH, FACOEM

Evidence-based Practice COVID-19 Consultants

Steven Mandel, MD, FACOEM

William Niehaus, MD

Greg S. Vanichkachorn, MD, MPH, FACOEM

These panel members represent expertise in occupational medicine, internal medicine, pulmonary medicine, and infectious disease. As required for quality guidelines – Institute of Medicine’s (IOM’s) Standards for Developing Trustworthy Clinical Practice Guidelines and Appraisal of Guidelines for Research and Evaluation (AGREE) – a detailed application process captured conflicts of interest. The above Panel has none to declare relevant to this guideline.

Research Team

Kurt T. Hegmann, MD, MPH, FACOEM, FACP

Matthew S. Thiese, PhD, MSPH

Emilee Eden, MPH

Kristine Hegmann, MSPH, CIC

Jenna L. Praggastis, BS

Alison Mancuso, BA

Braydon R. Black

Madison N. Tallman
Madeleine Smith, BS
Elise D. Chan
Andrew S. Barbee
Michael R. Langston, BS
Christina P. Pick, BS
Chapman B. Cox, BS
Derrick K. Wong
Jenny L. Echeverria
Uchenna C. Ogbonnnaya MS, CSCS
Katherine C. Castro, MPH

Specialty Society and Society Representative Listing: Coronavirus (COVID-19) Guideline

ACOEM acknowledges the following organizations and their representatives who served as reviewers of the Coronavirus (COVID-19) Guideline. Their contributions are greatly appreciated. By listing the following individuals or organizations, it does not infer that these individuals or organizations support or endorse the Coronavirus (COVID-19) Guideline developed by ACOEM. Reviewers from three additional societies wished to remain anonymous.

American College of Chest Physicians

Holly Keyt, MD
Steven Q. Simpson, MD

Society for Healthcare Epidemiology of America

Meghan A. Baker, MD, ScD
David J. Weber, MD, MPH

Other Reviewers:

James W. Butler, MD, MPH, MRO, FAADEP, CIME, FACOEM
Victoria A. Cassano, MD, MPH, MPhil, FACOEM, FACPM
Glenn S. Pransky, MD, MOCCH, FACOEM
Kenji Saito, MD, JD, FACOEM
Tanisha K. Taylor, MD, MPH, FACP, CIME, FACOEM

10.2. CONFLICT OF INTEREST DISCLOSURES

Kurt T. Hegmann, MD, MPH, FACOEM, FACP (Editor-in-Chief and Methodology Committee Consultant)

Professor and Center Director, Rocky Mountain Center for Occupational and Environmental Health, University of Utah

National, Regional, Local Committee Affiliations—Board of Trustees (2003-12) and Chair (2010-12), American Board of Preventive Medicine; Chair, Federal Motor Carrier Safety Administration's Medical Review Board (2006-2010); Member, *Journal of Occupational and Environmental Medicine* Editorial Board

Other Guidelines Related Professional Activities—Chair, Evidence-based Practice Committee, ACOEM (2006-present)

Research Grants/Other Support—NIOSH (CDC) Training grants and research grants primarily on the epidemiology of musculoskeletal disorders (e.g., CTS, shoulder tendinosis, LBP) and truck driver safety.

Financial/Non-Financial Conflict of Interest—Consultant, Reed Group (ACOEM Guidelines Editor-in-Chief); Consulting with companies, unions and insurers

Clayton T. Cowl, MD, MS, FACOEM (Panel Member)

Associate Professor of Medicine, Mayo Clinic

National, Regional, Local Committee Affiliations—Past President, American College of Chest Physicians

Other Guidelines Related Professional Activities—None

Research Grants/Other Support—Co-Co-Principal Investigator, U.S. Department of Labor RETAIN Grant

Financial/Non-Financial Conflict of Interest—None

Philip Harber, MD, MPH, FACOEM, ATSF, FCCP (Panel Member)

Adjunct Professor of Public Health, Mel and Enid Zuckerman College of Public Health, University of Arizona; Professor Emeritus, UCLA; Dr Philip Harber PLLC

National, Regional, Local Committee Affiliations— Member, ACOEM Council on Scientific Affairs; ISO (respirators)

Other Guidelines Related Professional Activities—Member, ACOEM Evidence-based Practice Asthma Panel

Research Grants/Other Support— National Library of Medicine (NIH)

Financial/Non-Financial Conflict of Interest—None

Mark H. Hyman, MD, FACP, FAADEP

HymanHealth; Internal Medicine, Emeritus Associate Clinical Professor of Medicine at UCLA

National, Regional, Local Committee Affiliations— Veritas Medicus Board Officer (2017-2020); Director/Chairman, International Academy of Independent Medical Evaluators (2008-2015); Member, ACOEM Public Safety Medicine Section; American College of Physicians Governor's Advisory Council (2008-2013)

Other Guidelines Related Professional Activities—Evidence-based Practice Occupational Asthma Panel, ACOEM; Medical Advisory Board, *The Medical Disability Advisor*, 6th Edition; Advisory Committee, American Medical Association (AMA) *Guides to the Evaluation of Permanent Impairment*; Editorial Board, *AMA Guides Newsletter*

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest— Honoraria: Teaching honoraria from various courses, ACOEM and IAIME related

Karin A. Pacheco, MD, MSPH, FAAAAI (Panel Member)

Staff Physician, Division of Environmental and Occupational Health Sciences, National Jewish Medical and Research Center; Assistant Professor, Preventive Medicine and Allergy/ Immunology, University of Colorado Health Sciences Center

National, Regional, Local Committee Affiliations—Reviewer/contributor to AMA Guides, 5th Edition and 6th edition; Chair of EORD Interest Section, American Academy of Allergy, Asthma & Immunology; Member of Occupational Disease Committee, American Academy of Allergy, Asthma & Immunology; Reviewer for American Journal of Respiratory and Critical Care Medicine, Allergy and Clinical Immunology, Annals of Allergy, European Respiratory Journal, and Clinical Experimental Allergy, Chest, and Allergy

Other Guidelines Related Professional Activities—Occupational Asthma Committee (Contributing Author), American College of Chest Physicians: “An Evidence-based Guide for the Diagnosis and Management of Occupational Asthma;” “Contact Dermatitis Practice Parameters Update” for American Academy of Allergy, Asthma & Immunology/American College of Allergy, Asthma and Immunology/Joint Council of Allergy, Asthma and Immunology

Research Grants/Other Support—Colorado Bioscience: Development of diagnostic tests for metal allergy, Colorado Challenge: Mechanisms and diagnosis of metal sensitization

Financial/Non-Financial Conflict of Interest—None

Thomas Winters, MD, FACOEM, FACPM, FACP (Panel Member)

Chief Medical Officer & Principal, OEHN; Visiting Scientist, Occupational and Environmental Health, Harvard School of Public Health; Visiting Lecturer, Harvard Medical School

National, Regional, Local Committee Affiliations—Member, Residency Advisory Committee, Occupational/Environmental Health, Harvard School of Public Health

Other Guidelines Related Professional Activities—Member, ACOEM Evidence-based Practice Opioids Panel

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—Consultations: healthcare and business organizations

Eric M. Wood, MD, MPH, FACOEM (Panel Member)

Occupational Medicine Program Director and Residency Program Director in Occupational and Environmental Medicine, Rocky Mountain Center for Occupational and Environmental Health, University of Utah; Associate Professor (Clinical), School of Medicine, University of Utah; Department of Family and Preventive Medicine, University of Utah

National, Regional, Local Committee Affiliations—Vice-Chair, Occupational Medicine, American Board of Preventive Medicine; Member, ACOEM Maintenance of Certification Committee; Member, ACOEM Council on Education and Academic Affairs; Co-Chair, NIOSH, National Occupational Research Agenda, TWU Sector; Chair, American Board of Preventive Medicine Occupational Medicine Examination Committee; Member, Education Committee, Department of Family and Preventive Medicine, University of Utah; Residency Advisory Committee, Rocky Mountain Center for Occupational and Environmental Health, University of Utah

Guidelines Related Professional Activities— Hospital committee guideline for clinical practice; occupational medicine outpatient clinic guidelines

Research Grants/Other Support— NIOSH (CDC) Training grants and research grants primarily on the epidemiology of musculoskeletal disorders (e.g., CTS, shoulder tendinosis, LBP) and preventing work injuries and chronic illnesses in truck drivers

Financial/Non-Financial Conflict of Interest—Honoraria: Teaching honoraria from various courses, mostly ACOEM-related; Clinical: Primary, secondary and tertiary clinical management of occupational injuries and diseases

Steven Mandel, MD, FACOEM (Panel Consultant)

Clinical Professor of Neurology, Zucker School of Medicine at Hofstra/Northwell; Adjunct Professor of Medicine NY Medical College; Adjunct Professor of Podiatry at Temple University and NY College of Pediatric Medicine; Steven Mandel MD PC

National, Regional, Local Committee Affiliations—Medical Review Officer; Certified Medical Examiner; Reviewer, Journal of Voice; Expert Opinion Editor, Practical Neurology; Past President, NY Occupational and Environmental Medical Association; Diversity and Inclusion Committee, NY State and NY County Medical Society; Wellness Committee, NY State and NY County Medical Society; AMA Ambassador; NY State Liaison of the Senior Physician Section, AMA

Other Guidelines Related Professional Activities—Member, ACOEM Evidence-based Practice Traumatic Brain Injury Panel; Member, ACOEM Evidence-based Practice Ankle/Foot Panel; Member, ACOEM Evidence-based Practice Opioids Panel; Member, ACOEM Evidence-based Practice Disability Prevention and Treatment Panel; Member, Sports Concussion, American Academy of Neurology

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

William Niehaus, MD (Panel Consultant)

Assistant Professor, University of Colorado School of Medicine; Associate Program Director, CU PM&R Residency Program; President and Founder, Free Rehabilitation Services Volunteer Partnership Clinic; Co-Founder, Combined Neuro Therapy Clinic; Medical Director, Outpatient PM&R Services

National, Regional, Local Committee Affiliations—Social Media Editor, Physical Medicine & Rehabilitation Journal; Reviewer, PM&R Journal; Founder and Physician Lead for UCH Proactive Rehabilitative Interdisciplinary Management & Evaluation of Disposition Team QI Initiative; PM&R Physician Advisor, UCH MICU Early Mobility Project; Member, Surgical & Trauma Spinal Cord Injury QI Team; Member, Quality Leadership Team on the Acute Inpatient Rehabilitation Unit

Other Guidelines Related Professional Activities— Member, AAPMR Clinical Guidance on Assessment and Treatment of Fatigue in Postacute Sequelae of SARS-CoV-2 Infection

Research Grants/Other Support—None

Financial/Non-Financial Conflict of Interest—None

Greg S. Vanichkachorn, MD, MPH, FACOEM (Panel Consultant)

Senior Associate Consultant, Mayo Clinic – Rochester; Medical Director, COVID Activity Rehabilitation Program

National, Regional, Local Committee Affiliations— Co-lead, Mayo Clinic Multidisciplinary Approach Guiding Post COVID Investigation, Education, and Symptom Management (MAGPIES) team

Other Guidelines Related Professional Activities—None

Research Grants/Other Support—Principal Investigator, Mayo Clinic – Rochester: Treatment Outcomes from Early Intervention in Post COVID Syndrome, Post COVID Syndrome: Patient Population and Treatment Outcomes

Financial/Non-Financial Conflict of Interest—Consultant for Brain FX™

REFERENCES

1. van Tulder, M., Furlan, A., Bombardier, C., Bouter, L. Updated method guidelines for systematic reviews in the cochrane collaboration back review group. *Spine (Phila Pa 1976)*; Jun 15 2003.
2. Murti, M., Achonu, C., Smith, B., Brown, K., Kim, J., Johnson, J., Ravindran, S., Buchan, S. COVID-19 workplace outbreaks by industry sector and their associated household transmission, Ontario, Canada, January to June, 2020. *JOEM*; 2021.
3. Baptista, M., Burton, W., Pawlecki, B., Pransky, G. A physician's guide for workers' return to work during COVID-19 pandemic. *JOEM*; Mar 1 2021.
4. Ritchie, Hannah, Ortiz-Ospina, Esteban, Beltekian, Diana, Mathieu, Edouard, Hasell, Joe, Macdonald, Bobbie, Giattino, Charlie, Roser, Max. Coronavirus (COVID-19) Vaccinations. 2021.
5. Herby, J., Jonung, L., Hanke, S. A literature review and meta-analysis of the effects of lockdowns on COVID-19 mortality. *Appl. Econ.*; 2022.
6. Chinazzi, Matteo, Davis, Jessica T, Ajelli, Marco, Gioannini, Corrado, Litvinova, Maria, Merler, Stefano, y Piontti, Ana Pastore, Mu, Kunpeng, Rossi, Luca, Sun, Kaiyuan. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science*; 2020.
7. Johns Hopkins University. Coronavirus COVID-19 Global Cases by the Center for Systems Science and Engineering (CSSE) at Johns Hopkins University (JHU). 2020.
8. Braine, Theresa. Planes almost entirely safe from coronavirus transmission if everyone wears masks: study. 2020.
9. Khan, Rabeea F, Meyer, John D. How Does the Hierarchy of Controls Integrate With the Epidemiologic Triangle to Help Address and Understand Transmission of SARS-CoV-2?. *Journal of Occupational and Environmental Medicine*; 2020.
10. Chu, Derek K, Akl, Elie A, Duda, Stephanie, Solo, Karla, Yaacoub, Sally, Schünemann, Holger J, El-harakeh, Amena, Bognanni, Antonio, Lotfi, Tamara, Loeb, Mark. Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *The Lancet*; 2020.
11. WHO. COVID-19 table of PPE with description and related standard (simplified version). 2020.
12. WHO. COVID-19 Technical Specifications for personal protective equipment, list of standards and checklists. 2020.
13. WHO. Coronavirus disease (COVID-19): Masks. 2020.
14. WHO. Mask use in the context of COVID-19: interim guidance, 1 December 2020. 2020.
15. Perencevich, Eli N., Diekema, Daniel J., Edmond, Michael B. Moving Personal Protective Equipment Into the Community: Face Shields and Containment of COVID-19. *JAMA*; 2020.
16. ASHRAE. Core Recommendations for Reducing Airborne Infectious Aerosol Exposure. 2021.
17. ASHRAE. General Information. 2021.
18. CDC. Ventilation in Buildings. 2021.
19. ASHRAE. ASHRAE Position Document on Infectious Aerosols. 2020.
20. ACGIH. White Paper on Ventilation for Industrial Settings during the COVID-19 Pandemic. 2020.
21. CDC. Interim Guidance for Businesses and Employers to Plan and Respond to Coronavirus Disease 2019 (COVID-19). 2020.
22. ASHRAE. ASHRAE Epidemic Task Force. 2020.
23. CDC. Frequently Asked Questions. 2020.
24. Safelane Health. COVID Symptoms Tracker and Automated Decision Support Tool. 2020.
25. Johns Hopkins Medicine. Coronavirus (COVID-19) Self-Checker. 2020.
26. CDC. Symptoms of Coronavirus. 2020.
27. Massachusetts General Hospital. COVID-19 Symptom Tracker App. 2020.
28. WHO. Report of the WHO-China joint mission on coronavirus disease 2019 (COVID-19). 2020.
29. Zhu, Y, Bloxham, C, Hulme, K, Sinclair, J, Tong, ZWM, Steele, L, Noye, E, Lu, J, Chew, KY, Pickering, J, Gilks, C, Bowen, A, Short, K. Children are unlikely to have been the primary source of household SARS-CoV-2 infections. *medRxiv*; 2020.
30. Liu, Z., Xing, B., Xue Za, Z. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) in China. Feb 10 2020.

31. Cristiani, L., Mancino, E., Matera, L., Nenna, R., Pierangeli, A., Scagnolari, C., Midulla, F. Will children reveal their secret? The coronavirus dilemma. *Eur. Clin. Respir. J.*; 2020.
32. Rubin, R. School superintendents confront COVID-19—"There are no good options for next year". *JAMA*; 2020.
33. Davies, N., Klepac, P., Liu, Y., Prem, K., Jit, M., Pearson, C., Quilty, B., Kucharski, A., Gibbs, H., Clifford, S., Gimma, A., van Zandvoort, K., Munday, J., Diamond, C., Edmunds, W., Houben, R., Hellewell, J., Russell, T., Abbott, S., Funk, S., Bosse, N., Sun, Y., Flasche, S., Rosello, A., Jarvis, C., Eggo, R., group, Cmmid,Covid-,working. Age-dependent effects in the transmission and control of COVID-19 epidemics. *Nat. Med.*; 2020/06/16.
34. Dorn, E., Hancock, B., Sarakatsannis, J., Viruleg, E. COVID-19 and student learning in the United States: The hurt could last a lifetime. *McKinsey & Company*; 2020.
35. Horowitz, J. Lower-income parents most concerned about their children falling behind amid COVID-19 school closures. *Pew Research Center*; 2020.
36. Bacher-Hicks, A., Goodman, J., Mulhern, C. Inequality in household adaptation to schooling shocks: Covid-induced online learning engagement in real time. 2020.
37. Van Lancker, W., Parolin, Z. COVID-19, school closures, and child poverty: a social crisis in the making. *Lancet Public Health*; 2020.
38. Burgess, S., Sievertsen, H. Schools, skills, and learning: the impact of COVID-19 on education. 2020.
39. Azevedo, J. P., Hasan, A., Goldemberg, D., Iqbal, S. A., Geven, K. Simulating the potential impacts of covid-19 school closures on schooling and learning outcomes: a set of global estimates. 2020.
40. Fitzpatrick, BR, Berends, M, Ferrare, JJ, Waddington, RJ. Virtual illusion: Comparing student achievement and teacher and classroom characteristics in online and brick-and-mortar charter schools. *Educ. Res.*; 2020.
41. Durlak, J., Dymnicki, A., Taylor, R., Weissberg, R., Schellinger, K., Dubois, D., Lipsey, M., Greenberg, M., O'brien, M. Collaborative for academic, social, and emotional learning (CASEL). 2007.
42. Loades, M., Chatburn, E., Higson-Sweeney, N., Reynolds, S., Shafran, R., Brigden, A., Linney, C., McManus, M., Borwick, C., Crawley, E. Rapid systematic review: the impact of social isolation and loneliness on the mental health of children and adolescents in the context of COVID-19. *J Am Acad Child Adolesc Psychiatry*; 2020/06/03/.
43. Birnbaum, M. Reopened schools in Europe and Asia have largely avoided coronavirus outbreaks. They have lessons for the U.S. 2020.
44. Stein-Zamir, C., Abramson, N., Shoob, H., Libal, E., Bitan, M., Cardash, T., Cayam, R., Miskin, I. A large COVID-19 outbreak in a high school 10 days after schools' reopening, Israel, May 2020. *Euro Surveill.*; 2020.
45. Schwartz, F, Lieber, D. Israelis fear schools reopened too soon as Covid-19 cases climb. 2020.
46. Bank, The,World, UNICEF, , UNESCO. The state of the global education crisis: a path to recovery. December 3, 2021.
47. CDC. Late Sequelae of COVID-19. 2020.
48. Banerji, Aleena, Wickner, Paige G., Saff, Rebecca, Stone, Cosby A., Robinson, Lacey B., Long, Aidan A., Wolfson, Anna R., Williams, Paul, Khan, David A., Phillips, Elizabeth, Blumenthal, Kimberly G. mRNA Vaccines to Prevent COVID-19 Disease and Reported Allergic Reactions: Current Evidence and Suggested Approach. *The Journal of Allergy and Clinical Immunology: In Practice*; 2020/12/31/.
49. Zhou, Zhao-Hua, Stone, Cosby A., Jakubovic, Baruch, Phillips, Elizabeth J., Sussman, Gordon, Park, JuMe, Hoang, Uyen, Kirshner, Susan L., Levin, Robert, Kozlowski, Steven. Anti-PEG IgE in anaphylaxis associated with polyethylene glycol. *The Journal of Allergy and Clinical Immunology: In Practice*; 2020/11/17/.
50. Blumenthal, Kimberly G., Freeman, Esther E., Saff, Rebecca R., Robinson, Lacey B., Wolfson, Anna R., Foreman, Ruth K., Hashimoto, Dean, Banerji, Aleena, Li, Lily, Anvari, Sara, Shenoy, Erica S. Delayed Large Local Reactions to mRNA-1273 Vaccine against SARS-CoV-2. *New England Journal of Medicine*; 2021.
51. Cecinati, Valerio, Principi, Nicola, Brescia, Letizia, Giordano, Paola, Esposito, Susanna. Vaccine administration and the development of immune thrombocytopenic purpura in children. *Human Vaccines & Immunotherapeutics*; 2013/05/14.
52. Arnold, D. T., Milne, A., Samms, E., Staddon, L., Maskell, N.A., Hamilton, F.W. Symptoms after COVID-19 vaccination in patients with persistent symptoms after acute infection: a case series. *Ann. Intern. Med.*; 2021.
53. Zhang, Lizhou, Jackson, Cody B, Mou, Huihui, Ojha, Amrita, Peng, Haiyong, Quinlan, Brian D, Rangarajan, Erumbi S, Pan, Andi, Vanderheiden, Abigail, Suthar, Mehul S. SARS-CoV-2 spike-protein D614G mutation increases virion spike density and infectivity. *Nature communications*; 2020.

54. Hou, Yixuan J, Chiba, Shiho, Halfmann, Peter, Ehre, Camille, Kuroda, Makoto, Dinnon, Kenneth H, Leist, Sarah R, Schäfer, Alexandra, Nakajima, Noriko, Takahashi, Kenta. SARS-CoV-2 D614G variant exhibits efficient replication ex vivo and transmission in vivo. *Science*; 2020.
55. Kemp, SA, Collier, DA, Datir, R, Gayed, S, Jahun, A, Hosmillo, M, Ferreira, IATM, Rees-Spear, C, Mlcochova, P, Lumb, Ines Ushiro, Roberts, David, Chandra, Anita, Temperton, N, Sharrocks, K, Blane, E, Briggs, JAG, van Gils, MJ, Smith, KGC, Bradley, JR, Smith, C, Goldstein, RA, Goodfellow, IG, Smielewska, A, Skittrall, JP, Gouliouris, T, Gkrania-Klotsas, E, Illingworth, CJR, McCoy, LE, Gupta, RK. Neutralising antibodies drive Spike mediated SARS-CoV-2 evasion. *medRxiv*; 2020.
56. Galloway, Summer E, Paul, Prabasaj, MacCannell, Duncan R, Johansson, Michael A, Brooks, John T, MacNeil, Adam, Slayton, Rachel B, Tong, Suxiang, Silk, Benjamin J, Armstrong, Gregory L. Emergence of SARS-CoV-2 b. 1.1. 7 lineage—united states, december 29, 2020–january 12, 2021. *Morbidity and Mortality Weekly Report*; 2021.
57. Madhi, S., Baillie, V., Cutland, C., Voysey, M., Koen, A., Fairlie, L., Padayachee, S., Dheda, K., Barnabas, S., Bhorat, Q. Efficacy of the ChAdOx1 nCoV-19 Covid-19 vaccine against the B. 1.351 variant. *NEJM*; 2021.
58. Liu, Yang, Liu, Jianying, Xia, Hongjie, Zhang, Xianwen, Fontes-Garfias, Camila R., Swanson, Kena A., Cai, Hui, Sarkar, Ritu, Chen, Wei, Cutler, Mark, Cooper, David, Weaver, Scott C., Muik, Alexander, Sahin, Ugur, Jansen, Kathrin U., Xie, Xuping, Dormitzer, Philip R., Shi, Pei-Yong. Neutralizing Activity of BNT162b2-Elicited Serum. *New England Journal of Medicine*; 2021.
59. CDC. SARS-CoV-2 variant classifications and definitions. December 1, 2021.
60. CDC. What you need to know about variants. February 25, 2022.
61. CDC. Omicron variant: what you need to know. February 2, 2022.
62. Wang, Zijun, Schmidt, Fabian, Weisblum, Yiska, Muecksch, Frauke, Barnes, Christopher O., Finkin, Shlomo, Schaefer-Babajew, Dennis, Cipolla, Melissa, Gaebler, Christian, Lieberman, Jenna A., Oliveira, Thiago Y., Yang, Zhi, Abernathy, Morgan E., Huey-Tubman, Kathryn E., Hurley, Arlene, Turroja, Martina, West, Kamille A., Gordon, Kristie, Millard, Katrina G., Ramos, Victor, Da Silva, Justin, Xu, Jianliang, Colbert, Robert A., Patel, Roshni, Dizon, Juan, Unson-O'Brien, Cecille, Shimeliovich, Irina, Gazumyan, Anna, Caskey, Marina, Bjorkman, Pamela J., Casellas, Rafael, Hatzioannou, Theodora, Bieniasz, Paul D., Nussenzweig, Michel C. mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants. *bioRxiv*; 2021.
63. Tarke, Alison, Sidney, John, Methot, Nils, Zhang, Yun, Dan, Jennifer M., Goodwin, Benjamin, Rubiro, Paul, Sutherland, Aaron, da Silva Antunes, Ricardo, Frazier, April, Rawlings, Stephen A., Smith, Davey M., Peters, Bjoern, Scheuermann, Richard H., Weiskopf, Daniela, Crotty, Shane, Grifoni, Alba, Sette, Alessandro. Negligible impact of SARS-CoV-2 variants on CD4⁺ and CD8⁺ T cell reactivity in COVID-19 exposed donors and vaccinees. *bioRxiv*; 2021.
64. Willyard, C. What the omicron wave is revealing about human immunity. *Nature*; February 2, 2022.
65. Chou, R., Dana, T., Jungbauer, R. Update alert 6: masks for prevention of respiratory virus infections, including SARS-CoV-2, in health care and community settings. *Ann. Intern. Med.*; 2021.
66. Honein, Margaret A, Christie, Athalia, Rose, Dale A, Brooks, John T, Meaney-Delman, Dana, Cohn, Amanda, Sauber-Schatz, Erin K, Walker, Allison, McDonald, L Clifford, Liburd, Leandris C. Summary of Guidance for Public Health Strategies to Address High Levels of Community Transmission of SARS-CoV-2 and Related Deaths, December 2020. *Morbidity and Mortality Weekly Report*; 2020.
67. Alfano, Vincenzo, Ercolano, Salvatore. The efficacy of lockdown against COVID-19: a cross-country panel analysis. *Applied health economics and health policy*; 2020.
68. Makinde, Olusola S, Adeola, Abiodun M, Abiodun, Gbenga J, Olusola-Makinde, Olubukola O, Alejandro, Aceves. Comparison of Predictive Models and Impact Assessment of Lockdown for COVID-19 over the United States. *Journal of Epidemiology and Global Health*; 2021.
69. Guy, Gery P. Association of State-Issued Mask Mandates and Allowing On-Premises Restaurant Dining with County-Level COVID-19 Case and Death Growth Rates—United States, March 1–December 31, 2020. *MMWR. Morbidity and Mortality Weekly Report*; 2021.
70. Shlomai, A. National lockdown not cost effective in Israel during COVID-2 pandemic. *PharmacoEcon Outcomes News*; 2021.
71. Miles, D. K., Stedman, M., Heald, A. H. "Stay at Home, Protect the National Health Service, Save Lives": A cost benefit analysis of the lockdown in the United Kingdom. *Int J Clin Pract*; Aug 13 2020.
72. Holland, Kristin M., Jones, Christopher, Vivolo-Kantor, Alana M., Idaikkadar, Nimi, Zwald, Marissa, Hoots, Brooke, Yard, Ellen, D'Inverno, Ashley, Swedo, Elizabeth, Chen, May S., Petrosky, Emiko, Board, Amy, Martinez, Pedro, Stone, Deborah M., Law, Royal, Coletta, Michael A., Adjemian, Jennifer, Thomas, Craig, Puddy, Richard W., Peacock, Georgina, Dowling,

- Nicole F., Houry, Debra. Trends in US Emergency Department Visits for Mental Health, Overdose, and Violence Outcomes Before and During the COVID-19 Pandemic. *JAMA Psychiatry*; 2021.
73. Henderson, Emily. FAIR Health study highlights the impact of COVID-19 on pediatric mental health. 2021.
74. Amsalem, Doron, Dixon, Lisa B., Neria, Yuval. The Coronavirus Disease 2019 (COVID-19) Outbreak and Mental Health: Current Risks and Recommended Actions. *JAMA Psychiatry*; 2021.
75. Magson, Natasha R., Freeman, Justin Y. A., Rapee, Ronald M., Richardson, Cele E., Oar, Ella L., Fardouly, Jasmine. Risk and Protective Factors for Prospective Changes in Adolescent Mental Health during the COVID-19 Pandemic. *Journal of Youth and Adolescence*; 2021/01/01.
76. Ravens-Sieberer, Ulrike, Kaman, Anne, Erhart, Michael, Devine, Janine, Schlack, Robert, Otto, Christiane. Impact of the COVID-19 pandemic on quality of life and mental health in children and adolescents in Germany. *European Child & Adolescent Psychiatry*; 2021/01/25.
77. Rodriguez-Morales, Alfonso J, Cardona-Ospina, Jaime A, Gutiérrez-Ocampo, Estefanía, Villamizar-Peña, Rhuvi, Holguin-Rivera, Yeimer, Escalera-Antezana, Juan Pablo, Alvarado-Arnez, Lucia Elena, Bonilla-Aldana, D Katterine, Franco-Paredes, Carlos, Henao-Martinez, Andrés F. Clinical, laboratory and imaging features of COVID-19: A systematic review and meta-analysis. *Travel medicine and infectious disease*; 2020.
78. Xu, Xi, Yu, Chengcheng, Qu, Jing, Zhang, Lieguang, Jiang, Songfeng, Huang, Deyang, Chen, Bihua, Zhang, Zhiping, Guan, Wanhua, Ling, Zhoukun. Imaging and clinical features of patients with 2019 novel coronavirus SARS-CoV-2. *European Journal of Nuclear Medicine and Molecular Imaging*; 2020.
79. Liang, Wenhua, Liang, Hengrui, Ou, Limin, Chen, Binfeng, Chen, Ailan, Li, Caichen, Li, Yimin, Guan, Weijie, Sang, Ling, Lu, Jiatao. Development and validation of a clinical risk score to predict the occurrence of critical illness in hospitalized patients with COVID-19. *JAMA Internal Medicine*; 2020.
80. Health, Guangzhou, Institute, of, Respiratory. Calculation Tool For Predicting Critical-ill COVID-19 At Admission. 2020.
81. Yuan, Jing, Zou, Rougrong, Zeng, Lijiao, Kou, Shanglong, Lan, Jianfeng, Li, Xiaohe, Liang, Yanhua, Ding, Xiaoyan, Tan, Guoyu, Tang, Shenghong. The correlation between viral clearance and biochemical outcomes of 94 COVID-19 infected discharged patients. *Inflammation Research*; 2020.
82. Retallack, H., Di Lullo, E., Arias, C., Knopp, K. A., Laurie, M. T., Sandoval-Espinosa, C., Mancía Leon, W. R., Krencik, R., Ullian, E. M., Spatazza, J., Pollen, A. A., Mandel-Brehm, C., Nowakowski, T. J., Kriegstein, A. R., DeRisi, J. L. Zika virus cell tropism in the developing human brain and inhibition by azithromycin. *Proc Natl Acad Sci U S A*; Dec 13 2016.
83. Madrid, P. B., Panchal, R. G., Warren, T. K., Shurtleff, A. C., Endsley, A. N., Green, C. E., Kolokoltsov, A., Davey, R., Manger, I. D., Gilfillan, L., Bavari, S., Tanga, M. J. Evaluation of Ebola Virus Inhibitors for Drug Repurposing. *ACS Infect Dis*; Jul 10 2015.
84. Bosseboeuf, E, Aubry, M, Nhan, T, Pina, JJ, Rolain, JM, Raoult, D, Musso, D. Azithromycin Inhibits the Replication of Zika Virus. *J. Antivir. Antiretrovir*; 2018.
85. Bacharier, Leonard B, Guilbert, Theresa W, Mauger, David T, Boehmer, Susan, Beigelman, Avraham, Fitzpatrick, Anne M, Jackson, Daniel J, Baxi, Sachin N, Benson, Mindy, Burnham, Carey-Ann D. Early administration of azithromycin and prevention of severe lower respiratory tract illnesses in preschool children with a history of such illnesses: a randomized clinical trial. *Jama*; 2015.
86. Albani, Filippo, Fusina, Federica, Giovannini, Alessia, Ferretti, Pierluigi, Granato, Anna, Prezioso, Chiara, Divizia, Danilo, Sabaini, Alessandra, Marri, Marco, Malpetti, Elena. Impact of Azithromycin and/or Hydroxychloroquine on Hospital Mortality in COVID-19. *Journal of clinical medicine*; 2020.
87. Hsia, Brian C, Greige, Nicolas, Quiroz, Jose A, Khokhar, Ahmed S, Daily, Johanna, Di Biase, Luigi, Ferrick, Kevin J, Fisher, John D, Krumer, Andrew. QT prolongation in a diverse, urban population of COVID-19 patients treated with hydroxychloroquine, chloroquine, or azithromycin. *Journal of Interventional Cardiac Electrophysiology*; 2020.
88. Rodríguez-Molinero, Alejandro, Pérez-López, Carlos, Gálvez-Barrón, César, Miñarro, Antonio, Macho, Oscar, López, Gabriela F, Robles, Maria Teresa, Dapena, María Dolores, Martínez, Sergi, Rodríguez, Ezequiel. Observational study of azithromycin in hospitalized patients with COVID-19. *PLoS one*; 2020.
89. Irie, Kei, Nakagawa, Atsushi, Fujita, Hiroto, Tamura, Ryo, Eto, Masaaki, Ikeshue, Hiroaki, Muroi, Nobuyuki, Tomii, Keisuke, Hashida, Tohru. Pharmacokinetics of Favipiravir in Critically Ill Patients with COVID-19. *Clinical and Translational Science*; 2020.
90. Pongpirul, Wannarat A, Wiboonthukul, Surasak, Charoenpong, Lantharita, Panitatum, Nayot, Vachiraphan, Apichart, Uttayamakul, Sumonmal, Pongpirul, Krit, Manosuthi, Weerawat, Prasithsirikul, Wisit. Clinical course and potential predictive factors for pneumonia of adult patients with Coronavirus Disease 2019 (COVID-19): A retrospective observational analysis of 193 confirmed cases in Thailand. *PLoS neglected tropical diseases*; 2020.

91. Sano, Tomoya, Kimizuka, Yoshifumi, Fujikura, Yuji, Hisada, Tetsuya, Watanabe, Chie, Suematsu, Ryohei, Izumi, Kaori, Sugiura, Hiroaki, Miyata, Jun, Shinmoto, Hiroshi. COVID-19 in older adults: Retrospective cohort study in a tertiary hospital in Japan. *Geriatrics & gerontology international*; 2020.
92. Yaylaci, Selçuk, Dheir, Hamad, Şenocak, Didar, Genc, Ahmed Bilal, Kocayigit, Havva, Çekiç, Deniz, Varım, Ceyhun, Aydın, Abdülkadir, Koroglu, Mehmet, Karabay, Oğuz. The effects of favipiravir on hematological parameters of covid-19 patients. *Revista da Associação Médica Brasileira*; 2020.
93. Yamamura, Hitoshi, Matsuura, Hiroshi, Nakagawa, Junichiro, Fukuoka, Hiroshi, Domi, Hisaya, Chujoh, Satoru. Effect of favipiravir and an anti-inflammatory strategy for COVID-19. *Critical Care*; 2020.
94. Ivashchenko, Andrey A, Dmitriev, Kirill A, Vostokova, Natalia V, Azarova, Valeria N, Blinow, Andrew A, Egorova, Alina N, Gordeev, Ivan G, Ilin, Alexey P, Karapetian, Ruben N, Kravchenko, Dmitry V. AVIFAVIR for treatment of patients with moderate COVID-19: interim results of a phase II/III multicenter randomized clinical trial. *medRxiv*; 2020.
95. Khamis, Faryal, Al Naabi, Hanan, Al Lawati, Adil, Ambusaidi, Zaiyana, Al Sharji, Mariam, Al Barwani, Umkulthum, Pandak, Nenad, Al Balushi, Zakariya, Al Bahrani, Maher, Al-Zakwani, Ibrahim. Randomized Controlled Open Label Trial on the Use of Favipiravir Combined with Inhaled Interferon beta-1b in Hospitalized Patients with Moderate to Severe COVID-19 Pneumonia. *International Journal of Infectious Diseases*; 2020.
96. Dabbous, Hany M, El-Sayed, Manal H, El Assal, Gihan, Elghazaly, Hesham, Ebeid, Fatma FS, Sherief, Ahmed F, Elgaafary, Maha, Fawzy, Ehab, Hassany, Sahar M, Riad, Ahmed R. A Randomized Controlled Study Of Favipiravir Vs Hydroxychloroquine In COVID-19 Management: What Have We Learned So Far?. 2020.
97. WHO. “Solidarity” clinical trial for COVID-19 treatments. 2020.
98. Yao, Tian-Tian, Qian, Jian-Dan, Zhu, Wen-Yan, Wang, Yan, Wang, Gui-Qiang. A systematic review of lopinavir therapy for SARS coronavirus and MERS coronavirus—A possible reference for coronavirus disease-19 treatment option. *Journal of medical virology*; 2020.
99. Verdugo-Paiva, Francisca, Izcovich, Ariel, Ragusa, Martín, Rada, Gabriel, Group, COVID-19, L·OVE, Working. Lopinavir/ritonavir for COVID-19: a living systematic review. *Medwave*; 2020.
100. Lê, Minh Patrick, Jaquet, Pierre, Patrier, Juliette, Wicky, Paul-Henri, Le Hingrat, Quentin, Veyrier, Marc, Kauv, Juliette, Sonnevile, Romain, Visseaux, Benoit, Laouénan, Cédric. Pharmacokinetics of lopinavir/ritonavir oral solution to treat COVID-19 in mechanically ventilated ICU patients. *Journal of Antimicrobial Chemotherapy*; 2020.
101. Gao, Guiju, Wang, Aibin, Wang, Sa, Qian, Fang, Chen, Meiling, Yu, Fengting, Zhang, Ju, Wang, Xudong, Ma, Xiaoyang, Zhao, Tianwei. Brief Report: Retrospective Evaluation on the Efficacy of Lopinavir/Ritonavir and Chloroquine to Treat Nonsevere COVID-19 Patients. *Journal of acquired immune deficiency syndromes (1999)*; 2020.
102. Karolyi, Mario, Pawelka, Erich, Mader, Theresa, Omid, Sara, Kelani, Hasan, Ely, Sarah, Jilma, Bernd, Baumgartner, Sebastian, Laferl, Hermann, Ott, Clemens. Hydroxychloroquine versus lopinavir/ritonavir in severe COVID-19 patients. *Wiener Klinische Wochenschrift*; 2020.
103. Kim, Ji-Won, Kim, Eun Jin, Kwon, Hyun Hee, Jung, Chi Young, Kim, Kyung Chan, Choe, Jung-Yoon, Hong, Hyo-Lim. Lopinavir-ritonavir versus hydroxychloroquine for viral clearance and clinical improvement in patients with mild to moderate coronavirus disease 2019. *The Korean journal of internal medicine*; 2020.
104. Lecronier, Marie, Beurton, Alexandra, Burrel, Sonia, Haudebourg, Luc, Deleris, Robin, Le Marec, Julien, Virolle, Sara, Nemlaghi, Safaa, Bureau, Côme, Mora, Pierre. Comparison of hydroxychloroquine, lopinavir/ritonavir, and standard of care in critically ill patients with SARS-CoV-2 pneumonia: an opportunistic retrospective analysis. *Critical Care*; 2020.
105. Zhu, Zhen, Lu, Zhaohui, Xu, Tianmin, Chen, Cong, Yang, Gang, Zha, Tao, Lu, Jianchun, Xue, Yuan. Arbidol monotherapy is superior to lopinavir/ritonavir in treating COVID-19. *Journal of Infection*; 2020.
106. Cao, Bin, Wang, Yeming, Wen, Danning, Liu, Wen, Wang, Jingli, Fan, Guohui, Ruan, Lianguo, Song, Bin, Cai, Yanping, Wei, Ming. A trial of lopinavir–ritonavir in adults hospitalized with severe Covid-19. *New England Journal of Medicine*; 2020.
107. Shih, Weichung J., Shen, Xin, Zhang, Peng, Xie, Tai. Remdesivir is Effective for Moderately Severe Patients: A Re-Analysis of the First Double-Blind, Placebo-Controlled, Randomized Trial on Remdesivir for Treatment of Severe COVID-19 Patients Conducted in Wuhan City. *Open Access Journal of Clinical Trials*; 2020.
108. Susan A. Olender, Katherine K. Perez, Alan S. Go, Bindu Balani, Eboni G. Price-Haywood, Nirav S. Shah, Su Wang, Theresa L. Walunas, Shobha Swaminathan, Jihad Slim, BumSik Chin, Stéphane De Wit. Remdesivir for Severe COVID-19 versus a Cohort Receiving Standard of Care. *Oxford University Press for the Infectious Diseases Society of America*; 2020.
109. Pasquini, Z., Montalti, R., Temperoni, C., Canovari, B., Mancini, M., Tempesta, M., Pimpini, D., Zallocco, N., Barchiesi, F. Effectiveness of remdesivir in patients with COVID-19 under mechanical ventilation in an Italian ICU. *J Antimicrob Chemother*; Nov 1 2020.

110. Kalligeros, M., Tashima, K. T., Mylonas, E. K., Rybak, N., Flanagan, T. P., Farmakiotis, D., Beckwith, C. G., Sanchez, M., Neill, M., Johnson, J. E., Garland, J. M., Aung, S., Byrd, K. M., O'Brien, T., Pandita, A., Aridi, J., Macias Gil, R., Larkin, J., Shehadeh, F., Mylonakis, E. Remdesivir Use Compared With Supportive Care in Hospitalized Patients With Severe COVID-19: A Single-Center Experience. *Open Forum Infect Dis*; Oct 2020.
111. Lee, C., Ahn, M. Y., Byeon, K., Choi, J. P., Hahm, C., Kim, H., Kim, S., Kim, T. H., Oh, J., Oh, D. H. Clinical Experience with Use of Remdesivir in the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2: a Case Series. *Infect Chemother*; Sep 2020.
112. Hsu, Chen-Yang, Lai, Chao-Chih, Yen, Amy Ming-Fang, Chen, Sam Li-She, Chen, Hsiu-Hsi. Efficacy of remdesivir in COVID-19 patients with a simulated two-arm controlled study. *medRxiv*; 2020.
113. Kaka, Anjum S, MacDonald, Roderick, Greer, Nancy, Vela, Kathryn, Duan-Porter, Wei, Obley, Adam, Wilt, Timothy J. Major Update: Remdesivir for Adults With COVID-19: A Living Systematic Review and Meta-analysis for the American College of Physicians Practice Points. *Annals of Internal Medicine*; 2021.
114. Qaseem, Amir, Yost, Jennifer, Etcheandia-Ikobaltzeta, Itziar, Abraham, George M, Jokela, Janet A, Forciea, Mary Ann, Miller, Matthew C, Humphrey, Linda L. Should Remdesivir Be Used for the Treatment of Patients With COVID-19? Rapid, Living Practice Points From the American College of Physicians (Version 1). *Annals of internal medicine*; 2020.
115. Qaseem, A., Yost, J., Etcheandia-Ikobaltzeta, I., Abraham, G., Jokela, J., Forciea, M., Miller, M., Humphrey, L. Should remdesivir be used for the treatment of patients with COVID-19? Rapid, living practice points from the American College of Physicians (Version 1). *Ann Intern Med*; 2020.
116. Hasan, SS, Radford, S, Kow, CS, Zaidi, STR. Venous thromboembolism in critically ill COVID-19 patients receiving prophylactic or therapeutic anticoagulation: a systematic review and meta-analysis. *J. Thromb. Thrombolysis*; 2020.
117. Falcone, M., Tiseo, G., Barbieri, G., Galfo, V., Russo, A., Viridis, A., Forfori, F., Corradi, F., Guarracino, F., Carrozzi, L., Celi, A., Santini, M., Monzani, F., De Marco, S., Pistello, M., Danesi, R., Ghiadoni, L., Alessio, F., Menichetti, F., Group, on behalf of the Pisa Covid Study. Role of low-molecular weight heparin in hospitalized patients with SARS-CoV-2 pneumonia: a prospective observational study. *Open Forum Infect. Dis.*; 2020.
118. Viecca, M, Radovanovic, D, Forleo, G, Santus, P. Enhanced platelet inhibition treatment improves hypoxemia in patients with severe Covid-19 and hypercoagulability. A case control, proof of concept study. *Pharmacol. Res. Commun.*; 2020/08/01/.
119. Di Perri, G. The rationale for low-molecular weight heparin (LMWH) use in SARS-CoV-2 infection. *Infez Med*; Jun 1 2020.
120. Fontana, P., Casini, A., Robert-Ebadi, H., Glauser, F., Righini, M., Blondon, M. Venous thromboembolism in COVID-19: systematic review of reported risks and current guidelines. *Swiss Med Wkly*; Jun 15 2020.
121. Nittari, G., Pallotta, G., Amenta, F., Tayebati, S. Current pharmacological treatments for SARS-COV-2: a narrative review. *Eur J Pharmacol*; 2020.
122. van Haren, F., Page, C., Laffey, J., Artigas, A., Camprubi-Rimblas, M., Nunes, Q., Smith, R., Shute, J., Carroll, M., Tree, J., Carroll, M., Singh, D., Wilkinson, T., Dixon, B. Nebulised heparin as a treatment for COVID-19: scientific rationale and a call for randomised evidence. *Critical Care*; 2020/07/22.
123. Ayerbe, L., Risco, C., Ayis, S. The association between treatment with heparin and survival in patients with COVID-19. *J. Thromb. Thrombolysis*; 2020.
124. Hippensteel, J., LaRiviere, W., Colbert, J., Langouët-Astrié, C., Schmidt, E. Heparin as a therapy for COVID-19: current evidence and future possibilities. *Am. J. Physiol. - Lung Cell. Mol. Physiol.*; 2020.
125. Birocchi, S., Manzoni, M., Podda, G. M., Casazza, G., Cattaneo, M. High rates of pulmonary artery occlusions in COVID-19. A meta-analysis. *Eur J Clin Invest*; Jan 2021.
126. Shah, A, Donovan, K, McHugh, A, Pandey, M, Aaron, L, Bradbury, CA., Stanworth, SJ., Alikhan, R, Von Kier, S, Maher, K, Curry, N, Shapiro, S, Rowland, MJ., Thomas, M, Mason, R, Holland, M, Holmes, T, Ware, M, Gurney, S, McKechnie, SR. Thrombotic and haemorrhagic complications in critically ill patients with COVID-19: a multicentre observational study. *J. Crit. Care*; 2020/09/18.
127. Rosovsky, R. P., Sanfilippo, K. M., Wang, T. F., Rajan, S. K., Shah, S., Martin, K. A., F, Ní,Áinle, Huisman, M., Hunt, B. J., Kahn, S. R., Kevane, B., Lee, A. Y., McLintock, C., Baumann Kreuziger, L. Anticoagulation practice patterns in COVID-19: A global survey. *Res Pract Thromb Haemost*; Jul 9 2020.
128. Lu, Y., Pan, L., Zhang, W., Cheng, F., Hu, S., Zhang, X., Jiang, H. A meta-analysis of the incidence of venous thromboembolic events and impact of anticoagulation on mortality in patients with COVID-19. *J. Glob. Infect. Dis.*; 2020/11/01/.

129. Moores, L., Tritschler, T., Brosnahan, S., Carrier, M., Collen, J., Doerschug, K., Holley, A., Jimenez, D., Le Gal, G., Rali, P. Prevention, diagnosis, and treatment of VTE in patients with coronavirus disease 2019: CHEST guideline and expert panel report. *Chest*; 2020.
130. De Havenon, A., Yaghi, S., Mistry, E., Delic, A., Hohmann, S., Shippey, E., Stulberg, E., Tirschwell, D., Frontera, J., Petersen, N. Endovascular thrombectomy in acute ischemic stroke patients with COVID-19: prevalence, demographics, and outcomes. *J. Neurointerv. Surg.*; 2020.
131. Reis, Gilmar, dos Santos Moreira-Silva, Eduardo Augusto, Silva, Daniela Carla Medeiros, Thabane, Lehana, Milagres, Aline Cruz, Ferreira, Thiago Santiago, Dos Santos, Castilho Vitor Quirino, de Souza Campos, Vitoria Helena, Nogueira, Ana Maria Ribeiro, de Almeida, Ana Paula Figueiredo Guimaraes. Effect of early treatment with flvoxamine on risk of emergency care and hospitalisation among patients with COVID-19: the TOGETHER randomised, platform clinical trial. *The Lancet Global Health*; 2022.
132. Lenze, Eric J, Mattar, Caline, Zorumski, Charles F, Stevens, Angela, Schweiger, Julie, Nicol, Ginger E, Miller, J Philip, Yang, Lei, Yingling, Michael, Avidan, Michael S. Fluvoxamine vs placebo and clinical deterioration in outpatients with symptomatic COVID-19: a randomized clinical trial. *Jama*; 2020.
133. Seftel, David, Boulware, David R. Prospective cohort of fluvoxamine for early treatment of coronavirus disease 19. *Open Forum Infectious Diseases*; 2021.
134. Oskotsky, T., Maric, I., Tang, A., Oskotsky, B., Wong, R., Aghaeepour, N., Sirota, M., Stevenson, D. Mortality risk among patients with COVID-19 prescribed selective serotonin reuptake inhibitor antidepressants. *JAMA Netw Open*; Nov 1 2021.
135. Mahase, E. Covid-19: Pfizer's paxlovid is 89% effective in patients at risk of serious illness, company reports. *BMJ*; 2021.
136. Fischer, William A, Eron, Joseph J, Holman, Wayne, Cohen, Myron S, Fang, Lei, Szewczyk, Laura J, Sheahan, Timothy P, Baric, Ralph S, Mollan, Katie R, Wolfe, Cameron R. Molnupiravir, an Oral Antiviral Treatment for COVID-19. *medRxiv*; 2021.
137. Painter, Wendy P, Holman, Wayne, Bush, Jim A, Almazedi, Firas, Malik, Hamzah, Eraut, Nicola CJE, Morin, Merribeth J, Szewczyk, Laura J, Painter, George R. Human safety, tolerability, and pharmacokinetics of molnupiravir, a novel broad-spectrum oral antiviral agent with activity against SARS-CoV-2. *Antimicrobial agents and chemotherapy*; 2021.
138. Khoo, Saye H, Fitzgerald, Richard, Fletcher, Thomas, Ewings, Sean, Jaki, Thomas, Lyon, Rebecca, Downs, Nichola, Walker, Lauren, Tansley-Hancock, Olana, Greenhalf, William. Optimal dose and safety of molnupiravir in patients with early SARS-CoV-2: a Phase I, open-label, dose-escalating, randomized controlled study. *Journal of Antimicrobial Chemotherapy*; 2021.
139. Jayk Bernal, A., Gomes da Silva, M., Musungaie, D., Kovalchuk, E., Gonzalez, A., Delos Reyes, V., Martín-Quirós, A., Caraco, Y., Williams-Diaz, A., Brown, M. Molnupiravir for oral treatment of Covid-19 in nonhospitalized patients. *NEJM*; 2022.
140. Administration, U.S., Food, and Drug, O'Shaughnessy, J. Molnupiravir Letter of Authorization. March 23, 2022.
141. Capra, R., De Rossi, N., Mattioli, F., Romanelli, G., Scarpazza, C., Sormani, M. P., Cossi, S. Impact of low dose tocilizumab on mortality rate in patients with COVID-19 related pneumonia. *Eur J Intern Med*; Jun 2020.
142. Xu, X, Han, M, Li, T, Sun, W, Wang, D, Fu, B, Zhou, Y, Zheng, X, Yang, Y, Li, X. Effective treatment of severe COVID-19 patients with tocilizumab. *ChinaXiv*; 2020.
143. Gritti, G, Raimondi, F, Ripamonti, D, Riva, I, Landi, F, Alborghetti, L, Frigeni, M, Damiani, M, Micò, C, Fagioli, S. Use of siltuximab in patients with COVID-19 pneumonia requiring ventilatory support. *medRxiv*; 2020.
144. Gritti, G, Alborghetti, L, Damiani, M, Jonathan, PMMD, Benjamin, MJ, Kanhai, KJK, Reljanovic, GT, Rizzi, M, Rambaldi, A, Alborghetti, L, Frigeni, M. IL-6 signalling pathway inactivation with siltuximab in patients with COVID-19 respiratory failure: an observational cohort study. 2020.
145. Campochiaro, C., Della-Torre, E., Cavalli, G., De Luca, G., Ripa, M., Boffini, N., Tomelleri, A., Baldissera, E., Rovere-Querini, P., Ruggeri, A., Monti, G., De Cobelli, F., Zangrillo, A., Tresoldi, M., Castagna, A., Dagna, L. Efficacy and safety of tocilizumab in severe COVID-19 patients: a single-centre retrospective cohort study. *Eur J Intern Med*; Jun 2020.
146. Somers, E., Eschenauer, G., Troost, J., Golob, J., Gandhi, T., Wang, L., Zhou, N., Petty, L., Baang, J., Dillman, N. Tocilizumab for treatment of mechanically ventilated patients with COVID-19. *medRxiv*; 2020.
147. Alattar, R., Ibrahim, T., Shaar, S. H., Abdalla, S., Shukri, K., Dagfal, J. N., Khatib, M. Y., Aboukamar, M., Abukhattab, M., Alsoub, H. A. Tocilizumab for the treatment of severe COVID-19. *J. Med. Virol.*; 2020.
148. Biran, N., Ip, A., Ahn, J., Go, R., Wang, S., Mathura, S., Sinclair, B., Bednars, U., Marafelias, M., Hansen, E. Tocilizumab among patients with COVID-19 in the intensive care unit: a multicentre observational study. *The Lancet Rheumatology*; 2020.

149. Canziani, L., Trovati, S., Brunetta, E., Testa, A., De Santis, M., Bombardieri, E., Guidelli, G., Albano, G., Folci, M., Squadroni, M. Interleukin-6 receptor blocking with intravenous tocilizumab in COVID-19 severe acute respiratory distress syndrome: a retrospective case-control survival analysis of 128 patients. *J. Autoimmun.*; 2020.
150. Chilimuri, S., Sun, H., Alemam, A., Kang, K., Lao, P., Mantri, N., Schiller, L., Sharabun, M., Shehi, E., Tejada, J., Yugay, A., Nayudu, S. Tocilizumab use in patients with moderate to severe COVID-19: a retrospective cohort study. *J Clin Pharm Ther*; Oct 24 2020.
151. Dastan, F., Saffaei, A., Haseli, S., Marjani, M., Moniri, A., Abtahian, Z., Abedini, A., Kiani, A., Seifi, S., Jammati, H., Hashemian, S., Pourabdollah Toutkaboni, M., Eslaminejad, A., Heshmatnia, J., Sadeghi, M., Nadji, S., Dastan, A., Baghaei, P., Varahram, M., Yousefian, S., Salamzadeh, J., Tabarsi, P. Promising effects of tocilizumab in COVID-19: A non-controlled, prospective clinical trial. *Int. Immunopharmacol.*; 2020.
152. de Cáceres, C., Martínez, R., Bachiller, P., Marín, L., García, J. The effect of tocilizumab on cytokine release syndrome in COVID-19 patients. *Pharmacol Rep*; 2020.
153. Della-Torre, E., Campochiaro, C., Cavalli, G., De Luca, G., Napolitano, A., La Marca, S., Boffini, N., Da Prat, V., Di Terlizzi, G., Lanzillotta, M. Interleukin-6 blockade with sarilumab in severe COVID-19 pneumonia with systemic hyperinflammation: an open-label cohort study. *Ann. Rheum. Dis.*; 2020.
154. Guaraldi, G., Meschiari, M., Cozzi-Lepri, A., Milic, J., Tonelli, R., Menozzi, M., Franceschini, E., Cuomo, G., Orlando, G., Borghi, V. Tocilizumab in patients with severe COVID-19: a retrospective cohort study. *The Lancet Rheumatology*; 2020.
155. Sanz Herrero, F., Puchades Gimeno, F., Ortega Garcia, P., Ferrer Gomez, C., Ocete Mochon, MD, Garcia Deltoro, M. Methylprednisolone added to tocilizumab reduces mortality in SARS-CoV-2 pneumonia: An observational study. *J. Intern. Med.*; 2020.
156. Jurado, A., Martín, M. C., Abad-Molina, C., Orduña, A., Martínez, A., Ocaña, E., Yarce, O., Navas, A. M., Trujillo, A., Fernández, L., Vergara, E., Rodríguez, B., Quirant, B., Martínez-Cáceres, E., Hernández, M., Perurena-Prieto, J., Gil, J., Cantenys, S., González-Martínez, G., Martínez-Saavedra, M. T., Rojo, R., Marco, F. M., Mora, S., Ontañón, J., López-Hoyos, M., Ocejo-Vinyals, G., Melero, J., Aguilar, M., Almeida, D., Medina, S., Vegas, M. C., Jiménez, Y., Prada, Á, Monzón, D., Boix, F., Cunill, V., Molina, J. COVID-19: Age, Interleukin-6, C-reactive protein, and lymphocytes as key clues from a multicentre retrospective study. *Immunity and Ageing*; 2020.
157. Kaminski, MA, Sunny, S, Balabayova, K, Kaur, A, Gupta, A, Abdallah, M, Quale, J. Tocilizumab therapy for COVID-19: A comparison of subcutaneous and intravenous therapies. *J. Glob. Infect. Dis.*; 2020.
158. Kewan, T, Covut, F, Al-Jaghbeer, MJ, Rose, L, Gopalakrishna, KV, Akbik, B. Tocilizumab for treatment of patients with severe COVID-19: A retrospective cohort study. *EClinicalMedicine*; 2020.
159. Luo, P., Liu, Y., Qiu, L., Liu, X., Liu, D., Li, J. Tocilizumab treatment in COVID-19: a single center experience. *J. Med. Virol.*; 2020.
160. Malekzadeh, R., Abedini, A., Mohsenpour, B., Sharifipour, E., Ghasemian, R., Javad-Mousavi, S., Khodashahi, R., Darban, M., Kalantari, S., Abdollahi, N. Subcutaneous tocilizumab in adults with severe and critical COVID-19: a prospective open-label uncontrolled multicenter trial. *Int. Immunopharmacol.*; 2020.
161. Masiá, M., Fernández-González, M., Padilla, S., Ortega, P., García, J., Agulló, V., García-Abellán, J., Telenti, G., Guillén, L., Gutiérrez, F. Impact of interleukin-6 blockade with tocilizumab on SARS-CoV-2 viral kinetics and antibody responses in patients with COVID-19: a prospective cohort study. *EBioMedicine*; 2020.
162. Mastroianni, A., Greco, S., Apuzzo, G., De Santis, S., Oriolo, C., Zanolini, A., Chidichimo, L., Vangeli, V. Subcutaneous tocilizumab treatment in patients with severe COVID-19-related cytokine release syndrome: an observational cohort study. *EClinicalMedicine*; 2020.
163. Meng, J., Xiao, G., Zhang, J., He, X., Ou, M., Bi, J., Yang, R., Di, W., Wang, Z., Li, Z., Gao, H., Liu, L., Zhang, G. Renin-angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension. *Emerg. microbes & infect.*; 2020.
164. Menzella, F., Fontana, M., Salvarani, C., Massari, M., Ruggiero, P., Scelfo, C., Barbieri, C., Castagnetti, C., Catellani, C., Gibellini, G. Efficacy of tocilizumab in patients with COVID-19 ARDS undergoing noninvasive ventilation. *Critical Care*; 2020.
165. Mikulska, M., Nicolini, L., Signori, A., Di Biagio, A., Sepulcri, C., Russo, C., Dettori, S., Berruti, M., Sormani, M., Giacobbe, D. Tocilizumab and steroid treatment in patients with COVID-19 pneumonia. *Plos one*; 2020.
166. Perrone, F., Piccirillo, M., Ascierto, P., Salvarani, C., Parrella, R., Marata, A., Popoli, P., Ferraris, L., Marrocco-Trischitta, M., Ripamonti, D. Tocilizumab for patients with COVID-19 pneumonia. The single-arm TOCIVID-19 prospective trial. *J Transl Med*; 2020.

167. Toniati, P., Piva, S., Cattalini, M., Garrafa, E., Regola, F., Castelli, F., Franceschini, F., Focà, E., Andreoli, L., Latronico, N. Tocilizumab for the treatment of severe COVID-19 pneumonia with hyperinflammatory syndrome and acute respiratory failure: A single center study of 100 patients in Brescia, Italy. *Autoimmun. Rev.*; 2020.
168. Price, C., Altice, F., Shyr, Y., Koff, A., Pischel, L., Goshua, G., Azar, M., Mcmanus, D., Chen, S., Gleeson, S. Tocilizumab treatment for cytokine release syndrome in hospitalized patients with coronavirus disease 2019: survival and clinical outcomes. *Chest*; 2020.
169. Ramiro, S., Mostard, RLM, Magro-Checa, C, Van Dongen, CMP, Dormans, T, Buijs, J, Gronenschild, M, De Kruif, MD, Van Haren, EHJ, Van Kraaij, T. Historically controlled comparison of glucocorticoids with or without tocilizumab versus supportive care only in patients with COVID-19-associated cytokine storm syndrome: results of the CHIC study. *Ann. Rheum. Dis.*; 2020.
170. Rossotti, R, Travi, G, Ughi, N, Corradin, M, Baiguera, C, Fumagalli, R, Bottiroli, M, Mondino, M, Merli, M, Bellone, A. Safety and efficacy of anti-il6-receptor tocilizumab use in severe and critical patients affected by coronavirus disease 2019: A comparative analysis. *J. Infect.*; 2020.
171. Sciascia, S, Aprà, F, Baffa, A, Baldovino, S, Boaro, D, Boero, R, Bonora, S, Calcagno, A, Cecchi, I, Cinnirella, G. Pilot prospective open, single-arm multicentre study on off-label use of tocilizumab in severe patients with COVID-19. *Clin Exp Rheumatol*; 2020.
172. Tomasiewicz, K., Piekarska, A., Stempkowska-Rejek, J., Serafińska, S., Gawkowska, A., Parczewski, M., Niścigorska-Olsen, J., Łapiński, T., Zarębska-Michaluk, D., Kowalska, J. Tocilizumab for patients with severe COVID-19: a retrospective, multi-center study. *Expert Rev Anti Infect Ther.*; 2020.
173. Tsai, A., Diawara, O., Nahass, R., Brunetti, L. Impact of tocilizumab administration on mortality in severe COVID-19. *Scientific reports*; 2020.
174. Borku, U., Ikitimur, H., Yavuzer, S. "Tocilizumab challenge: a series of cytokine storm therapy experience in hospitalized Covid-19 pneumonia patients". *J. Med. Virol.*; 2020.
175. Zhang, J., Hao, Y., Ou, W., Ming, F., Liang, G., Qian, Y., Cai, Q., Dong, S., Hu, S., Wang, W., Wei, S. Serum interleukin-6 is an indicator for severity in 901 patients with SARS-CoV-2 infection: a cohort study. *J Transl Med*; Oct 29 2020.
176. Milligan, P., Amsden, J., Norris, S., Baker, R., Myers, J., Preston, F., Gregory, J., Zoppi, K., Tayebi, K., Yeleti, R. Clinical outcomes in a cohort of non-ventilated COVID-19 patients with progressive hypoxemia and hyper-inflammatory response treated with baricitinib. *Lancet*; 2020.
177. Rodriguez-Garcia, Jose Luis, Sanchez-Nievas, Gines, Arevalo-Serrano, Juan, Garcia-Gomez, Cristina, Jimenez-Vizuite, Jose Maria, Martinez-Alfaro, Elisa. Baricitinib improves respiratory function in patients treated with corticosteroids for SARS-CoV-2 pneumonia: an observational cohort study. *Rheumatology (Oxford, England)*; 2020.
178. Titanji, B., Farley, M., Mehta, A., Connor-Schuler, R., Moanna, A., Cribbs, S., O'Shea, J., DeSilva, K., Chan, B., Edwards, A. Use of baricitinib in patients with moderate and severe COVID-19. *Clin. Infect. Dis*; 2020.
179. Bronte, V., Ugel, S., Tinazzi, E., Vella, A., De Sanctis, F., Canè, S., Batani, V., Trovato, R., Fiore, A., Petrova, V. Baricitinib restrains the immune dysregulation in patients with severe COVID-19. *J. Clin. Investig.*; 2020.
180. Cantini, F., Niccoli, L., Matarrese, D., Nicastrì, E., Stobbione, P., Goletti, D. Baricitinib therapy in COVID-19: a pilot study on safety and clinical impact. *J. Infect.*; 2020.
181. Administration, US Food and Drug. Fact sheet for healthcare providers: emergency use authorization (EUA) of casirivimab and imdevimab. 2020.
182. NIH. The COVID-19 Treatment Guidelines Panel's Statement on the Emergency Use Authorization of Bamlanivimab for the Treatment of COVID-19. 2020.
183. Administration, US Food and Drug. Fact sheet for healthcare providers: emergency use authorization (EUA) of bamlanivimab. 2020.
184. National Institute of Health (NIH). Therapeutic Management of Nonhospitalized Adults with COVID-19. February 1, 2022.
185. Administration, US Food & Drug. Coronavirus (COVID-19) update: FDA authorizes additional monoclonal antibody for treatment of COVID-19. May 26, 2021. 2021.
186. Administration, US Food & Drug, O'Shaughnessy, J. A. GSK Sotrovimab Letter Of Authorization 02232022. February 23, 2022.
187. GlaxoSmithKline LLC. Fact sheet for healthcare providers emergency use authorization (EUA) of sotrovimab. 2022.

188. López-Medina, E., López, P., Hurtado, I., Dávalos, D., Ramirez, O., Martínez, E., Díazgranados, J., Oñate, J., Chavarriaga, H., Herrera, S. Effect of ivermectin on time to resolution of symptoms among adults with mild COVID-19: a randomized clinical trial. *JAMA*; 2021.
189. Podder, C., Chowdhury, N., Sina, M., Haque, W. Outcome of ivermectin treated mild to moderate COVID-19 cases: a single-centre, open-label, randomised controlled study. *IMCIMS*; 2020.
190. Vanichkachorn, G., Newcomb, R., Cowl, C., Murad, M., Breeher, L., Miller, S., Trenary, M., Neveau, D., Higgins, S. Post-COVID-19 syndrome (long haul syndrome): description of a multidisciplinary clinic at Mayo Clinic and characteristics of the initial patient cohort. *Mayo Clin Proc*; Jul 2021.
191. Krolewiecki, A, Lifschitz, A, Moragas, M, Travacio, M, Valentini, R, Alonso, DF, Solari, R, Tinelli, MA, Cimino, RO, Álvarez, L. Antiviral effect of high-dose ivermectin in adults with COVID-19: A pilot randomised, controlled, open label, multicentre trial. 2020.
192. Niaee, M., Namdar, P., Allami, A., Zolghadr, L., Javadi, A., Karampour, A., Varnaseri, M., Bijani, B., Cheraghi, F., Naderi, Y. Ivermectin as an adjunct treatment for hospitalized adult COVID-19 patients: a randomized multi-center clinical trial. *Asian Pac J Trop Dis*; 2021.
193. Chowdhury, A., Shahbaz, M., Karim, M., Islam, J., Guo, D., He, S. A randomized trial of ivermectin-doxycycline and hydroxychloroquine-azithromycin therapy on COVID19 patients. 2020.
194. Hashim, HA, Maulood, MF, Rasheed, AM, Fatak, DF, Kabah, KK, Abdulamir, AS. Controlled randomized clinical trial on using ivermectin with doxycycline for treating COVID-19 patients in Baghdad, Iraq. *medRxiv*; 2020.
195. Elgazzar, A., Hany, B., Youssef, S., Hafez, M., Moussa, H. Efficacy and safety of ivermectin for treatment and prophylaxis of COVID-19 pandemic. 2020.
196. Kory, P., Meduri, G., Varon, J., Iglesias, J., Marik, P. Review of the emerging evidence demonstrating the efficacy of ivermectin in the prophylaxis and treatment of COVID-19. *Am J Ther*; Apr 22 2021.
197. Caly, L., Druce, J., Catton, M., Jans, D., Wagstaff, K. The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro. *Antiviral Res*; Jun 2020.
198. Joyner, MJ., Bruno, KA., Klassen, SA., Kunze, KL., Johnson, PW., Lesser, ER., Wiggins, CC., Senefeld, JW., Klompas, AM., Hodge, DO., Shepherd, JRA, Rea, RF., Whelan, ER., Clayburn, AJ., Spiegel, MR., Baker, SE., Larson, KF., Ripoll, JG., Andersen, KJ., Buras, MR., Vogt, MNP, Herasevich, V, Dennis, JJ., Regimbal, RJ., Bauer, PR., Blair, JE., van Buskirk, CM., Winters, JL., Stubbs, JR., van Helmond, N, Butterfield, BP., Sexton, MA., Diaz Soto, JC., Paneth, NS., Verdun, NC., Marks, P, Casadevall, A, Fairweather, DL, Carter, RE., Wright, R.S. Safety update: COVID-19 convalescent plasma in 20,000 hospitalized patients. *Mayo Clin. Proc.*; 2020/09/01/.
199. Joyner, MJ, Senefeld, JW, Klassen, SA, Mills, JR, Johnson, PW, Theel, ES, Wiggins, CC, Bruno, KA, Klompas, AM, Lesser, ER, Kunze, KL, Sexton, MA, Diaz Soto, JC, Baker, SE, Shepherd, JRA, van Helmond, N, van Buskirk, CM., Winters, JL, Stubbs, JR, Rea, RF, Hodge, DO, Herasevich, V, Whelan, ER, Clayburn, AJ, Larson, KF, Ripoll, JG, Andersen, KJ, Buras, MR, Vogt, MNP, Dennis, JJ, Regimbal, RJ, Bauer, PR, Blair, JE, Paneth, NS, Fairweather, DL, Wright, R.S, Carter, RE, Casadevall, A. Effect of convalescent plasma on mortality among hospitalized patients with COVID-19: Initial three-month experience. *medRxiv*; 2020.
200. Joyner, MJ, Wright, RS, Fairweather, DL, Senefeld, JW, Bruno, KA, Klassen, SA, Carter, RE, Klompas, AM, Wiggins, CC, Shepherd, JRA. Early safety indicators of COVID-19 convalescent plasma in 5,000 patients. *J. Clin. Investig.*; 2020.
201. Salazar, E, Christensen, PA., Graviss, EA., Nguyen, DT., Castillo, B, Chen, J, Lopez, BV., Eagar, TN., Yi, X, Zhao, P, Rogers, J, Shehabeldin, A, Joseph, D, Leveque, C, Olsen, RJ., Bernard, DW., Gollihar, J, Musser, JM. Treatment of coronavirus disease 2019 patients with convalescent plasma reveals a signal of significantly decreased mortality. *Am. J. Clin. Pathol.*; 2020/11/01/.
202. Wang, Y, Zhang, L, Sang, L, Ye, F, Ruan, S, Zhong, B, Song, T, Alshukairi, A, Chen, R, Zhang, Z. Kinetics of viral load and antibody response in relation to COVID-19 severity. *J. Clin. Investig.*; 2020.
203. Salazar, E, Kuchipudi, SV., Christensen, PA., Eagar, TN., Yi, X, Zhao, P, Jin, Z, Long, SW, Olsen, RJ., Chen, J, Castillo, B, Leveque, C, Towers, DM., Lavinder, J, Gollihar, JD., Cardona, J, Ippolito, GC., Nissly, RH., Bird, IM., Greenawalt, D, Rossi, RM., Gontu, A, Srinivasan, S, Poojary, IB., Cattadori, IM., Hudson, PJ., Joselyn, N, Prugar, L, Huie, K, Herbert, A, Bernard, DW., Dye, J, Kapur, V, Musser, JM. Relationship between anti-spike protein antibody titers and SARS-CoV-2 &emph;in Vitro&emph;; virus neutralization in convalescent plasma. *bioRxiv*; 2020.
204. Olivares-Gazca, J., Priesca-Marín, J., Ojeda-Laguna, M., Garcés-Eisele, J., Soto-Olvera, S., Palacios-Alonso, A., Izquierdo-Vega, J., Chacon-Cano, R., Arizpe-Bravo, D., López-Trujillo, M. Infusion of convalescent plasma is associated with clinical improvement in critically ill patients with COVID-19: a pilot study. *Rev Invest Clin*; 2020.
205. Nora, H., Philippos, E., Marcel, A., Cornelius, D., Dunja, B., Ortwin, A., Jan-Steffen, K., Petra, B. Assessment of SARS-CoV-2 in human semen-a cohort study. *Fertil Steril*; 2020.

206. Klein, SL, Pekosz, A, Park, HS, Ursin, RL, Shapiro, JR, Benner, SE, Littlefield, K, Kumar, S, Naik, HM, Betenbaugh, MJ. Sex, age, and hospitalization drive antibody responses in a COVID-19 convalescent plasma donor population. *J. Clin. Investig.*; 2020.
207. Isho, B., Abe, K., Zuo, M., Jamal, A., Rathod, B., Wang, J., Li, Z., Chao, G., Rojas, O., Bang, Y., Pu, A., Christie-Holmes, N., Gervais, C., Ceccarelli, D., Samavarchi-Tehrani, P., Guvenc, F., Budylowski, P., Li, A., Paterson, A., Yue, F., Marin, L., Caldwell, L., Wrana, J., Colwill, K., Sicheri, F., Mubareka, S., Gray-Owen, S., Drews, S., Siqueira, W., Barrios-Rodiles, M., Ostrowski, M., Rini, J., Durocher, Y., McGeer, A., Gommerman, J., Gingras, A. Persistence of serum and saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. *Sci. Immunol.*; 2020.
208. Hansen, J, Baum, A, Pascal, KE, Russo, V, Giordano, S, Wloga, E, Fulton, BO, Yan, Y, Koon, K, Patel, K. Studies in humanized mice and convalescent humans yield a SARS-CoV-2 antibody cocktail. *Science*; 2020.
209. Faqihi, F., Alharthy, A., Alodat, M., Kutsogiannis, D., Brindley, P., Karakitsos, D. Therapeutic plasma exchange in adult critically ill patients with life-threatening SARS-CoV-2 disease: a pilot study. *J. Crit. Care*; 2020.
210. Bradfute, S., Hurwitz, I., Yingling, A., Ye, C., Cheng, Q., Noonan, T., Raval, J., Sosa, N., Mertz, G., Perkins, D., Harkins, M. Severe acute respiratory syndrome coronavirus 2 neutralizing antibody titers in convalescent plasma and recipients in New Mexico: an open treatment study in patients with coronavirus disease 2019. *J. Infect. Dis.*; 2020.
211. Liu, S., Lin, H., Baine, I., Wajnberg, A., Gumprecht, J., Rahman, F., Rodriguez, D., Tandon, P., Bassily-Marcus, A., Bander, J., Sanky, C., Dupper, A., Zheng, A., Nguyen, F., Amanat, F., Stadlbauer, D., Altman, D., Chen, B., Krammer, F., Mendu, D., Firpo-Betancourt, A., Levin, M., Bagiella, E., Casadevall, A., Cordon-Cardo, C., Jhang, J., Arinsburg, S., Reich, D., Aberg, J., Bouvier, N. Convalescent plasma treatment of severe COVID-19: a propensity score-matched control study. *Nature Med*; 2020/11/01.
212. Wang, X, Guo, X, Xin, Q, Pan, Y, Hu, Y, Li, J, Chu, Y, Feng, Y, Wang, Q. Neutralizing antibody responses to severe acute respiratory syndrome coronavirus 2 in coronavirus disease 2019 inpatients and convalescent patients. *Clin Infect Dis*; 2020.
213. Gharbharan, A, Jordans, CCE, GeurtsvanKessel, C, den Hollander, JG, Karim, F, Mollema, FPN, Stalenhoef, JE, Dofferhoff, A, Ludwig, I, Koster, A. Convalescent plasma for COVID-19. A randomized clinical trial. *MEDRxiv*; 2020.
214. Tian, X., Li, C., Huang, A., Xia, S., Lu, S., Shi, Z., Lu, L., Jiang, S., Yang, Z., Wu, Y., Ying, T. Potent binding of 2019 novel coronavirus spike protein by a SARS coronavirus-specific human monoclonal antibody. *Emerg Microbes Infect*; 2020.
215. Wrapp, D., Wang, N., Corbett, K. S., Goldsmith, J. A., Hsieh, C. L., Abiona, O., Graham, B. S., McLellan, J. S. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science*; Mar 13 2020.
216. Hung, I., To, K., Lee, C., Lee, K., Chan, K., Yan, W., Liu, R., Watt, C., Chan, W., Lai, K. Convalescent plasma treatment reduced mortality in patients with severe pandemic influenza A (H1N1) 2009 virus infection. *Arch. Clin. Infect. Dis.*; 2011.
217. Winck, J, Ambrosino, N. COVID-19 pandemic and non invasive respiratory management: every Goliath needs a David. An evidence based evaluation of problems. *Pulmonology*; 2020.
218. Eslami, G., Mousaviasl, S., Radmanesh, E., Jelvay, S., Bitaraf, S., Simmons, B., Wentzel, H., Hill, A., Sadeghi, A., Freeman, J., Salmanzadeh, S., Esmaeilian, H., Mobarak, M., Tabibi, R., Jafari Kashi, A. H., Lotfi, Z., Talebzadeh, S. M., Wickramatillake, A., Momtazan, M., Hajizadeh Farsani, M., Marjani, S., Mobarak, S. The impact of sofosbuvir/daclatasvir or ribavirin in patients with severe COVID-19. *J Antimicrob Chemother*; Nov 1 2020.
219. Gong, WJ, Zhou, T, Wu, SL, Ye, JL, Xu, JQ, Zeng, F, Su, YY, Han, Y, Lv, YN, Zhang, Y. Clinical efficacy of ribavirin in adults hospitalized with severe Covid-19: A retrospective analysis of 208 patients. 2020.
220. Tong, S., Su, Y., Yu, Y., Wu, C., Chen, J., Wang, S., Jiang, J. Ribavirin therapy for severe COVID-19: a retrospective cohort study. *Int J Antimicrob Agents*; Sep 2020.
221. Cheng, C., Lee, Y., Chen, C., Lin, Y., Liu, C., Liao, C., Cheng, S. Lopinavir/ritonavir did not shorten the duration of SARS CoV-2 shedding in patients with mild pneumonia in Taiwan. *J Microbiol Immunol Infect*; 2020.
222. Jothimani, D, Kailasam, E, Danielraj, S, Nallathambi, B, Ramachandran, H, Sekar, P, Manoharan, S, Ramani, V, Narasimhan, G, Kaliamoorthy, I. COVID-19: Poor outcomes in patients with zinc deficiency. *J. Glob. Infect. Dis.*; 2020.
223. Yasui, Y, Yasui, H, Suzuki, K, Saitou, T, Yamamoto, Y, Ishizaka, T, Nishida, K, Yoshihara, S, Gohma, I, Ogawa, Y. Analysis of the predictive factors for a critical illness of COVID-19 during treatment-relationship between serum zinc level and critical illness of COVID-19-. *Int. J. Infect. Dis.*; 2020.
224. Vogel-González, M, Talló-Parra, M, Herrera-Fernández, V, Pérez-Vilaró, G, Chillón, M, Nogués, X, Gómez-Zorrilla, S, López-Montesinos, I, Villar, J, Sorlí-Redó, L. Low zinc levels at clinical admission associates with poor outcomes in COVID-19. 2020.
225. Carlucci, P., Ahuja, T., Petrilli, C., Rajagopalan, H., Jones, S., Rahimian, J. Hydroxychloroquine and azithromycin plus zinc vs hydroxychloroquine and azithromycin alone: outcomes in hospitalized COVID-19 patients. *medRxiv*; 2020.

226. Frontera, JA, Rahimian, JO, Yaghi, S, Liu, M, Lewis, A, de Havenon, A, Mainali, S, Huang, J, Scher, E, Wisniewski, T. Treatment with zinc is associated with reduced in-hospital mortality among COVID-19 patients: A multi-center cohort study. *Res Sq*; 2020.
227. Butler-Laporte, Guillaume, Nakanishi, Tomoko, Mooser, Vincent, Morrison, David R, Abdullah, Tala, Adeleye, Olumide, Mamlook, Noor, Kimchi, Nofar, Afrasiabi, Zaman, Rezk, Nardin. Vitamin D and Covid-19 Susceptibility and Severity: a Mendelian Randomization Study. *medRxiv*; 2020.
228. Hastie, CE, Mackay, DF, Ho, F, Celis-Morales, CA, Katikireddi, SV, Niedzwiedz, CL, Jani, BD, Welsh, P, Mair, FS, Gray, SR. Vitamin D concentrations and COVID-19 infection in UK Biobank. *Diabetes Metab Syndr*; 2020.
229. Raisi-Estabragh, Z, McCracken, C, Bethell, MS, Cooper, J, Cooper, C, Caulfield, MJ, Munroe, PB, Harvey, NC, Petersen, SE. Greater risk of severe COVID-19 in black, asian and minority ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25 (OH)-vitamin D status: study of 1326 cases from the UK Biobank. *J Public Health*; 2020.
230. De Smet, D., De Smet, K., Herroelen, P., Gryspeerdt, S., Martens, G. Vitamin D deficiency as risk factor for severe COVID-19: a convergence of two pandemics. *MedRxiv*; 2020.
231. Darling, A., Ahmadi, K., Ward, K., Harvey, N., Alves, A., Dunn-Waters, D., Lanham-New, S., Cooper, C., Blackbourn, D. Vitamin D status, body mass index, ethnicity and COVID-19: Initial analysis of the first-reported UK Biobank COVID-19 positive cases (n 580) compared with negative controls (n 723). *MedRxiv*; 2020.
232. Brenner, H., Holleczer, B., Schöttker, B. Vitamin D insufficiency and deficiency and mortality from respiratory diseases in a cohort of older adults: potential for limiting the death toll during and beyond the COVID-19 pandemic?. *Nutrients*; 2020.
233. Tan, C., Ho, L., Kalimuddin, S., Chong, B., Teh, Y., Thien, S., Wong, H., Tern, P., Chay, J., Nagarajan, C. A cohort study to evaluate the effect of combination vitamin D, magnesium and vitamin B12 (DMB) on progression to severe outcome in older COVID-19 patients. *medRxiv*; 2020.
234. Raharusun, P. Patterns of COVID-19 mortality and vitamin D: An Indonesian study. *Available at SSRN 3585561*; 2020.
235. Radujkovic, A, Hippchen, T, Tiwari-Heckler, S, Dreher, S, Boxberger, M, Merle, U. Vitamin D deficiency and outcome of COVID-19 patients. *Nutrients*; 2020.
236. Pizzini, A., Aichner, M., Sahanic, S., Böhm, A., Egger, A., Hoermann, G., Kurz, K., Widmann, G., Bellmann-Weiler, R., Weiss, G. Impact of vitamin d deficiency on covid-19—a prospective analysis from the covid registry. *Nutrients*; 2020.
237. Meltzer, D., Best, T., Zhang, H., Vokes, T., Arora, V., Solway, J. Association of vitamin D status and other clinical characteristics with COVID-19 test results. *JAMA Netw. Open*; 2020.
238. Merzon, E., Tworowski, D., Gorohovski, A., Vinker, S., Golan Cohen, A., Green, I., Frenkel-Morgenstern, M. Low plasma 25 (OH) vitamin D level is associated with increased risk of COVID-19 infection: an Israeli population-based study. *FEBS J.*; 2020.
239. Maghbooli, Z., Sahraian, M., Ebrahimi, M., Pazoki, M., Kafan, S., Tabriz, H., Hadadi, A., Montazeri, M., Nasiri, M., Shirvani, A. Vitamin D sufficiency, a serum 25-hydroxyvitamin D at least 30 ng/mL reduced risk for adverse clinical outcomes in patients with COVID-19 infection. *PloS one*; 2020.
240. Macaya, Fernando, Paeres, Carolina Espejo, Carbó, Adrián Valls, Fernández-Ortiz, Antonio, Del Castillo, Juan González, Sánchez, Francisco Javier Martín, de la Vega, Isabelle Runkle, Herrera, Miguel Angel Rubio. Interaction between age and vitamin D deficiency in severe COVID-19 infection. *Nutrición hospitalaria: Organo oficial de la Sociedad española de nutrición parenteral y enteral*; 2020.
241. Kaufman, HW, Niles, JK, Kroll, MH, Bi, C, Holick, MF. SARS-CoV-2 positivity rates associated with circulating 25-hydroxyvitamin D levels. *PloS one*; 2020.
242. Jain, A., Chaurasia, R., Sengar, N., Singh, M., Mahor, S., Narain, S. Analysis of vitamin D level among asymptomatic and critically ill COVID-19 patients and its correlation with inflammatory markers. *Sci. Rep.*; 2020.
243. Ilie, P., Stefanescu, S., Smith, L. The role of vitamin D in the prevention of coronavirus disease 2019 infection and mortality. *Aging Clin Exp Res*; 2020.
244. Hernández, JL, Nan, D, Fernandez-Ayala, M, García-Unzueta, M, Hernández-Hernández, MA, López-Hoyos, M, Muñoz-Cacho, P, Olmos, JM, Gutiérrez-Cuadra, M, Ruiz-Cubillán, JJ. Vitamin D status in hospitalized patients with SARS-CoV-2 infection. *J. Clin. Endocrinol*; 2020.
245. LaFond, E., Weidman, K., Lief, L. Care of the postcoronavirus disease 2019 patient. *Curr Opin Pulm Med*; 2021.
246. Ayoubkhani, D., Khunti, K., Nafilyan, V., Maddox, T., Humberstone, B., Diamond, I., Banerjee, A. Epidemiology of post-COVID syndrome following hospitalisation with coronavirus: a retrospective cohort study. *MedRxiv*; 2021.

247. Halpin, Stephen J, McIvor, Claire, Whyatt, Gemma, Adams, Anastasia, Harvey, Olivia, McLean, Lyndsay, Walshaw, Christopher, Kemp, Steven, Corrado, Joanna, Singh, Rajinder. Postdischarge symptoms and rehabilitation needs in survivors of COVID-19 infection: A cross-sectional evaluation. *Journal of medical virology*; 2021.
248. Maley, J., Alba, G., Barry, J., Bartels, M., Fleming, T., Oleson, C., Rydberg, L., Sampsel, S., Silver, J., Sipes, S. Multi-disciplinary collaborative consensus guidance statement on the assessment and treatment of breathing discomfort and respiratory sequelae in patients with post-acute sequelae of SARS-CoV-2 infection (PASC). *PM&R*; 2022.
249. Fumagalli, A., Misuraca, C., Bianchi, A., Borsa, N., Limonta, S., Maggiolini, S., Bonardi, D. R., Corsonello, A., Di Rosa, M., Soraci, L., Lattanzio, F., Colombo, D. Pulmonary function in patients surviving to COVID-19 pneumonia. *Infection*; Feb 2021.
250. Mo, Xiaoneng, Jian, Wenhua, Su, Zhuquan, Chen, Mu, Peng, Hui, Peng, Ping, Lei, Chunliang, Chen, Ruchong, Zhong, Nanshan, Li, Shiyue. Abnormal pulmonary function in COVID-19 patients at time of hospital discharge. *The European respiratory journal*; 2020.
251. Akhilesh, Kunoor, Aditya, Ashok, Merlin, Moni, Asmita, Mehta, Elizabeth Mary, Thomas, Veena, Viswan, Harikrishnan, K. A case series on post-COVID pulmonary rehabilitation: Early experiences from Kerala, South India. *Indian Journal of Case Reports*; 02/09 2021.
252. McCarthy, B., Casey, D., Devane, D., Murphy, K., Murphy, E., Lacasse, Y. Pulmonary rehabilitation for chronic obstructive pulmonary disease. *Cochrane Database of Systematic Reviews*; 2015.
253. Hill, N. S. Pulmonary rehabilitation. *Proc Am Thorac Soc*; 2006.
254. Cheng, H. H., Chou, W. Rehabilitation can reduce mortality rate in patients who were intubated due to pneumonia. *Annals of Physical and Rehabilitation Medicine*; 2018/07/01/.
255. Dowman, L., Hill, C. J., May, A., Holland, A. E. Pulmonary rehabilitation for interstitial lung disease. *Cochrane Database of Systematic Reviews*; 2021.
256. Lau, H. M., Ng, G. Y., Jones, A. Y., Lee, E. W., Siu, E. H., Hui, D. S. A randomised controlled trial of the effectiveness of an exercise training program in patients recovering from severe acute respiratory syndrome. *Aust J Physiother*; 2005.
257. Spruit, Martijn A, Singh, Sally J, Garvey, Chris, ZuWallack, Richard, Nici, Linda, Rochester, Carolyn, Hill, Kylie, Holland, Anne E, Lareau, Suzanne C, Man, William D-C. An official American Thoracic Society/European Respiratory Society statement: key concepts and advances in pulmonary rehabilitation. *American journal of respiratory and critical care medicine*; 2013.
258. Guo, Tao, Fan, Yongzhen, Chen, Ming, Wu, Xiaoyan, Zhang, Lin, He, Tao, Wang, Hairong, Wan, Jing, Wang, Xinghuan, Lu, Zhibing. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA cardiology*; 2020.
259. Lindner, Diana, Fitzek, Antonia, Bräuninger, Hanna, Aleshcheva, Ganna, Edler, Caroline, Meissner, Kira, Scherschel, Katharina, Kirchhof, Paulus, Escher, Felicitas, Schultheiss, Heinz-Peter. Association of Cardiac Infection With SARS-CoV-2 in Confirmed COVID-19 Autopsy Cases. *JAMA cardiology*; 2020.
260. Driggin, Elissa, Madhavan, Mahesh V., Bikdeli, Behnood, Chuich, Taylor, Laracy, Justin, Biondi-Zoccai, Giuseppe, Brown, Tyler S., Der Nigoghossian, Caroline, Zidar, David A., Haythe, Jennifer, Brodie, Daniel, Beckman, Joshua A., Kirtane, Ajay J., Stone, Gregg W., Krumholz, Harlan M., Parikh, Sahil A. Cardiovascular Considerations for Patients, Health Care Workers, and Health Systems During the COVID-19 Pandemic. *Journal of the American College of Cardiology*; 2020.
261. Puntmann, Valentina O., Carerj, M. Ludovica, Wieters, Imke, Fahim, Masia, Arendt, Christophe, Hoffmann, Jedrzej, Shchendrygina, Anastasia, Escher, Felicitas, Vasa-Nicotera, Mariuca, Zeiher, Andreas M., Vehreschild, Maria, Nagel, Eike. Outcomes of Cardiovascular Magnetic Resonance Imaging in Patients Recently Recovered From Coronavirus Disease 2019 (COVID-19). *JAMA Cardiology*; 2020.
262. Madjid, M., Safavi-Naeini, P., Solomon, S. D., Vardeny, O. Potential Effects of Coronaviruses on the Cardiovascular System: A Review. *JAMA Cardiol*; Jul 1 2020.
263. Kochi, A. N., Tagliari, A. P., Forleo, G. B., Fassini, G. M., Tondo, C. Cardiac and arrhythmic complications in patients with COVID-19. *J Cardiovasc Electrophysiol*; May 2020.
264. Ng, M. Y., Ferreira, V. M., Leung, S. T., Yin Lee, J. C., Ho-Tung Fong, A., To Liu, R. W., Man Chan, J. W., Wu, A. K. L., Lung, K. C., Crean, A. M., Fan-Ngai Hung, I., Siu, C. W. Patients Recovered From COVID-19 Show Ongoing Subclinical Myocarditis as Revealed by Cardiac Magnetic Resonance Imaging. *JACC Cardiovasc Imaging*; Nov 2020.
265. Hermann, M., Pekacka-Egli, A. M., Witassek, F., Baumgaertner, R., Schoendorf, S., Spielmanns, M. Feasibility and Efficacy of Cardiopulmonary Rehabilitation After COVID-19. *Am J Phys Med Rehabil*; Oct 2020.
266. Cowie, Aynsley, Buckley, John, Doherty, Patrick, Furze, Gill, Hayward, Jo, Hinton, Sally, Jones, Jennifer, Speck, Linda, Dalal, Hasnain, Mills, Joseph. Standards and core components for cardiovascular disease prevention and rehabilitation. *Heart*; 2019.

267. Dalal, Hasnain M, Doherty, Patrick, Taylor, Rod S. Cardiac rehabilitation. *Bmj*; 2015.
268. Piepoli, M. F., Corrà, U., Adamopoulos, S., Benzer, W., Bjarnason-Wehrens, B., Cupples, M., Dendale, P., Doherty, P., Gaita, D., Höfer, S., McGee, H., Mendes, M., Niebauer, J., Pogosova, N., Garcia-Porrero, E., Rauch, B., Schmid, J. P., Giannuzzi, P. Secondary prevention in the clinical management of patients with cardiovascular diseases. Core components, standards and outcome measures for referral and delivery: a policy statement from the cardiac rehabilitation section of the European Association for Cardiovascular Prevention & Rehabilitation. Endorsed by the Committee for Practice Guidelines of the European Society of Cardiology. *Eur J Prev Cardiol*; Jun 2014.
269. Balady, G. J., Williams, M. A., Ades, P. A., Bittner, V., Comoss, P., Foody, J. M., Franklin, B., Sanderson, B., Southard, D. Core components of cardiac rehabilitation/secondary prevention programs: 2007 update: a scientific statement from the American Heart Association Exercise, Cardiac Rehabilitation, and Prevention Committee, the Council on Clinical Cardiology; the Councils on Cardiovascular Nursing, Epidemiology and Prevention, and Nutrition, Physical Activity, and Metabolism; and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*; May 22 2007.
270. Leon, A. S., Franklin, B. A., Costa, F., Balady, G. J., Berra, K. A., Stewart, K. J., Thompson, P. D., Williams, M. A., Lauer, M. S. Cardiac rehabilitation and secondary prevention of coronary heart disease: an American Heart Association scientific statement from the Council on Clinical Cardiology (Subcommittee on Exercise, Cardiac Rehabilitation, and Prevention) and the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity), in collaboration with the American association of Cardiovascular and Pulmonary Rehabilitation. *Circulation*; Jan 25 2005.
271. Kortebein, P. Rehabilitation for hospital-associated deconditioning. *Am J Phys Med Rehabil*; Jan 2009.
272. Galloway, R. V., Karmarkar, A. M., Graham, J. E., Tan, A., Raji, M., Granger, C. V., Ottenbacher, K. J. Hospital Readmission Following Discharge From Inpatient Rehabilitation for Older Adults With Debility. *Phys Ther*; Feb 2016.
273. Timmer, A. J., Unsworth, C. A., Taylor, N. F. Rehabilitation interventions with deconditioned older adults following an acute hospital admission: a systematic review. *Clin Rehabil*; Nov 2014.
274. Herrera, J., Niehaus, W., Whiteson, J., Azola, A., Baratta, J., Fleming, T., Kim, S., Naqvi, H., Sampsel, S., Silver, J., Gutierrez, M., Maley, J., Herman, E., Abramoff, B. Multidisciplinary collaborative consensus guidance statement on the assessment and treatment of fatigue in postacute sequelae of SARS-CoV-2 infection (PASC) patients. *PM&R*; Sep 2021.
275. Barker-Davies, Robert M, O'Sullivan, Oliver, Senaratne, Kahawalage Pumi Prathima, Baker, Polly, Cranley, Mark, Dharm-Datta, Shreshth, Ellis, Henrietta, Goodall, Duncan, Gough, Michael, Lewis, Sarah. The Stanford Hall consensus statement for post-COVID-19 rehabilitation. *British journal of sports medicine*; 2020.
276. Alexander, T., Thomson, V. S., Malviya, A., Mohan, B., Wander, G. S., S, H., Seth, S., Reddy, S., Arulraj, S., Shah, S., Joshi, S., Tiwaskar, M., Nadkar, M., Tewary, K. Guidance for Health Care Providers on Management of Cardiovascular Complications in Patients Suspected or Confirmed with COVID 19 Virus Infection. *J Assoc Physicians India*; May 2020.
277. Johnson, Joshua K, Lapin, Brittany, Green, Karen, Stilphen, Mary. Frequency of Physical Therapist Intervention Is Associated With Mobility Status and Disposition at Hospital Discharge for Patients With COVID-19. *Physical Therapy*; 2021.
278. Sun, Tiantian, Guo, Liyun, Tian, Fei, Dai, Tiantian, Xing, Xiaohong, Zhao, Junqing, Li, Qiang. Rehabilitation of patients with COVID-19. *Expert Review of Respiratory Medicine*; 2020.
279. Rooney, Scott, Webster, Amy, Paul, Lorna. Systematic Review of Changes and Recovery in Physical Function and Fitness After Severe Acute Respiratory Syndrome-Related Coronavirus Infection: Implications for COVID-19 Rehabilitation. *Physical Therapy*; 2020.
280. Vickory, F., Ridgeway, K., Falvey, J., Houwer, B., Gunlikson, J., Payne, K., Niehaus, W. Safety, feasibility, and outcomes of frequent, long-duration rehabilitation in an inpatient rehabilitation facility after prolonged hospitalization for severe COVID-19: an observational study. *Phys Ther*; Nov 1 2021.
281. CDC. What is ME/CFS?. 2021.
282. Chilazi, M., Duffy, E., Thakkar, A., Michos, E. COVID and cardiovascular disease: what we know in 2021. *Curr Atheroscler Rep*; May 13 2021.
283. Dayrit, J., Verdusco-Gutierrez, M., Teal, A., Shah, S. Enhanced external counterpulsation as a novel treatment for post-acute COVID-19 sequelae. *Cureus*; Apr 7 2021.
284. Stefano, G. Historical insight into infections and disorders associated with neurological and psychiatric sequelae similar to long COVID. *Med Sci Monit*; Feb 26 2021.
285. Stefano, G., Büttiker, P., Weissenberger, S., Martin, A., Ptacek, R., Kream, R. Editorial: the pathogenesis of long-term neuropsychiatric COVID-19 and the role of microglia, mitochondria, and persistent neuroinflammation: a hypothesis. *Med Sci Monit*; May 10 2021.
286. Theoharides, T., Cholevas, C., Polyzoidis, K., Politis, A. Long-COVID syndrome-associated brain fog and chemofog: Luteolin to the rescue. *Biofactors*; Mar 2021.

287. Theoharides, T., Conti, P. COVID-19 and multisystem inflammatory syndrome, or is it mast cell activation syndrome. *J Biol Regul Homeost Agents*; 2020.
288. Stefano, G., Ptacek, R., Ptackova, H., Martin, A., Kream, R. Selective neuronal mitochondrial targeting in SARS-CoV-2 infection affects cognitive processes to induce 'brain fog' and results in behavioral changes that favor viral survival. *Med Sci Monit*; Jan 25 2021.
289. Østergaard, L. SARS CoV-2 related microvascular damage and symptoms during and after COVID-19: Consequences of capillary transit-time changes, tissue hypoxia and inflammation. *Physiol Rep*; Feb 2021.
290. Kingstone, T., Taylor, A., O'Donnell, C., Atherton, H., Blane, D., Chew-Graham, C. Finding the 'right' GP: a qualitative study of the experiences of people with long-COVID. *BJGP Open*; Dec 2020.
291. Novak, P. Post COVID-19 syndrome associated with orthostatic cerebral hypoperfusion syndrome, small fiber neuropathy and benefit of immunotherapy: a case report. *eNeurologicalSci*; Dec 2020.
292. Hellmuth, J., Barnett, T., Asken, B., Kelly, J., Torres, L., Stephens, M., Greenhouse, B., Martin, J., Chow, F., Deeks, S., Greene, M., Miller, B., Annan, W., Henrich, T., Peluso, M. Persistent COVID-19-associated neurocognitive symptoms in non-hospitalized patients. *J Neurovirol*; Feb 2021.
293. Graham, E., Clark, J., Orban, Z., Lim, P., Szymanski, A., Taylor, C., DiBiase, R., Jia, D., Balabanov, R., Ho, S., Batra, A., Liotta, E., Korolnik, I. Persistent neurologic symptoms and cognitive dysfunction in non-hospitalized Covid-19 "long haulers". *Ann Clin Transl Neurol*; May 2021.
294. Humphreys, H., Kilby, L., Kudiersky, N., Copeland, R. Long COVID and the role of physical activity: a qualitative study. *BMJ Open*; Mar 10 2021.
295. Gulick, S., Mandel, S., Maitz, E. Special report: cognitive screening after COVID-19. *Practical Neurology*; 2021.
296. Havervall, S., Rosell, A., Phillipson, M., Mangsbo, S., Nilsson, P., Hober, S., Thålin, C. Symptoms and functional impairment assessed 8 months after mild COVID-19 among health care workers. *JAMA*; 2021.
297. Rogers, J., David, A. A longer look at COVID-19 and neuropsychiatric outcomes. *The Lancet Psychiatry*; 2021.
298. Taquet, M., Geddes, J., Husain, M., Luciano, S., Harrison, P. 6-month neurological and psychiatric outcomes in 236 379 survivors of COVID-19: a retrospective cohort study using electronic health records. *Lancet Psychiatry*; 2021.
299. Taquet, M., Luciano, S., Geddes, J., Harrison, P. Bidirectional associations between COVID-19 and psychiatric disorder: retrospective cohort studies of 62 354 COVID-19 cases in the USA. *Lancet Psychiatry*; 2021.
300. Boivin, M. J., Nakasujja, N., Sikorskii, A., Opoka, R. O., Giordani, B. A Randomized Controlled Trial to Evaluate if Computerized Cognitive Rehabilitation Improves Neurocognition in Ugandan Children with HIV. *AIDS Res Hum Retroviruses*; Aug 2016.
301. Boivin, M. J., Nakasujja, N., Sikorskii, A., Ruiseñor-Escudero, H., Familiar-Lopez, I., Walhof, K., van der Lugt, E. M., Opoka, R. O., Giordani, B. Neuropsychological benefits of computerized cognitive rehabilitation training in Ugandan children surviving severe malaria: A randomized controlled trial. *Brain Res Bull*; Feb 2019.
302. Fine, J., Ambrose, A., Didehbani, N., Fleming, T., Glashan, L., Longo, M., Merlino, A., Ng, R., Nora, G., Rolin, S., Silver, J., Terzic, C., Verduzco-Gutierrez, M., Sampsel, S. Multi-disciplinary collaborative consensus guidance statement on the assessment and treatment of cognitive symptoms in patients with post-acute sequelae of SARS-CoV-2 infection (PASC). *PM&R*; Jan 2022.
303. Carfi, Angelo, Bernabei, Roberto, Landi, Francesco. Persistent symptoms in patients after acute covid-19. *JAMA*; 2020.
304. Linden, Wolfgang. Psychological treatments in cardiac rehabilitation: review of rationales and outcomes. *Journal of Psychosomatic Research*; 2000/04/01/.
305. Kemp, Harriet I., Corner, Eve, Colvin, Lesley A. Chronic pain after COVID-19: implications for rehabilitation. *British Journal of Anaesthesia*; 2020.
306. Li, Jinzhi, Li, Xiuchuan, Jiang, Jie, Xu, Xuexue, Wu, Jing, Xu, Yuanyuan, Lin, Xin, Hall, John, Xu, Huashan, Xu, Jincheng. The Effect of Cognitive Behavioral Therapy on Depression, Anxiety, and Stress in Patients With COVID-19: A Randomized Controlled Trial. *Frontiers in Psychiatry*; 2020.
307. Liu, Kai, Chen, Ying, Wu, Duoqi, Lin, Ruzheng, Wang, Zaisheng, Pan, Liqing. Effects of progressive muscle relaxation on anxiety and sleep quality in patients with COVID-19. *Complementary Therapies in Clinical Practice*; 2020.
308. Wei, Ning, Huang, Bo-Chao, Lu, Shao-Jia, Hu, Jian-Bo, Zhou, Xiao-Yi, Hu, Chan-Chan, Chen, Jing-Kai, Huang, Jin-Wen, Li, Shu-Guang, Wang, Zheng. Efficacy of internet-based integrated intervention on depression and anxiety symptoms in patients with COVID-19. *Journal of Zhejiang University. Science. B*; 2020.
309. CDC. Situation Summary. 2020.

310. Lu, Roujian, Zhao, Xiang, Li, Juan, Niu, Peihua, Yang, Bo, Wu, Honglong, Wang, Wenling, Song, Hao, Huang, Baoying, Zhu, Na. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *The Lancet*; 2020.
311. Lam, Tommy Tsan-Yuk, Jia, Na, Zhang, Ya-Wei, Shum, Marcus Ho-Hin, Jiang, Jia-Fu, Zhu, Hua-Chen, Tong, Yi-Gang, Shi, Yong-Xia, Ni, Xue-Bing, Liao, Yun-Shi, Li, Wen-Juan, Jiang, Bao-Gui, Wei, Wei, Yuan, Ting-Ting, Zheng, Kui, Cui, Xiao-Ming, Li, Jie, Pei, Guang-Qian, Qiang, Xin, Cheung, William Yiu-Man, Li, Lian-Feng, Sun, Fang-Fang, Qin, Si, Huang, Ji-Cheng, Leung, Gabriel M., Holmes, Edward C., Hu, Yan-Ling, Guan, Yi, Cao, Wu-Chun. Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins. *Nature*; 2020/07/01.
312. Ma, Virginia L., Nair, Meera S. Coronavirus May Have Spread in China Last August, Preliminary Harvard Study Suggests. 2020.
313. Kuo, Lily, Boseley, Sarah. Coronavirus may have been in Wuhan in August, study suggests. 2020.
314. Faulconbridge, Guy, Tian, Yew Lun, Kelland, Kate, Steenhuysen, Julie. China, scientists dismiss Harvard study suggesting COVID-19 was spreading in Wuhan in August. 2020.
315. Everington, Keoni. US investigating 'hazardous event' in Wuhan lab in October. 2020.
316. Dilanian, Ken, Arrow, Ruairidh, Kube, Courtney, Lee, Carol E., Jones, Louise, Bodo, Lorand. Report says cellphone data suggests October shutdown at Wuhan lab, but experts are skeptical. 2020.
317. Cohen, J. Call of the wild why many scientists say it's unlikely that SARS-CoV2 originated from a "lab leak". *Science*; September 2, 2021.
318. Calisher, C., Carroll, D., Colwell, R., Corley, R. B., Daszak, P., Drosten, C., Enjuanes, L., Farrar, J., Field, H., Golding, J., Gorbalenya, A., Haagmans, B., Hughes, J., Karesh, W., Keusch, G., Lam, S., Lubroth, J., Mackenzie, J., Madoff, L., Mazet, J., Palese, P., Perlman, S., Poon, L., Roizman, B., Saif, L., Subbarao, K., Turner, M. Statement in support of the scientists, public health professionals, and medical professionals of China combatting COVID-19. *Lancet*; Mar 7 2020.
319. Quay, Q., Muller, R. The science suggests a wuhan lab leak. *WSJ*; 2021.
320. van Helden, J., Butler, C., Achaz, G., Canard, B., Casane, D., Claverie, J., Colombo, F., Courtier, V., Ebright, R., Graner, F., Leitenberg, M., Morand, S., Petrovsky, N., Segreto, R., Decroly, E., Halloy, J. An appeal for an objective, open, and transparent scientific debate about the origin of SARS-CoV-2. *Lancet*; Oct 16 2021.
321. Ghosal, Aniruddha, Neergaard, Lauran. Health officials worry as untraceable virus clusters emerge. 2020.
322. Johansson, M., Quandelacy, T., Kada, S., Prasad, P., Steele, M., Brooks, J., Slayton, R., Biggerstaff, M., Butler, J. SARS-CoV-2 transmission from people without COVID-19 symptoms. *JAMA network open*; 2021.
323. CDC. COVID-19 Pandemic Planning Scenarios. 2020.
324. Bai, Yan, Yao, Lingsheng, Wei, Tao, Tian, Fei, Jin, Dong-Yan, Chen, Lijuan, Wang, Meiyun. Presumed asymptomatic carrier transmission of COVID-19. *Jama*; 2020.
325. Dowdle, W. The principles of disease elimination and eradication. December 31, 1999.
326. Corona, A. Disease eradication: what does it take to wipe out a disease?. March 6, 2020.
327. Kuchipudi, S., Surendran-Nair, M., Ruden, R., Yon, M., Nissly, R., Vandegrift, K., Nelli, R., Li, L., Jayarao, B., Maranas, C., Levine, N., Willgert, K., Conlan, A., Olsen, R., Davis, J., Musser, J., Hudson, P., Kapur, V. Multiple spillovers from humans and onward transmission of SARS-CoV-2 in white-tailed deer. *Proc Natl Acad Sci U S A*; Feb 8 2022.
328. Marques, A., Sherrill-Mix, S., Everett, J., Adhikari, H., Reddy, S., Ellis, J., Zelif, H., Greening, S., Cannuscio, C., Strelau, K. Evolutionary trajectories of SARS-CoV-2 alpha and delta variants in white-tailed deer in Pennsylvania. *medRxiv*; 2022.
329. Murphy, H., Ly, H. Understanding the prevalence of SARS-CoV-2 (COVID-19) exposure in companion, captive, wild, and farmed animals. *Virulence*; Dec 2021.
330. Pahuja, Sourabh, Madan, Manu, Mittal, Saurabh, Pandey, Ravindra Mohan, Nilima, Nilima, Madan, Karan, Mohan, Anant, Hadda, Vijay, Tiwari, Pawan, Guleria, Randeep. Weather Parameters and COVID-19: A Correlational Analysis. *Journal of Occupational and Environmental Medicine*; 2020.
331. Adnan, Shahzada, Hanif, Muhammad, Khan, Azmat Hayat, Latif, Muhammad, Ullah, Kalim, Bashir, Furrukh, Kamil, Shahzad, Haider, Sajjad. Impact of Heat Index and Ultraviolet Index on COVID-19 in Major Cities of Pakistan. *Journal of Occupational and Environmental Medicine*; 2021.
332. Ahlawat, Ajit, Wiedensohler, Alfred, Mishra, Sumit Kumar. An Overview on the role of relative humidity in airborne transmission of SARS-CoV-2 in indoor environments. *Aerosol Air Qual. Res*; 2020.
333. Vally, Hassan. 6 countries, 6 curves: how nations that moved fast against COVID-19 avoided disaster. 2020.

334. Wikipedia. COVID-19 pandemic lockdowns. 2020.
335. Kaplan, Juliana, Frias, Lauren, McFall-Johnsen, Morgan. Countries around the world are reopening — here's our constantly updated list of how they're doing it and who remains under lockdown. 2020.
336. Worldometer. Sweden. 2020.
337. Worldometer. Denmark. 2020.
338. Worldometer. Japan. 2020.
339. Statista. Number of tested and confirmed coronavirus (COVID-19) cases in Denmark since January 2020. 2020.
340. Paton, Callum. Coronavirus: Sweden's neighbours wary of reopening borders after surge in deaths. 2020.
341. Milne, Richard. First to close - first to reopen: Denmark's gain from virus response. 2020.
342. Centeno, Chris. Did Our Coronavirus Shutdown Work? You Decide. 2020.
343. Mortensen, Andreas, Skydsgaard, Nikolaj. Reopening schools in Denmark did not worsen outbreak, data shows. 2020.
344. WHO. Tracking SARS-CoV-2 variants. February 22, 2022.
345. Grannis, S. J., Rowley, E. A., Ong, T. C., Stenehjem, E., Klein, N. P., DeSilva, M. B., Naleway, A. L., Natarajan, K., Thompson, M. G. Interim estimates of COVID-19 vaccine effectiveness against COVID-19-associated emergency department or urgent care clinic encounters and hospitalizations among adults during SARS-CoV-2 B.1.617.2 (delta) variant predominance - nine states, June-August 2021. *MMWR*; Sep 17 2021.
346. CDC. The possibility of COVID-19 after vaccination: breakthrough infections. December 17, 2021.
347. Scobie, H. M., Johnson, A. G., Suthar, A. B., Severson, R., Alden, N. B., Balter, S., Bertolino, D., Blythe, D., Brady, S., Cadwell, B., Cheng, I., Davidson, S., Delgadillo, J., Devinney, K., Duchin, J., Duwell, M., Fisher, R., Fleischauer, A., Grant, A., Griffin, J., Haddix, M., Hand, J., Hanson, M., Hawkins, E., Herlihy, R. K., Hicks, L., Holtzman, C., Hoskins, M., Hyun, J., Kaur, R., Kay, M., Kidrowski, H., Kim, C., Komatsu, K., Kugeler, K., Lewis, M., Lyons, B. C., Lyons, S., Lynfield, R., McCaffrey, K., McMullen, C., Milroy, L., Meyer, S., Nolen, L., Patel, M. R., Pogojans, S., Reese, H. E., Saupe, A., Sell, J., Sokol, T., Sosin, D., Stanislawski, E., Stevens, K., Vest, H., White, K., Wilson, E., MacNeil, A., Ritchey, M. D., Silk, B. J. Monitoring incidence of COVID-19 cases, hospitalizations, and deaths, by vaccination status - 13 U.S. jurisdictions, April 4-July 17, 2021. *MMWR*; Sep 17 2021.
348. Bajema, K. L., Dahl, R. M., Prill, M. M., Meites, E., Rodriguez-Barradas, M. C., Marconi, V. C., Beenhouwer, D. O., Brown, S. T., Holodniy, M., Lucero-Obusan, C., Rivera-Dominguez, G., Morones, R. G., Whitmire, A., Goldin, E. B., Evener, S. L., Tremarelli, M., Tong, S., Hall, A. J., Schrag, S. J., McMorro, M., Kobayashi, M., Verani, J. R., Surie, D. Effectiveness of COVID-19 mRNA vaccines against COVID-19-associated hospitalization - five veterans affairs medical centers, United States, February 1-August 6, 2021. *MMWR*; Sep 17 2021.
349. Britton, A., Fleming-Dutra, K., Shang, N., Smith, Z., Dorji, T., Derado, G., Accorsi, E., Ajani, U., Miller, J., Schrag, S., Verani, J. Association of COVID-19 vaccination with symptomatic SARS-CoV-2 infection by time since vaccination and delta variant predominance. *JAMA*; Mar 15 2022.
350. El Sahly, H., Baden, L., Essink, B., Doblecki-Lewis, S., Martin, J., Anderson, E., Campbell, T., Clark, J., Jackson, L., Fichtenbaum, C., Zervos, M., Rankin, B., Eder, F., Feldman, G., Kennelly, C., Han-Conrad, L., Levin, M., Neuzil, K., Corey, L., Gilbert, P., Janes, H., Follmann, D., Marovich, M., Polakowski, L., Mascola, J., Ledgerwood, J., Graham, B., August, A., Clouting, H., Deng, W., Han, S., Leav, B., Manzo, D., Pajon, R., Schödel, F., Tomassini, J., Zhou, H., Miller, J. Efficacy of the mRNA-1273 SARS-CoV-2 vaccine at completion of blinded phase. *NEJM*; Nov 4 2021.
351. Moline, H., Whitaker, M., Deng, L., Rhodes, J., Milucky, J., Pham, H., Patel, K., Anglin, O., Reingold, A., Chai, S., Alden, N., Kawasaki, B., Meek, J., Yousey-Hindes, K., Anderson, E., Farley, M., Ryan, P., Kim, S., Nunez, V., Como-Sabetti, K., Lynfield, R., Sosin, D. M., McMullen, C., Muse, A., Barney, G., Bennett, N., Bushey, S., Shiltz, J., Sutton, M., Abdullah, N., Talbot, H., Schaffner, W., Chatelain, R., Ortega, J., Murthy, B., Zell, E., Schrag, S., Taylor, C., Shang, N., Verani, J., Havers, F. Effectiveness of COVID-19 vaccines in preventing hospitalization among adults aged ≥65 Years - COVID-NET, 13 States, February-April 2021. *MMWR*; Aug 13 2021.
352. Lutrick, K., Rivers, P., Yoo, Y., Grant, L., Hollister, J., Jovel, K., Khan, S., Lowe, A., Baccam, Z., Hanson, H., Olsho, L., Fowlkes, A., Caban-Martinez, A., Porter, C., Yoon, S., Meece, J., Gaglani, M., Burns, J., Mayo-Lamberte, J., Nakayima Miiro, F., Bissonnette, A., LeClair, L., Kutty, P., Romine, J., Stefanski, E., Edwards, L., Ellingson, K., Gerald, J., Bedrick, E., Madhivanan, P., Krupp, K., Gerald, L., Thompson, M., Burgess, J. Interim estimate of vaccine effectiveness of BNT162b2 (Pfizer-BioNTech) vaccine in preventing SARS-CoV-2 infection among adolescents aged 12-17 years - Arizona, July-December 2021. *MMWR*; Dec 31 2021.
353. Accorsi, E., Britton, A., Fleming-Dutra, K., Smith, Z., Shang, N., Derado, G., Miller, J., Schrag, S., Verani, J. Association between 3 doses of mRNA COVID-19 vaccine and symptomatic infection caused by the SARS-CoV-2 omicron and delta variants. *JAMA*; Feb 15 2022.

354. CDC. Science brief: omicron (B.2.2.529) variant. December 2, 2021.
355. Schreiber, M. The scientist in Botswana who identified omicron was saddened by the world's reaction. December 16, 2021.
356. Mallapaty, S. Where did Omicron come from? Three key theories. *Nature*; 2022.
357. Wei, C., Shan, K., Wang, W., Zhang, S., Huan, Q., Qian, W. Evidence for a mouse origin of the SARS-CoV-2 Omicron variant. *J Genet Genomics*; Dec 2021.
358. CDC. Monitoring variant proportions. February 26, 2022.
359. Emont, J. Covid-19 factory closures prompt some U.S. businesses to rethink vietnam. *Wall Street Journal*; 2021.
360. Lynch, D. Inside America's broken supply chain. *Washington Post*; September 20, 2021.
361. Kay, G. The workers who fuel the global supply chain warned of a possible 'systems collapse' if solutions aren't found. *Business Insider Africa*; September 30, 2021.
362. Kurtenbach, E. Shortages, shipping, shutdowns hit Asian factory output. September 30, 2021.
363. Lewnard, J., Hong, V., Patel, M., Kahn, R., Lipsitch, M., Tartof, S. Clinical outcomes among patients infected with Omicron (B. 1.1. 529) SARS-CoV-2 variant in southern California. *medRxiv*; 2022.
364. Riediker, M., Briceno-Ayala, L., Ichihara, G., Albani, D., Poffet, D., Tsai, D. H., Iff, S., Monn, C. Higher viral load and infectivity increase risk of aerosol transmission for Delta and Omicron variants of SARS-CoV-2. *Swiss Med Wkly*; Jan 3 2022.
365. Chen, J., Wang, R., Gilby, N., Wei, G. Omicron (B.1.1.529): infectivity, vaccine breakthrough, and antibody resistance. *ArXiv*; Dec 1 2021.
366. Nishiura, H., Ito, K., Anzai, A., Kobayashi, T., Piantham, C., Rodríguez-Morales, A. Relative reproduction number of SARS-CoV-2 omicron (B. 1.1. 529) compared with delta variant in South Africa. 2022.
367. Sharma, S. Handwashing an effective tool to prevent COVID-19, other diseases. October 15, 2020.
368. Patel, A., Jernigan, D. Initial public health response and interim clinical guidance for the 2019 novel coronavirus outbreak—United States, December 31, 2019–February 4, 2020. *MMWR*; 2020.
369. Review, OSHA. Preventing spread of COVID-19 – Don't forget to wash your hands!. *OSHA Review Incorporated*; February 11, 2020.
370. Meyerowitz, E., Richterman, A., Gandhi, R., Sax, P. Transmission of SARS-CoV-2: a review of viral, host, and environmental factors. *Ann Intern Med*; Jan 2021.
371. Wiersinga, W Joost, Rhodes, Andrew, Cheng, Allen C, Peacock, Sharon J, Prescott, Hallie C. Pathophysiology, transmission, diagnosis, and treatment of coronavirus disease 2019 (COVID-19): a review. *Jama*; 2020.
372. Council, National, Research. Rapid expert consultation on the possibility of bioaerosol spread of SARS-CoV-2 for the COVID-19 pandemic (April 1, 2020). 2020.
373. Wei, Wycliffe E, Li, Zongbin, Chiew, Calvin J, Yong, Sarah E, Toh, Matthias P, Lee, Vernon J. Presymptomatic Transmission of SARS-CoV-2—Singapore, January 23–March 16, 2020. *Morbidity and Mortality Weekly Report*; 2020.
374. Yu, I. T., Li, Y., Wong, T. W., Tam, W., Chan, A. T., Lee, J. H., Leung, D. Y., Ho, T. Evidence of airborne transmission of the severe acute respiratory syndrome virus. *N Engl J Med*; Apr 22 2004.
375. Ames, Michael. Why an Idaho ski destination has one of the highest COVID-19 infection rates in the nation. 2020.
376. Read, Richard. A choir decided to go ahead with rehearsal. Now dozens of members have COVID-19 and two are dead. 2020.
377. Guo, Zhen-Dong, Wang, Zhong-Yi, Zhang, Shou-Feng, Li, Xiao, Li, Lin, Li, Chao, Cui, Yan, Fu, Rui-Bin, Dong, Yun-Zhu, Chi, Xiang-Yang. Aerosol and surface distribution of severe acute respiratory syndrome coronavirus 2 in hospital wards, Wuhan, China, 2020. *Emerg Infect Dis*; 2020.
378. Liu, Yuan, Ning, Zhi, Chen, Yu, Guo, Ming, Liu, Yingle, Gali, Nirmal Kumar, Sun, Li, Duan, Yusen, Cai, Jing, Westerdahl, Dane. Aerodynamic analysis of SARS-CoV-2 in two Wuhan hospitals. *Nature*; 2020.
379. Santarpia, Joshua L, Rivera, Danielle N, Herrera, Vicki, Morwitzer, M Jane, Creager, Hannah, Santarpia, George W, Crown, Kevin K, Brett-Major, David, Schnaubelt, Elizabeth, Broadhurst, M Jana. Transmission potential of SARS-CoV-2 in viral shedding observed at the University of Nebraska Medical Center. *MedRxiv*; 2020.
380. Anderson, Elizabeth L., Turnham, Paul, Griffin, John R., Clarke, Chester C. Consideration of the Aerosol Transmission for COVID-19 and Public Health. *Risk analysis : an official publication of the Society for Risk Analysis*; 04/20/received 04/20/accepted 5/2/pubmed 5/15/medline 5/2/entrez 2020.

381. Tang, Song, Mao, Yixin, Jones, Rachael M., Tan, Qiyue, Ji, John S., Li, Na, Shen, Jin, Lv, Yuebin, Pan, Lijun, Ding, Pei, Wang, Xiaochen, Wang, Youbin, MacIntyre, C. Raina, Shi, Xiaoming. Aerosol transmission of SARS-CoV-2? Evidence, prevention and control. *Environment international*; 2020.
382. Jimenez, Jose-Luis. COVID-19 Data Dives: Why Arguments Against SARS-CoV-2 Aerosol Transmission Don't Hold Water. 2020.
383. Hamner, Lea, Dubbel, Polly, Capron, Ian, Ross, Andy, Jordan, Amber, Lee, Jaxon, Lynn, Joanne, Ball, Amelia, Narwal, Simranjit, Russell, Sam, Patrick, Dale, Leibrand, Howard. High SARS-CoV-2 Attack Rate Following Exposure at a Choir Practice — Skagit County, Washington, March 2020. 2020.
384. Miller, Shelly L., Nazaroff, William W., Jimenez, Jose L., Boerstra, Atze, Buonanno, Giorgio, Dancer, Stephanie J., Kurnitski, Jarek, Marr, Linsey C., Morawska, Lidia, Noakes, Catherine. Transmission of SARS-CoV-2 by inhalation of respiratory aerosol in the Skagit Valley Chorale superspreading event. *medRxiv*; 2020.
385. Bourouiba, Lydia. Turbulent gas clouds and respiratory pathogen emissions: potential implications for reducing transmission of COVID-19. *Jama*; 2020.
386. Lewis, D. Is the coronavirus airborne? Experts can't agree. *Nature*; Apr 2020.
387. Zhang, Renyi, Li, Yixin, Zhang, Annie L., Wang, Yuan, Molina, Mario J. Identifying airborne transmission as the dominant route for the spread of COVID-19. *Proceedings of the National Academy of Sciences*; 2020.
388. Morawska, L., Milton, D. K. It is Time to Address Airborne Transmission of COVID-19. *Clin Infect Dis*; Jul 6 2020.
389. Li, Yuguo, Qian, Hua, Hang, Jian, Chen, Xuguang, Hong, Ling, Liang, Peng, Li, Jiansen, Xiao, Shenglan, Wei, Jianjian, Liu, Li. Evidence for probable aerosol transmission of SARS-CoV-2 in a poorly ventilated restaurant. *medRxiv*; 2020.
390. van Doremalen, Neeltje, Bushmaker, Trenton, Morris, Dylan H, Holbrook, Myndi G, Gamble, Amandine, Williamson, Brandi N, Tamin, Azaibi, Harcourt, Jennifer L, Thornburg, Natalie J, Gerber, Susan I. Aerosol and surface stability of SARS-CoV-2 as compared with SARS-CoV-1. *New England Journal of Medicine*; 2020.
391. CDC. Scientific brief: SARS-CoV-2 transmission. May 7, 2021.
392. Wang, C.C., Prather, K.A., Sznitman, J., Jimenez, J.L., Lakdawala, S.S., Tufekci, Z., Marr, L.C. Airborne transmission of respiratory viruses. *Science*; 2021.
393. Shang, Y., Dong, J., Tian, L., He, F., Tu, J. An improved numerical model for epidemic transmission and infection risks assessment in indoor environment. *J Aerosol Sci*; May 2022.
394. Fears, AC, Klimstra, WB, Duprex, P, Hartman, A, Weaver, SC, Plante, KS, Mirchandani, D, Plante, JA, Aguilar, PV, Fernández, D. Persistence of severe acute respiratory syndrome coronavirus 2 in aerosol suspensions. *Emerg. Infect. Dis.*; 2020.
395. Verma, Siddhartha. Airborne Transmission & Face Masks: How Do Different Types of Masks Protect Against Various Ranges of Transmitted Droplets? How Far Can Aerosolized Droplets Travel in the Air From an Uncovered Cough & For How Long?. 2020.
396. Lewis, Dyani. Mounting evidence suggests coronavirus is airborne — but health advice has not caught up. 2020.
397. IES. IES Committee Report: Germicidal Ultraviolet (GUV) – Frequently Asked Questions. 2020.
398. Schuit, Michael, Ratnesar-Shumate, Shanna, Yolitz, Jason, Williams, Gregory, Weaver, Wade, Green, Brian, Miller, David, Krause, Melissa, Beck, Katie, Wood, Stewart, Holland, Brian, Bohannon, Jordan, Freeburger, Denise, Hooper, Idris, Biryukov, Jennifer, Altamura, Louis A., Wahl, Victoria, Hevey, Michael, Dabisch, Paul. Airborne SARS-CoV-2 Is Rapidly Inactivated by Simulated Sunlight. *The Journal of Infectious Diseases*; 2020.
399. Fears, Alyssa C, Klimstra, William B, Duprex, Paul, Hartman, Amy, Weaver, Scott C, Plante, Kenneth S, Mirchandani, Divya, Plante, Jessica Ann, Aguilar, Patricia V, Fernández, Diana. Persistence of severe acute respiratory syndrome coronavirus 2 in aerosol suspensions. *Emerging infectious diseases*; 2020.
400. Stadnytskyi, Valentyn, Bax, Christina E., Bax, Adriaan, Anfinrud, Philip. The airborne lifetime of small speech droplets and their potential importance in SARS-CoV-2 transmission. *Proceedings of the National Academy of Sciences*; 2020.
401. Karimzadeh, S., Bhopal, R., Nguyen Tien, H. Review of infective dose, routes of transmission and outcome of COVID-19 caused by the SARS-COV-2: comparison with other respiratory viruses. *Epidemiol Infect*; Apr 14 2021.
402. Flaxman, S, Mishra, S, Gandy, A. Estimating the number of infections and the impact of non-pharmaceutical interventions on COVID-19 in 11 European countries. *Imperial College preprint*; 2020.
403. WHO. Statement on the meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel coronavirus (2019-nCoV). 2020.

404. Adam, Dillon C., Wu, Peng, Wong, Jessica Y., Lau, Eric H. Y., Tsang, Tim K., Cauchemez, Simon, Leung, Gabriel M., Cowling, Benjamin J. Clustering and superspreading potential of SARS-CoV-2 infections in Hong Kong. *Nature Medicine*; 2020/11/01.
405. Majra, D., Benson, J., Pitts, J., Stebbing, J. SARS-CoV-2 (COVID-19) superspreader events. *J Infect*; Jan 2021.
406. Willebrand, K. S., Pischel, L., Malik, A. A., Jenness, S. M., Omer, S. B. A review of COVID-19 transmission dynamics and clinical outcomes on cruise ships worldwide, January to October 2020. *Euro Surveill*; Jan 2022.
407. Brandal, L., MacDonald, E., Veneti, L., Ravlo, T., Lange, H., Naseer, U., Feruglio, S., Bragstad, K., Hungnes, O., Ødeskaug, L. Outbreak caused by the SARS-CoV-2 Omicron variant in Norway, November to December 2021. *Eurosurveillance*; 2021.
408. Garret, N., Tapley, A., Andriesen, J., Seocharan, I., Fisher, L.H., Bunts, L., Espy, N., Wallis, C., Randhawa, A.K., Goga, A. High rate of asymptomatic carriage associated with variant strain Omicron. *MedRxiv*; 2021.
409. Rodriguez, A. 'Enormous spread of omicron' may bring 140M new COVID infections to US in the next two months, model predicts. *USA Today*; December 22, 2021.
410. Kaul, R. 70% Omicron cases are asymptomatic: govt. *Hindustan Times*; December 25, 2021.
411. Bajema, Kristina L., Wiegand, Ryan E., Cuffe, Kendra, Patel, Sadhna V., Iachan, Ronaldo, Lim, Travis, Lee, Adam, Moyse, Davia, Havers, Fiona P., Harding, Lee, Fry, Alicia M., Hall, Aron J., Martin, Kelly, Biel, Marjorie, Deng, Yangyang, Meyer, William A., III, Mathur, Mohit, Kyle, Tonja, Gundlapalli, Adi V., Thornburg, Natalie J., Petersen, Lyle R., Edens, Chris. Estimated SARS-CoV-2 Seroprevalence in the US as of September 2020. *JAMA Internal Medicine*; 2020.
412. CDC. Estimated Disease Burden of COVID-19. 2021.
413. Oran, Daniel P, Topol, Eric J. Prevalence of Asymptomatic SARS-CoV-2 Infection: A Narrative Review. *Annals of Internal Medicine*; 2020.
414. Nishiura, Hiroshi, Kobayashi, Tetsuro, Miyama, Takeshi, Suzuki, Ayako, Jung, Sung-mok, Hayashi, Katsuma, Kinoshita, Ryo, Yang, Yichi, Yuan, Baoyin, Akhmetzhanov, Andrei R., Linton, Natalie M. Estimation of the asymptomatic ratio of novel coronavirus infections (COVID-19). *International Journal of Infectious Diseases*; 2020.
415. Gudbjartsson, Daniel F., Helgason, Agnar, Jonsson, Hakon, Magnusson, Olafur T., Melsted, Pall, Norddahl, Gudmundur L., Saemundsdottir, Jona, Sigurdsson, Asgeir, Sulem, Patrick, Agustsdottir, Arna B., Eiriksdottir, Berglind, Fridriksdottir, Run, Gardarsdottir, Elisabet E., Georgsson, Gudmundur, Gretarsdottir, Olafia S., Gudmundsson, Kjartan R., Gunnarsdottir, Thora R., Gylfason, Arnaldur, Holm, Hilma, Jensson, Brynjar O., Jonasdottir, Aslaug, Jonsson, Frosti, Josefsdottir, Kamilla S., Kristjansson, Thordur, Magnusdottir, Droplaug N., le Roux, Louise, Sigmundsdottir, Gudrun, Sveinbjornsson, Gardar, Sveinsdottir, Kristin E., Sveinsdottir, Maney, Thorarensen, Emil A., Thorbjornsson, Bjarni, Löve, Arthur, Masson, Gisli, Jonsdottir, Ingileif, Möller, Alma D., Gudnason, Thorolfur, Kristinsson, Karl G., Thorsteinsdottir, Unnur, Stefansson, Kari. Spread of SARS-CoV-2 in the Icelandic Population. *New England Journal of Medicine*; 2020.
416. Lavezzo, Enrico, Franchin, Elisa, Ciavarella, Constanze, Cuomo-Dannenburg, Gina, Barzon, Luisa, Del Vecchio, Claudia, Rossi, Lucia, Manganelli, Riccardo, Loregian, Arianna, Navarin, Nicolò. Suppression of COVID-19 outbreak in the municipality of Vo, Italy. *medRxiv*; 2020.
417. Moriarty, Leah F. Public health responses to COVID-19 outbreaks on cruise ships—worldwide, February–March 2020. *MMWR. Morbidity and mortality weekly report*; 2020.
418. Baggett, Travis P., Keyes, Harrison, Sporn, Nora, Gaeta, Jessie M. COVID-19 outbreak at a large homeless shelter in Boston: Implications for universal testing. *medRxiv*; 2020.
419. Chou, Elizabeth. Dozens positive for coronavirus at LA's Skid Row homeless shelter, after all residents tested. 2020.
420. Sutton, Desmond, Fuchs, Karin, D'Alton, Mary, Goffman, Dena. Universal Screening for SARS-CoV-2 in Women Admitted for Delivery. *New England Journal of Medicine*; 2020.
421. Navy, U.S. U.S. Navy COVID-19 updates. Daily Update: April 24, 2020. 2020.
422. Figaro, Le. Coronavirus : bilan définitif de 1046 cas sur le Charles de Gaulle. 2020.
423. Lytras, Theodore, Dellis, George, Flountzi, Anastasia, Hatzianastasiou, Sophia, Nikolopoulou, Georgia, Tsekou, Katerina, Diamantis, Zafiris, Stathopoulou, Grigoria, Togka, Marianthi, Gerolymatos, Gerasimos, Rigakos, George, Sapounas, Spiridon, Tsiodras, Sotirios. High prevalence of SARS-CoV-2 infection in repatriation flights to Greece from three European countries. *Journal of Travel Medicine*; 2020.
424. Arons, Melissa M, Hatfield, Kelly M, Reddy, Sujan C, Kimball, Anne, James, Allison, Jacobs, Jesica R, Taylor, Joanne, Spicer, Kevin, Bardossy, Ana C, Oakley, Lisa P. Presymptomatic SARS-CoV-2 infections and transmission in a skilled nursing facility. *New England journal of medicine*; 2020.

425. So, Linda, Smith, Grant. In four US state prisons, nearly 3,300 inmates test positive for coronavirus—96% without symptoms. Reuters. 25 April 2020. 2020.
426. Barrett, E. S., Horton, D. B., Roy, J., Gennaro, M. L., Brooks, A., Tischfield, J., Greenberg, P., Andrews, T., Jagpal, S., Reilly, N., Blaser, M. J., Carson, J. L., Panettieri, R. A., Jr. Prevalence of SARS-CoV-2 infection in previously undiagnosed health care workers at the onset of the U.S. COVID-19 epidemic. *medRxiv*; Apr 24 2020.
427. Ratliff, Victoria. About 45 % of COVID-19-positive Hoosiers don't know it. 2020.
428. Ing, A. J., Cocks, C., Green, J. P. COVID-19: in the footsteps of Ernest Shackleton. *Thorax*; Aug 2020.
429. NBC San Diego. USS Roosevelt's Asymptomatic Cases Helping Scientists Understand Virus. 2020.
430. Peniston, Bradley. The Battle of USS Theodore Roosevelt: a Timeline. 2020.
431. Seligman, Lara. Sailors keep testing positive on aircraft carrier, despite 2-week isolation. 2020.
432. University, Indiana. IU, ISDH release preliminary findings about impact of COVID-19 in Indiana. 2020.
433. McFarling, Usha Lee. When hard data are 'heartbreaking': Testing blitz in San Francisco shows Covid-19 struck mostly low-wage workers. 2020.
434. Kampf, Günter, Todt, Daniel, Pfaender, Stephanie, Steinmann, Eike. Persistence of coronaviruses on inanimate surfaces and its inactivation with biocidal agents. *Journal of Hospital Infection*; 2020.
435. Hirose, Ryohei, Ikegaya, Hiroshi, Naito, Yuji, Watanabe, Naoto, Yoshida, Takuma, Bandou, Risa, Daidoji, Tomo, Itoh, Yoshito, Nakaya, Takaaki. Survival of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) and Influenza Virus on Human Skin: Importance of Hand Hygiene in Coronavirus Disease 2019 (COVID-19). *Clinical Infectious Diseases*; 2020.
436. Eslami, H., Jalili, M. The role of environmental factors to transmission of SARS-CoV-2 (COVID-19). *AMB Express*; May 15 2020.
437. Huang, Z., Huang, J., Gu, Q., Du, P., Liang, H., Dong, Q. Optimal temperature zone for the dispersal of COVID-19. *Sci Total Environ*; May 16 2020.
438. Beaubien, Jason. Will Heat And Humidity Kill The Coronavirus?. 2020.
439. Chan, K. H., Peiris, J. S. Malik, Lam, S. Y., Poon, L. L. M., Yuen, K. Y., Seto, W. H. The Effects of Temperature and Relative Humidity on the Viability of the SARS Coronavirus. *Advances in Virology*; 2011/10/01.
440. Ratnesar-Shumate, S., Williams, G., Green, B., Krause, M., Holland, B., Wood, S., Bohannon, J., Boydston, J., Freeburger, D., Hooper, I., Beck, K., Yeager, J., Altamura, L. A., Biryukov, J., Yolitz, J., Schuit, M., Wahl, V., Hevey, M., Dabisch, P. Simulated Sunlight Rapidly Inactivates SARS-CoV-2 on Surfaces. *J Infect Dis*; May 20 2020.
441. Liu, Z., Zhu, L., Wang, Y., Zhou, Z., Guo, Y. The correlation between COVID-19 activities and climate factors in different climate types areas. *JOEM*; 2021.
442. Amherst, University of Massachusetts. Median incubation period for COVID-19. 2020.
443. Lauer, Stephen A, Grantz, Kyra H, Bi, Qifang, Jones, Forrest K, Zheng, Qulu, Meredith, Hannah R, Azman, Andrew S, Reich, Nicholas G, Lessler, Justin. The incubation period of coronavirus disease 2019 (COVID-19) from publicly reported confirmed cases: estimation and application. *Annals of internal medicine*; 2020.
444. Jansen, L., Tegomoh, B., Lange, K., Showalter, K., Figliomeni, J., Abdalhamid, B., Iwen, P.C., Fauver, J., Buss, B., Donahue, M. Investigation of a Sars-cov-2 b. 1.1. 529 (omicron) variant cluster—nebraska, november–december 2021. *MMWR*; 2021.
445. CDC. Symptoms. 2020.
446. Guan, Wei-jie, Ni, Zheng-yi, Hu, Yu, Liang, Wen-hua, Ou, Chun-quan, He, Jian-xing, Liu, Lei, Shan, Hong, Lei, Chun-liang, Hui, David SC. Clinical characteristics of 2019 novel coronavirus infection in China. *MedRxiv*; 2020.
447. National Institute of Infectious Disease Control and Prevention Medicine and National Center for Global Health. Active epidemiological investigation on SARS-CoV-2 infection caused by Omicron variant (Pango lineage B.1.1.529) in Japan: preliminary report on infectious period. *NCID*; 2022.
448. Saul, D. Study of NBA omicron cases finds many still infectious after CDC recommended self-isolation period ends. *Forbes*; January 15, 2022.
449. Jones, T.C., Biele, G., Mühlemann, B., Veith, T., Schneider, J., Beheim-Schwarzbach, J., Bleicker, T., Tesch, J., Schmidt, M.L., Sander, L.E. Estimating infectiousness throughout SARS-CoV-2 infection course. *Science*; 2021.
450. Torjesen, I. Covid-19: peak of viral shedding is later with omicron variant, Japanese data suggest. 2022.

451. Wölfel, Roman, Corman, Victor M, Guggemos, Wolfgang, Seilmaier, Michael, Zange, Sabine, Müller, Marcel A, Niemeyer, Daniela, Jones, Terry C, Vollmar, Patrick, Rothe, Camilla. Virological assessment of hospitalized patients with COVID-2019. *Nature*; 2020.
452. Widders, Arabella, Broom, Alex, Broom, Jennifer. SARS-CoV-2: The viral shedding vs infectivity dilemma. *Infection, disease & health*; 2020.
453. Cevik, Muge, Tate, Matthew, Lloyd, Ollie, Maraolo, Alberto Enrico, Schafers, Jenna, Ho, Antonia. SARS-CoV-2, SARS-CoV, and MERS-CoV viral load dynamics, duration of viral shedding, and infectiousness: a systematic review and meta-analysis. *The Lancet Microbe*; 2020.
454. Benefield, Amy E, Skrip, Laura A, Clement, Andrea, Althouse, Rachel A, Chang, Stewart, Althouse, Benjamin Muir. SARS-CoV-2 viral load peaks prior to symptom onset: a systematic review and individual-pooled analysis of coronavirus viral load from 66 studies. *medRxiv*; 2020.
455. Disease Control and Prevention Agency, Korea. Findings from investigation and analysis of re-positive cases. 2020.
456. Li, L., Liang, Y., Hu, F., Yan, H., Li, Y., Xie, Z., Huang, L., Zhao, J., Wan, Z., Wang, H., Shui, J., Cai, W., Tang, S. Molecular and serological characterization of SARS-CoV-2 infection among COVID-19 patients. *Virology*; Dec 2020.
457. Xiao, Ai Tang, Tong, Yi Xin, Gao, Chun, Zhu, Li, Zhang, Yu Jie, Zhang, Sheng. Dynamic profile of RT-PCR findings from 301 COVID-19 patients in Wuhan, China: A descriptive study. *Journal of clinical virology : the official publication of the Pan American Society for Clinical Virology*; 2020.
458. Gousseff, M., Penot, P., Gallay, L., Batisse, D., Benech, N., Bouiller, K., Collarino, R., Conrad, A., Slama, D., Joseph, C., Lemaignen, A., Lescure, F. X., Levy, B., Mahevas, M., Pozzetto, B., Vignier, N., Wyplosz, B., Salmon, D., Goehring, F., Botelho-Nevers, E. Clinical recurrences of COVID-19 symptoms after recovery: Viral relapse, reinfection or inflammatory rebound?. *J Infect*; Nov 2020.
459. To, Kelvin Kai-Wang, Hung, Ivan Fan-Ngai, Chan, Kwok-Hung, Yuan, Shuofeng, To, Wing-Kin, Tsang, Dominic Ngai-Chong, Cheng, Vincent Chi-Chung, Chen, Zhiwei, Kok, Kin-Hang, Yuen, Kwok-Yung. Serum antibody profile of a patient with COVID-19 reinfection. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*; 2020.
460. Bonifácio, Livia Pimenta, Pereira, Ana Paula Sulino, Araújo, Daniel Cardoso de Almeida E., Balbão, Viviane da Mata Pasti, Fonseca, Benedito Antônio Lopes da, Passos, Afonso Dinis Costa, Bellissimo-Rodrigues, Fernando. Are SARS-CoV-2 reinfection and Covid-19 recurrence possible? a case report from Brazil. *Revista da Sociedade Brasileira de Medicina Tropical*; 2020.
461. Parry, Jane. Covid-19: Hong Kong scientists report first confirmed case of reinfection. *BMJ*; 2020.
462. To, K. K., Hung, I. F., Ip, J. D., Chu, A. W., Chan, W. M., Tam, A. R., Fong, C. H., Yuan, S., Tsoi, H. W., Ng, A. C., Lee, L. L., Wan, P., Tso, E., To, W. K., Tsang, D., Chan, K. H., Huang, J. D., Kok, K. H., Cheng, V. C., Yuen, K. Y. COVID-19 re-infection by a phylogenetically distinct SARS-coronavirus-2 strain confirmed by whole genome sequencing. *Clin Infect Dis*; Aug 25 2020.
463. Hong, Jinshan, Gale, Jason. Hong Kong Reports First Confirmed Coronavirus Re-Infection. 2020.
464. Ferguson, N., Ghani, A., Cori, A., Hogan, A., Hinsley, W., Volz, E. Report 49: growth, population distribution and immune escape of Omicron in England. *Imperial College London*; 2021.
465. Guzik, T. J., Mohiddin, S. A., Dimarco, A., Patel, V., Savvatis, K., Marelli-Berg, F. M., Madhur, M. S., Tomaszewski, M., Maffia, P., D'Acquisto, F., Nicklin, S. A., Marian, A. J., Nosalski, R., Murray, E. C., Guzik, B., Berry, C., Touyz, R. M., Kreutz, R., Wang, D. W., Bhella, D., Sagliocco, O., Crea, F., Thomson, E. C., McInnes, I. B. COVID-19 and the cardiovascular system: implications for risk assessment, diagnosis, and treatment options. *Cardiovasc Res*; Apr 30 2020.
466. Siddamreddy, S., Thotakura, R., Dandu, V., Kanuru, S., Meegada, S. Corona Virus Disease 2019 (COVID-19) Presenting as Acute ST Elevation Myocardial Infarction. *Cureus*; Apr 22 2020.
467. Hu, Y., Sun, J., Dai, Z., Deng, H., Li, X., Huang, Q., Wu, Y., Sun, L., Xu, Y. Prevalence and severity of corona virus disease 2019 (COVID-19): A systematic review and meta-analysis. *J Clin Virol*; Apr 14 2020.
468. Zheng, Y., Xu, H., Yang, M., Zeng, Y., Chen, H., Liu, R., Li, Q., Zhang, N., Wang, D. Epidemiological characteristics and clinical features of 32 critical and 67 noncritical cases of COVID-19 in Chengdu. *J Clin Virol*; Apr 10 2020.
469. Liu, P. P., Blet, A., Smyth, D., Li, H. The Science Underlying COVID-19: Implications for the Cardiovascular System. *Circulation*; Apr 15 2020.
470. Fried, Justin A, Ramasubbu, Kumudha, Bhatt, Reema, Topkara, Veli K, Clerkin, Kevin J, Horn, Evelyn, Rabbani, LeRoy, Brodie, Daniel, Jain, Sneha S, Kirtane, Ajay. The variety of cardiovascular presentations of COVID-19. *Circulation*; 2020.

471. Rajpal, Saurabh, Tong, Matthew S, Borchers, James, Zareba, Karolina M, Obarski, Timothy P, Simonetti, Orlando P, Daniels, Curt J. Cardiovascular magnetic resonance findings in competitive athletes recovering from COVID-19 infection. *JAMA cardiology*; 2020.
472. Iacobucci, G. Covid-19: runny nose, headache, and fatigue are commonest symptoms of omicron, early data show. 2021.
473. Van Loon, Nele, Verbrugghe, Mathieu, Cartuyvels, Reinoud, Ramaekers, Dirk. Diagnosis of COVID-19 Based on Symptomatic Analysis of Hospital Healthcare Workers in Belgium: Observational Study in a Large Belgian Tertiary Care Center during Early COVID-19 Outbreak. *Journal of Occupational and Environmental Medicine*; 2020.
474. WHO. Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19). 2020.
475. Zhu, J., Ji, P., Pang, J., Zhong, Z., Li, H., He, C., Zhang, J., Zhao, C. Clinical characteristics of 3,062 COVID-19 patients: a meta-analysis. *J Med Virol*; Apr 15 2020.
476. Lovato, A., de Filippis, C. Clinical Presentation of COVID-19: A Systematic Review Focusing on Upper Airway Symptoms. *Ear Nose Throat J*; Apr 13 2020.
477. Michelen, Melina, Jones, Nicholas, Stavropoulou, Charitini. In patients of COVID-19, what are the symptoms and clinical features of mild and moderate cases?. 2020.
478. Pan, Lei, Mu, Mi, Ren, Hong Gang, Yang, P, Sun, Y, Wang, R. Clinical characteristics of COVID-19 patients with digestive symptoms in Hubei, China: a descriptive, cross-sectional, multicenter study. *Am J Gastroenterol*; 2020.
479. Wu, Ping, Duan, Fang, Luo, Chunhua, Liu, Qiang, Qu, Xingguang, Liang, Liang, Wu, Kaili. Characteristics of Ocular Findings of Patients With Coronavirus Disease 2019 (COVID-19) in Hubei Province, China. *JAMA ophthalmology*; 2020.
480. Moein, S. T., Hashemian, S. M. R., Mansourafshar, B., Khorram-Tousi, A., Tabarsi, P., Doty, R. L. Smell dysfunction: a biomarker for COVID-19. *Int Forum Allergy Rhinol*; Apr 17 2020.
481. Gandhi, Monica, Beyrer, Chris, Goosby, Eric. Masks Do More Than Protect Others During COVID-19: Reducing the Inoculum of SARS-CoV-2 to Protect the Wearer. *Journal of General Internal Medicine*; 2020/10/01.
482. Gandhi, Monica, Rutherford, George W. Facial Masking for Covid-19 — Potential for “Variolation” as We Await a Vaccine. *New England Journal of Medicine*; 2020.
483. Nicolai, Leo, Leunig, Alexander, Brambs, Sophia, Kaiser, Rainer, Weinberger, Tobias, Weigand, Michael, Muenchhoff, Maximilian, Hellmuth, Johannes C, Ledderose, Stephan, Schulz, Heiko. Immunothrombotic dysregulation in COVID-19 pneumonia is associated with respiratory failure and coagulopathy. *Circulation*; 2020.
484. Wichmann, Dominic, Sperhake, Jan-Peter, Lütgehetmann, Marc, Steurer, Stefan, Edler, Carolin, Heinemann, Axel, Heinrich, Fabian, Mushumba, Herbert, Kniep, Inga, Schröder, Ann Sophie, Burdelski, Christoph, de Heer, Geraldine, Nierhaus, Axel, Frings, Daniel, Pfefferle, Susanne, Becker, Heinrich, Bredereke-Wiedling, Hanns, de Weerth, Andreas, Paschen, Hans-Richard, Sheikhzadeh-Eggers, Sara, Stang, Axel, Schmiedel, Stefan, Bokemeyer, Carsten, Addo, Marylyn M., Aepfelbacher, Martin, Püschel, Klaus, Kluge, Stefan. Autopsy Findings and Venous Thromboembolism in Patients With COVID-19. *Ann Intern Med*; 2020.
485. Zhang, Y., Xiao, M., Zhang, S., Xia, P., Cao, W., Jiang, W., Chen, H., Ding, X., Zhao, H., Zhang, H., Wang, C., Zhao, J., Sun, X., Tian, R., Wu, W., Wu, D., Ma, J., Chen, Y., Zhang, D., Xie, J., Yan, X., Zhou, X., Liu, Z., Wang, J., Du, B., Qin, Y., Gao, P., Qin, X., Xu, Y., Zhang, W., Li, T., Zhang, F., Zhao, Y., Li, Y., Zhang, S. Coagulopathy and Antiphospholipid Antibodies in Patients with Covid-19. *N Engl J Med*; Apr 23 2020.
486. Oxley, T. J., Mocco, J., Majidi, S., Kellner, C. P., Shoirah, H., Singh, I. P., De Leacy, R. A., Shigematsu, T., Ladner, T. R., Yaeger, K. A., Skliut, M., Weinberger, J., Dangayach, N. S., Bederson, J. B., Tuhim, S., Fifi, J. T. Large-Vessel Stroke as a Presenting Feature of Covid-19 in the Young. *N Engl J Med*; Apr 28 2020.
487. Avula, Akshay, Nalleballe, Krishna, Narula, Naureen, Sapozhnikov, Steven, Dandu, Vasuki, Toom, Sudhamshi, Glaser, Allison, Elsayegh, Dany. COVID-19 presenting as stroke. *Brain, Behavior, and Immunity*; 2020/07/01/.
488. Lax, S., Skok, K., Zechner, P., Kessler, H., Kaufmann, N., Koelblinger, C., Vander, Kl., Bargfrieder, U., Trauner, M. Pulmonary arterial thrombosis in COVID-19 with fatal outcome: results from a prospective, single-center, clinicopathologic case series. *Ann Int Med*; 2020.
489. Klok, F. A., Kruip, Mjha, van der Meer, N. J. M., Arbous, M. S., Gommers, Dampj, Kant, K. M., Kaptein, F. H. J., van Paassen, J., Stals, M. A. M., Huisman, M. V., Endeman, H. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res*; Jul 2020.
490. Middeldorp, S., Coppens, M., van Haaps, T. F., Foppen, M., Vlaar, A. P., Müller, M. C. A., Bouman, C. C. S., Beenen, L. F. M., Kootte, R. S., Heijmans, J., Smits, L. P., Bonta, P. I., van Es, N. Incidence of venous thromboembolism in hospitalized patients with COVID-19. *J Thromb Haemost*; May 5 2020.

491. Levi, M., Thachil, J., Iba, T., Levy, J. H. Coagulation abnormalities and thrombosis in patients with COVID-19. *Lancet Haematol*; Jun 2020.
492. Root-Bernstein, R. COVID-19 coagulopathies: human blood proteins mimic SARS-CoV-2 virus, vaccine proteins and bacterial co-infections inducing autoimmunity: combinations of bacteria and SARS-CoV-2 synergize to induce autoantibodies targeting cardiolipin, cardiolipin-binding proteins, platelet factor 4, prothrombin, and coagulation factors. *Bioessays*; Dec 2021.
493. Ehsani, A. H., Nasimi, M., Bigdelo, Z. Pityriasis rosea as a cutaneous manifestation of COVID-19 infection. *J Eur Acad Dermatol Venereol*; May 2 2020.
494. Bouaziz, J. D., Duong, T., Jachiet, M., Velter, C., Lestang, P., Cassius, C., Arsouze, A., Domergue Than Trong, E., Bagot, M., Begon, E., Sulimovic, L., Rybojad, M. Vascular skin symptoms in COVID-19: a french observational study. *J Eur Acad Dermatol Venereol*; Apr 27 2020.
495. van Damme, C., Berlingin, E., Saussez, S., Accaputo, O. Acute urticaria with pyrexia as the first manifestations of a COVID-19 infection. *J Eur Acad Dermatol Venereol*; Apr 24 2020.
496. Henry, D., Ackerman, M., Sancelme, E., Finon, A., Esteve, E. Urticarial eruption in COVID-19 infection. *J Eur Acad Dermatol Venereol*; Apr 15 2020.
497. Zhao, Qing, Fang, Xiaokai, Pang, Zheng, Zhang, Bowen, Liu, Hong, Zhang, Furen. COVID-19 and cutaneous manifestations: a systematic review. *Journal of the European Academy of Dermatology and Venereology*; 2020.
498. Al Saiegh, F., Ghosh, R., Leibold, A., Avery, M. B., Schmidt, R. F., Theofanis, T., Mouchtouris, N., Philipp, L., Peiper, S. C., Wang, Z. X., Rincon, F., Tjoumakaris, S. I., Jabbour, P., Rosenwasser, R. H., Gooch, M. R. Status of SARS-CoV-2 in cerebrospinal fluid of patients with COVID-19 and stroke. *J Neurol Neurosurg Psychiatry*; Apr 30 2020.
499. Baig, A. M. Updates on What ACS Reported: Emerging Evidences of COVID-19 with Nervous System Involvement. *ACS Chem Neurosci*; Apr 28 2020.
500. Chacón-Aguilar, R., Osorio-Cámara, J. M., Sanjurjo-Jimenez, I., González-González, C., López-Carnero, J., Pérez-Moneo-Agapito, B. COVID-19: Fever syndrome and neurological symptoms in a neonate. *An Pediatr (Engl Ed)*; Apr 27 2020.
501. Gutiérrez-Ortiz, C., Méndez, A., Rodrigo-Rey, S., San Pedro-Murillo, E., Bermejo-Guerrero, L., Gordo-Mañas, R., de Aragón-Gómez, F., Benito-León, J. Miller Fisher Syndrome and polyneuritis cranialis in COVID-19. *Neurology*; Apr 17 2020.
502. Mao, L., Jin, H., Wang, M., Hu, Y., Chen, S., He, Q., Chang, J., Hong, C., Zhou, Y., Wang, D., Miao, X., Li, Y., Hu, B. Neurologic Manifestations of Hospitalized Patients With Coronavirus Disease 2019 in Wuhan, China. *JAMA Neurol*; Apr 10 2020.
503. Poyiadji, N., Shahin, G., Noujaim, D., Stone, M., Patel, S., Griffith, B. COVID-19-associated Acute Hemorrhagic Necrotizing Encephalopathy: CT and MRI Features. *Radiology*; Mar 31 2020.
504. Yan, Y., Yang, Y., Wang, F., Ren, H., Zhang, S., Shi, X., Yu, X., Dong, K. Clinical characteristics and outcomes of patients with severe covid-19 with diabetes. *BMJ Open Diabetes Res Care*; Apr 2020.
505. Varatharaj, A., Thomas, N., Ellul, M. A., Davies, N. W. S., Pollak, T. A., Tenorio, E. L., Sultan, M., Easton, A., Breen, G., Zandi, M., Coles, J. P., Manji, H., Al-Shahi Salman, R., Menon, D. K., Nicholson, T. R., Benjamin, L. A., Carson, A., Smith, C., Turner, M. R., Solomon, T., Kneen, R., Pett, S. L., Galea, I., Thomas, R. H., Michael, B. D. Neurological and neuropsychiatric complications of COVID-19 in 153 patients: a UK-wide surveillance study. *Lancet Psychiatry*; Jun 25 2020.
506. Rogers, J.P., David, A.S. A longer look at COVID-19 and neuropsychiatric outcomes. *Lancet Psychiatry*; 2021.
507. Ronco, C., Reis, T. Kidney involvement in COVID-19 and rationale for extracorporeal therapies. *Nat Rev Nephrol*; Apr 9 2020.
508. Velez, Juan Carlos Q., Caza, Tiffany, Larsen, Christopher P. COVAN is the new HIVAN: the re-emergence of collapsing glomerulopathy with COVID-19. *Nature Reviews Nephrology*; 2020/10/01.
509. Durvasula, R., Wellington, T., McNamara, E., Watnick, S. COVID-19 and Kidney Failure in the Acute Care Setting: Our Experience From Seattle. *Am J Kidney Dis*; Apr 8 2020.
510. Tang, N., Li, D., Wang, X., Sun, Z. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost*; Apr 2020.
511. Wong, John EL, Leo, Yee Sin, Tan, Chorch Chuan. COVID-19 in Singapore—current experience: critical global issues that require attention and action. *Jama*; 2020.
512. Bernheim, Adam, Mei, Xueyan, Huang, Mingqian, Yang, Yang, Fayad, Zahi A, Zhang, Ning, Diao, Kaiyue, Lin, Bin, Zhu, Xiqi, Li, Kunwei. Chest CT findings in coronavirus disease-19 (COVID-19): relationship to duration of infection. *Radiology*; 2020.

513. He, X., Lau, E. H. Y., Wu, P., Deng, X., Wang, J., Hao, X., Lau, Y. C., Wong, J. Y., Guan, Y., Tan, X., Mo, X., Chen, Y., Liao, B., Chen, W., Hu, F., Zhang, Q., Zhong, M., Wu, Y., Zhao, L., Zhang, F., Cowling, B. J., Li, F., Leung, G. M. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med*; Apr 15 2020.
514. Matthews, Alex, McDermott, Nick. IT'S SPREADING First coronavirus case confirmed in London as woman diagnosed with deadly bug brings total in UK to nine. 2020.
515. Chua, G. T., Wong, J. S. C., To, K. K. W., Lam, I. C. S., Yau, F. Y. S., Chan, W. H., Ho, P. P. K., Duque, J. S. R., Yip, C. C. Y., Ng, A. C. K., Wong, W. H. S., Kwong, J. H. Y., Leung, K. F. S., Wan, P. T., Lam, K., Wong, I. C. K., Kwok, J., Ho, M. H. K., Chan, G. C. F., Lau, Y. L., Ip, P., Kwan, M. Y. W. Saliva viral load better correlates with clinical and immunological profiles in children with coronavirus disease 2019. *Emerg Microbes Infect*; Dec 2021.
516. Baggio, S., L'Huillier, A. G., Yerly, S., Bellon, M., Wagner, N., Rohr, M., Huttner, A., Blanchard-Rohner, G., Loevy, N., Kaiser, L., Jacquerioz, F., Eckerle, I. SARS-CoV-2 viral load in the upper respiratory tract of children and adults with early acute COVID-19. *Clin Infect Dis*; Aug 6 2020.
517. Czeisler, Mark É, Lane, Rashon I, Petrosky, Emiko, Wiley, Joshua F, Christensen, Aleta, Njai, Rashid, Weaver, Matthew D, Robbins, Rebecca, Facer-Childs, Elise R, Barger, Laura K. Mental health, substance use, and suicidal ideation during the COVID-19 pandemic—United States, June 24–30, 2020. *Morbidity and Mortality Weekly Report*; 2020.
518. Goldstein, Joseph, Belluck, Pam. Children Are Falling Ill With a Baffling Ailment Related to Covid-19. 2020.
519. CDC. COVID-19 Science Update released: November 24, 2020. 2020.
520. Maslo, C., Friedland, R., Toubkin, M., Laubscher, A., Akaloo, T., Kama, B. Characteristics and outcomes of hospitalized patients in South Africa during the COVID-19 Omicron wave compared with previous waves. *JAMA*; Feb 8 2022.
521. Kobayashi, T., Jung, S. M., Linton, N. M., Kinoshita, R., Hayashi, K., Miyama, T., Anzai, A., Yang, Y., Yuan, B., Akhmetzhanov, A. R., Suzuki, A., Nishiura, H. Communicating the Risk of Death from Novel Coronavirus Disease (COVID-19). *J Clin Med*; Feb 21 2020.
522. Ledberg, Anders. Mortality of the COVID-19 outbreak in Sweden in relation to previous severe disease outbreaks. *medRxiv*; 2020.
523. Payne, Daniel. Antibody tests continue to suggest COVID-19 far more widespread, less deadly than initially thought. 2020.
524. Hamilton, Jon. Antibody Tests Point To Lower Death Rate For The Coronavirus Than First Thought. 2020.
525. Ioannidis, John. The infection fatality rate of COVID-19 inferred from seroprevalence data. *medRxiv*; 2020.
526. Garibaldi, Brian T, Fiksel, Jacob, Muschelli, John, Robinson, Matthew L, Rouhizadeh, Masoud, Perin, Jamie, Schumock, Grant, Nagy, Paul, Gray, Josh H, Malapati, Harsha. Patient Trajectories Among Persons Hospitalized for COVID-19: A Cohort Study. *Annals of internal medicine*; 2020.
527. Society, American Geriatrics. American Geriatrics Society (AGS) Policy Brief: COVID-19 and Nursing Homes. *J Am Geriatr Soc*; Apr 8 2020.
528. McMichael, T. M., Clark, S., Pogojans, S., Kay, M., Lewis, J., Baer, A., Kawakami, V., Lukoff, M. D., Ferro, J., Brostrom-Smith, C., Riedo, F. X., Russell, D., Hiatt, B., Montgomery, P., Rao, A. K., Currie, D. W., Chow, E. J., Tobolowsky, F., Bardossy, A. C., Oakley, L. P., Jacobs, J. R., Schwartz, N. G., Stone, N., Reddy, S. C., Jernigan, J. A., Honein, M. A., Clark, T. A., Duchin, J. S. COVID-19 in a Long-Term Care Facility - King County, Washington, February 27-March 9, 2020. *MMWR Morb Mortal Wkly Rep*; Mar 27 2020.
529. Kimball, A., Hatfield, K. M., Arons, M., James, A., Taylor, J., Spicer, K., Bardossy, A. C., Oakley, L. P., Tanwar, S., Chisty, Z., Bell, J. M., Methner, M., Harney, J., Jacobs, J. R., Carlson, C. M., McLaughlin, H. P., Stone, N., Clark, S., Brostrom-Smith, C., Page, L. C., Kay, M., Lewis, J., Russell, D., Hiatt, B., Gant, J., Duchin, J. S., Clark, T. A., Honein, M. A., Reddy, S. C., Jernigan, J. A. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in Residents of a Long-Term Care Skilled Nursing Facility - King County, Washington, March 2020. *MMWR Morb Mortal Wkly Rep*; Apr 3 2020.
530. Davidson, P. M., Szanton, S. L. Nursing homes and COVID-19: we can and should do better. *J Clin Nurs*; Apr 12 2020.
531. Mills, J. P., Kaye, K. S., Mody, L. COVID-19 in older adults: clinical, psychosocial, and public health considerations. *JCI Insight*; Apr 17 2020.
532. Tartof, Sara Y, Qian, Lei, Hong, Vennis, Wei, Rong, Nadjafi, Ron F, Fischer, Heidi, Li, Zhuoxin, Shaw, Sally F, Caparosa, Susan L, Nau, Claudia L. Obesity and mortality among patients diagnosed with COVID-19: results from an integrated health care organization. *Annals of internal medicine*; 2020.
533. Sabourin, Katherine R., Schultz, Jonathan, Romero, Joshua, Lamb, Molly M., Larremore, Daniel, Morrison, Thomas E., Frazer-Abel, Ashley, Zimmer, Shanta, Kedl, Ross M., Jaenisch, Thomas, Rochford, Rosemary. Risk Factors of SARS-CoV-2

Antibodies in Arapahoe County First Responders - the COVID-19 Arapahoe Serosurveillance Study (CASES) Project. *Journal of Occupational and Environmental Medicine*; 2020.

534. CDC. People Who Are at Higher Risk for Severe Illness. 2020.

535. Zhang, J., Wang, X., Jia, X., Li, J., Hu, K., Chen, G., Wei, J., Gong, Z., Zhou, C., Yu, H., Yu, M., Lei, H., Cheng, F., Zhang, B., Xu, Y., Wang, G., Dong, W. Risk factors for disease severity, unimprovement, and mortality of COVID-19 patients in Wuhan, China. *Clin Microbiol Infect*; Apr 15 2020.

536. Li, X., Xu, S., Yu, M., Wang, K., Tao, Y., Zhou, Y., Shi, J., Zhou, M., Wu, B., Yang, Z., Zhang, C., Yue, J., Zhang, Z., Renz, H., Liu, X., Xie, J., Xie, M., Zhao, J. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol*; Apr 12 2020.

537. Zuin, M., Rigatelli, G., Zuliani, G., Rigatelli, A., Mazza, A., Roncon, L. Arterial hypertension and risk of death in patients with COVID-19 infection: systematic review and meta-analysis. *J Infect*; Apr 10 2020.

538. Garibaldi, BT, Fiksel, J, Muschelli, J, Robinson, ML, Rouhizadeh, M, Perin, J, Schumock, G, Nagy, P, Gray, JH, Malapati, H. Patient trajectories among persons hospitalized for COVID-19: A cohort study. *Ann. Intern. Med.*; 2020.

539. Zhang, Yanping. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. 2020.

540. Troya, J., Bastard, P., Planas-Serra, L., Ryan, P., Ruiz, M., de Carranza, M., Torres, J., Martínez, A., Abel, L., Casanova, J., Pujol, A. Neutralizing autoantibodies to type I IFNs in >10% of patients with severe COVID-19 pneumonia hospitalized in Madrid, Spain. *J Clin Immunol*; Jul 2021.

541. Ray, J., Schull, M., Vermeulen, M., Park, A. Association between ABO and rh blood groups and SARS-CoV-2 infection or severe COVID-19 illness : a population-based cohort study. *Ann Intern Med*; Mar 2021.

542. Ellinghaus, D., Degenhardt, F., Bujanda, L., Buti, M., Albillos, A., Invernizzi, P., Fernández, J., Prati, D., Baselli, G., Asselta, R., Grimsrud, M. M., Milani, C., Aziz, F., Kässens, J., May, S., Wendorff, M., Wienbrandt, L., Uellendahl-Werth, F., Zheng, T., Yi, X., de Pablo, R., Chercoles, A. G., Palom, A., Garcia-Fernandez, A. E., Rodriguez-Frias, F., Zanella, A., Bandera, A., Protti, A., Aghemo, A., Lleo, A., Biondi, A., Caballero-Garralda, A., Gori, A., Tanck, A., Carreras Nolla, A., Latiano, A., Fracanzani, A. L., Peschuck, A., Julià, A., Pesenti, A., Voza, A., Jiménez, D., Mateos, B., Nafria Jimenez, B., Quereda, C., Paccapelo, C., Gassner, C., Angelini, C., Cea, C., Solier, A., Pestaña, D., Muñoz-Díaz, E., Sandoval, E., Paraboschi, E. M., Navas, E., García Sánchez, F., Ceriotti, F., Martinelli-Boneschi, F., Peyvandi, F., Blasi, F., Téllez, L., Blanco-Grau, A., Hemmrich-Stanisak, G., Grasselli, G., Costantino, G., Cardamone, G., Foti, G., Aneli, S., Kurihara, H., ElAbd, H., My, I., Galván-Femenia, I., Martín, J., Erdmann, J., Ferrusquía-Acosta, J., Garcia-Etxebarria, K., Izquierdo-Sanchez, L., Bettini, L. R., Sumoy, L., Terranova, L., Moreira, L., Santoro, L., Scudeller, L., Mesonero, F., Roade, L., Rühlemann, M. C., Schaefer, M., Carrabba, M., Riveiro-Barciela, M., Figuera Basso, M. E., Valsecchi, M. G., Hernandez-Tejero, M., Acosta-Herrera, M., D'Angiò, M., Baldini, M., Cazzaniga, M., Schulzky, M., Cecconi, M., Wittig, M., Ciccirelli, M., Rodríguez-Gandía, M., Boccione, M., Miozzo, M., Montano, N., Braun, N., Sacchi, N., Martínez, N., Özer, O., Palmieri, O., Faverio, P., Preatoni, P., Bonfanti, P., Omodei, P., Tentorio, P., Castro, P., Rodrigues, P. M., Blandino Ortiz, A., de Cid, R., Ferrer, R., Gualtierotti, R., Nieto, R., Goerg, S., Badalamenti, S., Marsal, S., Matullo, G., Pelusi, S., Juzenas, S., Aliberti, S., Monzani, V., Moreno, V., Wesse, T., Lenz, T. L., Pumarola, T., Rimoldi, V., Bosari, S., Albrecht, W., Peter, W., Romero-Gómez, M., D'Amato, M., Duga, S., Banales, J. M., Hov, J. R., Folseraas, T., Valenti, L., Franke, A., Karlsen, T. H. Genomewide Association Study of Severe Covid-19 with Respiratory Failure. *N Engl J Med*; Jun 17 2020.

543. CDC. COVID-19 Vaccination Communication Toolkit. 2021.

544. CDC. Customizable COVID-19 Vaccine Content for Essential Workers. 2021.

545. Richardson, S., Hirsch, J. S., Narasimhan, M., Crawford, J. M., McGinn, T., Davidson, K. W., Barnaby, D. P., Becker, L. B., Chelico, J. D., Cohen, S. L., Cookingham, J., Coppa, K., Diefenbach, M. A., Dominello, A. J., Duer-Hefele, J., Falzon, L., Gitlin, J., Hajizadeh, N., Harvin, T. G., Hirschwerk, D. A., Kim, E. J., Kozel, Z. M., Marrast, L. M., Mogavero, J. N., Osorio, G. A., Qiu, M., Zanos, T. P. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *Jama*; Apr 22 2020.

546. Stave, G., Smith, S., Hymel, P., Heron, R. Worksite temperature screening for COVID-19. *J Occup Environ Med*; Aug 1 2021.

547. Kaleem, J. Sick staff. Endless COVID patients. Doctors 'just scraping by' as Omicron sweeps hospitals. *Los Angeles Times*; January 12, 2022.

548. Smith, H. As hospitals reel, California tells coronavirus positive medical workers to stay on the job. *Los Angeles Times*; January 12, 2022.

549. CDC. CDC updates and shortens recommended isolation and quarantine period for general population. December 27, 2021.

550. CDC. How to Protect Yourself & Others. 2020.

551. CDC. Duration of Isolation and Precautions for Adults with COVID-19. 2020.
552. CDC. Return-to-Work Criteria. 2020.
553. CDC. Discontinuation of Isolation for Persons with COVID-19 Not in Healthcare Settings. 2020.
554. CDC. Interim Clinical Considerations for Use of mRNA COVID-19 Vaccines Currently Authorized in the United States. 2021.
555. CDC. Appendices. 2020.
556. CDC. Contact Tracing. 2020.
557. CDC. Interim U.S. Guidance for Risk Assessment and Public Health Management of Healthcare Personnel with Potential Exposure in a Healthcare Setting to Patients with Coronavirus Disease 2019 (COVID-19). 2020.
558. Mass.gov. Information and Guidance for Persons in Quarantine due to COVID-19. 2020.
559. CDC. When to Quarantine. 2020.
560. CDC. Interim Guidance for Implementing Safety Practices for Critical Infrastructure Workers Who May Have Had Exposure to a Person with Suspected or Confirmed COVID-19. 2020.
561. Williamson, Elizabeth, Walker, Alex J, Bhaskaran, Krishnan J, Bacon, Seb, Bates, Chris, Morton, Caroline E, Curtis, Helen J, Mehrkar, Amir, Evans, David, Inglesby, Peter. OpenSAFELY: factors associated with COVID-19-related hospital death in the linked electronic health records of 17 million adult NHS patients. *medRxiv*; 2020.
562. Nehme, M., Braillard, O., Chappuis, F., Courvoisier, D.S., Guessous, I., Team, CoviCare, Study. Prevalence of symptoms more than seven months after diagnosis of symptomatic COVID-19 in an outpatient setting. *Ann Intern Med*; 2021.
563. Mahase, E., Kmiotowicz, Z. Covid-19: Doctors are told not to perform CPR on patients in cardiac arrest. *Bmj*; Mar 29 2020.
564. Bowles, Kathryn H, McDonald, Margaret, Barrón, Yolanda, Kennedy, Erin, O'Connor, Melissa, Mikkelsen, Mark. Surviving COVID-19 After Hospital Discharge: Symptom, Functional, and Adverse Outcomes of Home Health Recipients. *Annals of internal medicine*; 2020.
565. NICE. COVID-19 rapid guideline: managing the long-term effects of COVID-19. 2020.
566. Tenforde, Mark W., Kim, Sara, Lindsell, Christopher J., Rose, Erica Billig, Shapiro, Nathan I., Files, D. Clark, Gibbs, Kevin W., Erickson, Heidi L., Steingrub, Jay S., Smithline, Howard A., Gong, Michelle N., Aboodi, Michael S., Exline, Matthew C., Henning, Daniel J., Wilson, Jennifer G., Khan, Akram, Qadir, Nida, Brown, Samuel M., Peltan, Ithan D., Rice, Todd W., Hager, David N., Ginde, Adit A., Stubblefield, William B., Patel, Manish M., Self, Wesley H., Feldstein, Leora R., Investigators, IVY, Network, Team, CDC, COVID-19, Response. Symptom Duration and Risk Factors for Delayed Return to Usual Health Among Outpatients with COVID-19 in a Multistate Health Care Systems Network — United States, March–June 2020. 2020.
567. DiSilvio, Briana, Young, Meilin, Gordon, Ayla, Malik, Khalid, Singh, Ashley, Cheema, Tariq. Complications and outcomes of acute respiratory distress syndrome. *Critical care nursing quarterly*; 2019.
568. Dinglas, V. D., Chessare, C. M., Davis, W. E., Parker, A., Friedman, L. A., Colantuoni, E., Bingham, C. O., Turnbull, A. E., Needham, D. M. Perspectives of survivors, families and researchers on key outcomes for research in acute respiratory failure. *Thorax*; Jan 2018.
569. Chen, Jiajia, Wu, Jie, Hao, Shaorui, Yang, Meifang, Lu, Xiaoqing, Chen, Xiaoxiao, Li, Lanjuan. Long term outcomes in survivors of epidemic Influenza A (H7N9) virus infection. *Scientific reports*; 2017.
570. Chiumello, Davide, Coppola, Silvia, Froio, Sara, Gotti, Miriam. What's next after ARDS: long-term outcomes. *Respiratory care*; 2016.
571. Herridge, Margaret S, Moss, Marc, Hough, Catherine L, Hopkins, Ramona O, Rice, Todd W, Bienvenu, O Joseph, Azoulay, Elie. Recovery and outcomes after the acute respiratory distress syndrome (ARDS) in patients and their family caregivers. *Intensive care medicine*; 2016.
572. Mason, Christopher, Dooley, Nessa, Griffiths, Mark. Acute respiratory distress syndrome. *Clinical Medicine*; 2016.
573. Rogers, Jonathan P, Chesney, Edward, Oliver, Dominic, Pollak, Thomas A, McGuire, Philip, Fusar-Poli, Paolo, Zandi, Michael S, Lewis, Glyn, David, Anthony S. Psychiatric and neuropsychiatric presentations associated with severe coronavirus infections: a systematic review and meta-analysis with comparison to the COVID-19 pandemic. *The Lancet Psychiatry*; 2020.
574. Society, Intensive Care. Psychology of COVID-19 critical care patients. 2020.
575. Wilcox, M. E., Patsios, D., Murphy, G., Kudlow, P., Paul, N., Tansey, C. M., Chu, L., Matte, A., Tomlinson, G., Herridge, M. S. Radiologic outcomes at 5 years after severe ARDS. *Chest*; Apr 2013.

576. Files, D. C., Sanchez, M. A., Morris, P. E. A conceptual framework: the early and late phases of skeletal muscle dysfunction in the acute respiratory distress syndrome. *Crit Care*; Jul 2 2015.
577. Disser, Nathaniel P., De Micheli, Andrea J., Schonk, Martin M., Konnaris, Maxwell A., Piacentini, Alexander N., Edon, Daniel L., Toresdahl, Brett G., Rodeo, Scott A., Casey, Ellen K., Mendias, Christopher L. Musculoskeletal Consequences of COVID-19. *JBJS*; 2020.
578. Hyman, Mark H., Talmage, James B., Hegmann, Kurt T. Evaluating Covid-19 Injury Claims With a Focus on Workers' Compensation. *Journal of Occupational and Environmental Medicine*; 2020.
579. Andersson, Gunnar B.J., Cocchiarella, Linda. AMA Guides to the Evaluation of Permanent Impairment, 5th edition. 2000.
580. Rondinelli, Robert D. AMA Guides to the Evaluation of Permanent Impairment, Sixth Edition. 2008.
581. Parker, Edward P. K., Shrotri, Madhumita, Kampmann, Beate. Keeping track of the SARS-CoV-2 vaccine pipeline. *Nature Reviews Immunology*; 2020/11/01.
582. Callaway, Ewen. The race for coronavirus vaccines: a graphical guide. 2020.
583. Zhu, Feng-Cai, Li, Yu-Hua, Guan, Xu-Hua, Hou, Li-Hua, Wang, Wen-Juan, Li, Jing-Xin, Wu, Shi-Po, Wang, Bu-Sen, Wang, Zhao, Wang, Lei. Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: a dose-escalation, open-label, non-randomised, first-in-human trial. *The Lancet*; 2020.
584. Lanese, Nicoletta. When will a COVID-19 vaccine be ready?. 2020.
585. Butt, A., Omer, S., Yan, P., Shaikh, O., Mayr, F. SARS-CoV-2 vaccine effectiveness in a high-risk national population in a real-world setting. *Ann. Intern. Med.*; 2021.
586. Thompson, M., Burgess, J., Naleway, A., Tyner, H., Yoon, S., Meece, J., Olsho, L., Caban-Martinez, A., Fowlkes, A., Lutrick, K. Prevention and attenuation of Covid-19 with the BNT162b2 and mRNA-1273 vaccines. *NEJM*; 2021.
587. Thompson, M., Burgess, J., Naleway, A., Tyner, H., Yoon, S., Meece, J., Olsho, L., Caban-Martinez, A., Fowlkes, A., Lutrick, K. Interim estimates of vaccine effectiveness of BNT162b2 and mRNA-1273 COVID-19 vaccines in preventing SARS-CoV-2 infection among health care personnel, first responders, and other essential and frontline workers—eight US locations, December 2020–March 2021. *MMWR*; 2021.
588. London School of Hygiene & Tropical Medicine. COVID-19 Vaccine Tracker. 2020.
589. Amit, Sharon, Regev-Yochay, Gili, Afek, Arnon, Kreiss, Yitshak, Leshem, Eyal. Early rate reductions of SARS-CoV-2 infection and COVID-19 in BNT162b2 vaccine recipients. *The Lancet*; 2021.
590. Ferdinands, J., Rao, S., Dixon, B., Mitchell, P., DeSilva, M., Irving, S., Lewis, N., Natarajan, K., Stenehjem, E., Grannis, S., Han, J., McEvoy, C., Ong, T., Naleway, A., Reese, S., Embi, P., Dascomb, K., Klein, N., Griggs, E., Konatham, D., Kharbanda, A., Yang, D., Fadel, W., Grisel, N., Goddard, K., Patel, P., Liao, I. C., Birch, R., Valvi, N., Reynolds, S., Arndorfer, J., Zerbo, O., Dickerson, M., Murthy, K., Williams, J., Bozio, C., Blanton, L., Verani, J., Schrag, S., Dalton, A., Wondimu, M., Link-Gelles, R., Azziz-Baumgartner, E., Barron, M., Gaglani, M., Thompson, M., Fireman, B. Waning 2-dose and 3-dose effectiveness of mRNA vaccines against COVID-19-associated emergency department and urgent care encounters and hospitalizations among adults during periods of delta and omicron variant predominance - VISION Network, 10 states, August 2021-January 2022. *MMWR*; Feb 18 2022.
591. Hause, A., Baggs, J., Marquez, P., Myers, T., Su, J., Blanc, P., Gwira Baumblatt, J., Woo, E., Gee, J., Shimabukuro, T., Shay, D. Safety monitoring of COVID-19 vaccine booster doses among adults - United States, September 22, 2021-February 6, 2022. *MMWR*; Feb 18 2022.
592. Smith-Jeffcoat, S., Pomeroy, M., Sleweon, S., Sami, S., Ricaldi, J., Gebru, Y., Walker, B., Brady, S., Christenberry, M., Bart, S., Vostok, J., Meyer, S., Seys, S., Markelz, A., Ditto, N., Newbern, V., Thomas, F., Thomas, D., Cabredo, E., Kellner, S., Brown, V., Tate, J., Kirking, H. Multistate outbreak of SARS-CoV-2 B.1.1.529 (omicron) variant infections among persons in a social network attending a convention - New York City, November 18-December 20, 2021. *MMWR*; Feb 18 2022.
593. CDC. Science Brief: Background Rationale and Evidence for Public Health Recommendations for Fully Vaccinated People. 2021.
594. Gazit, S., Shlezinger, R., Perez, G., Lotan, R., Peretz, A., Ben-Tov, A., Cohen, D., Muhsen, K., Chodick, G., Patalon, T. Comparing SARS-CoV-2 natural immunity to vaccine-induced immunity: reinfections versus breakthrough infections. *MedRxiv*; 2021.
595. CDC. Evaluating and Testing Persons for Coronavirus Disease 2019 (COVID-19). 2020.
596. Hahn, Stephen M., Shuren, Jeffrey E. Coronavirus (COVID-19) Update: FDA Authorizes First Antigen Test to Help in the Rapid Detection of the Virus that Causes COVID-19 in Patients. 2020.

597. Kucirka, L. M., Lauer, S. A., Laeyendecker, O., Boon, D., Lessler, J. Variation in False-Negative Rate of Reverse Transcriptase Polymerase Chain Reaction-Based SARS-CoV-2 Tests by Time Since Exposure. *Ann Intern Med*; Aug 18 2020.
598. Pérez, Daniel Alberto Girón, Ruiz-Manzano, Rocío Alejandra, Benitez-Trinidad, Alma Betsaida, Ventura-Ramón, Guadalupe Herminia, Covantes-Rosales, Carlos Eduardo, Ojeda-Durán, Ansonny Jhovanny, Mercado-Salgado, Ulises, Toledo-Ibarra, Gladys Alejandra, Díaz-Reséndiz, Karina Janice, Girón-Pérez, Manuel Iván. Saliva Pooling Strategy for the Large-Scale Detection of SARS-CoV-2, Through Working-Groups Testing of Asymptomatic Subjects for Potential Applications in Different Workplaces. *Journal of Occupational and Environmental Medicine*; 2021.
599. Wyllie, Anne Louise, Fournier, John, Casanovas-Massana, Arnau, Campbell, Melissa, Tokuyama, Maria, Vijayakumar, Pavithra, Geng, Bertie, Muenker, M Catherine, Moore, Adam J, Vogels, Chantal BF. Saliva is more sensitive for SARS-CoV-2 detection in COVID-19 patients than nasopharyngeal swabs. *medRxiv*; 2020.
600. Bastos, M., Perlman-Arrow, S., Menzies, D., Campbell, J. The sensitivity and costs of testing for SARS-CoV-2 infection with saliva versus nasopharyngeal swabs: a systematic review and meta-analysis. *Ann Intern Med*; Apr 2021.
601. Azzi, Lorenzo, Carcano, Giulio, Gianfagna, Francesco, Grossi, Paolo, Gasperina, Daniela Dalla, Genoni, Angelo, Fasano, Mauro, Sessa, Fausto, Tettamanti, Lucia, Carinci, Francesco, Maurino, Vittorio, Rossi, Agostino, Tagliabue, Angelo, Baj, Andreina. Saliva is a reliable tool to detect SARS-CoV-2. *Journal of Infection*; 2020/04/14/.
602. Mina, Michael J, Parker, Roy, Larremore, Daniel B. Rethinking Covid-19 test sensitivity—A strategy for containment. *New England Journal of Medicine*; 2020.
603. Burbelo, Peter D., Riedo, Francis X., Morishima, Chihiro, Rawlings, Stephen, Smith, Davey, Das, Sanchita, Strich, Jeffrey R., Chertow, Daniel S., Davey, Richard T., Jr., Cohen, Jeffrey I. Sensitivity in Detection of Antibodies to Nucleocapsid and Spike Proteins of Severe Acute Respiratory Syndrome Coronavirus 2 in Patients With Coronavirus Disease 2019. *The Journal of Infectious Diseases*; 2020.
604. Woelfel, Roman, Corman, Victor Max, Guggemos, Wolfgang, Seilmaier, Michael, Zange, Sabine, Mueller, Marcel A, Niemeyer, Daniela, Vollmar, Patrick, Rothe, Camilla, Hoelscher, Michael. Clinical presentation and virological assessment of hospitalized cases of coronavirus disease 2019 in a travel-associated transmission cluster. *medRxiv*; 2020.
605. Yang, Yang, Yang, Minghui, Shen, Chenguang, Wang, Fuxiang, Yuan, Jing, Li, Jinxiu, Zhang, Mingxia, Wang, Zhaoqin, Xing, Li, Wei, Jinli. Laboratory diagnosis and monitoring the viral shedding of 2019-nCoV infections. *medRxiv*; 2020.
606. Carver, Catherine, Jones, Nick. Comparative accuracy of oropharyngeal and nasopharyngeal swabs for diagnosis of COVID-19. 2020.
607. Favresse, J., Gillot, C., Oliveira, M., Cadrobbi, J., Elsen, M., Eucher, C., Laffineur, K., Rosseels, C., Van Eeckhoudt, S., Nicolas, J. B., Morimont, L., Dogné, J. M., Douxfils, J. Head-to-head comparison of rapid and automated antigen detection tests for the diagnosis of SARS-CoV-2 infection. *J Clin Med*; Jan 13 2021.
608. Iyer, Anita S, Jones, Forrest K, Nodoushanian, Ariana, Kelly, Meagan, Becker, Margaret, Slater, Damien, Mills, Rachel, Teng, Erica, Kamruzzaman, Mohammad, Garcia-Beltran, Wilfredo F, Astudillo, Michael, Yang, Diane, Miller, Tyler E, Oiver, Elizabeth, Fischinger, Stephanie, Atyeo, Caroline, lafrate, Anthony John, Calderwood, Stephen B, Lauer, Stephen A, Yu, Jingyou, Li, Zhenfeng, Feldman, Jared, Hauser, Blake M, Cardonna, Timothy M, Branda, John A, Turbett, Sarah E, LaRocque, Regina C, Mellon, Guillaume, Barouch, Dan H, Schmidt, Aaron G, Azman, Andrew S, Alter, Galit, Ryan, Edward T, Harris, Jason B, Charles, Richelle C. Dynamics and significance of the antibody response to SARS-CoV-2 infection. *medRxiv*; 2020.
609. Cheng, Matthew P, Yansouni, Cedric P, Basta, Nicole E, Desjardins, Michaël, Kanjilal, Sanjat, Paquette, Katryn, Caya, Chelsea, Semret, Makeda, Quach, Caroline, Libman, Michael. Serodiagnostics for Severe Acute Respiratory Syndrome–Related Coronavirus-2: A Narrative Review. *Annals of Internal Medicine*; 2020.
610. Guo, L., Ren, L., Yang, S., Xiao, M., Chang, D., Yang, F., Dela Cruz, C. S., Wang, Y., Wu, C., Xiao, Y., Zhang, L., Han, L., Dang, S., Xu, Y., Yang, Q., Xu, S., Zhu, H., Xu, Y., Jin, Q., Sharma, L., Wang, L., Wang, J. Profiling Early Humoral Response to Diagnose Novel Coronavirus Disease (COVID-19). *Clin Infect Dis*; Mar 21 2020.
611. Zhao, J., Yuan, Q., Wang, H., Liu, W., Liao, X., Su, Y., Wang, X., Yuan, J., Li, T., Li, J., Qian, S., Hong, C., Wang, F., Liu, Y., Wang, Z., He, Q., Li, Z., He, B., Zhang, T., Fu, Y., Ge, S., Liu, L., Zhang, J., Xia, N., Zhang, Z. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *Clin Infect Dis*; Mar 28 2020.
612. Li, Z., Yi, Y., Luo, X., Xiong, N., Liu, Y., Li, S., Sun, R., Wang, Y., Hu, B., Chen, W., Zhang, Y., Wang, J., Huang, B., Lin, Y., Yang, J., Cai, W., Wang, X., Cheng, J., Chen, Z., Sun, K., Pan, W., Zhan, Z., Chen, L., Ye, F. Development and clinical application of a rapid IgM-IgG combined antibody test for SARS-CoV-2 infection diagnosis. *J Med Virol*; Feb 27 2020.
613. Bonelli, F., Sarasini, A., Zierold, C., Calleri, M., Bonetti, A., Vismara, C., Blocki, F. A., Pallavicini, L., Chinali, A., Campisi, D., Percivalle, E., DiNapoli, A. P., Perno, C. F., Baldanti, F. Clinical and Analytical Performance of an Automated Serological Test That Identifies S1/S2-Neutralizing IgG in COVID-19 Patients Semiquantitatively. *J Clin Microbiol*; Aug 24 2020.
614. Jacofsky, David, Jacofsky, Emilia M, Jacofsky, Marc. Understanding antibody testing for covid-19. *The Journal of Arthroplasty*; 2020.

615. FDA. EUA Authorized Serology Test Performance. 2020.
616. Pollán, Marina, Pérez-Gómez, Beatriz, Pastor-Barriuso, Roberto, Oteo, Jesús, Hernán, Miguel A, Pérez-Olmeda, Mayte, Sanmartín, Jose L, Fernández-García, Aurora, Cruz, Israel, de Larrea, Nerea Fernández. Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study. *The Lancet*; 2020.
617. Caturegli, Giorgio, Materi, Joshua, Howard, Brittney M., Caturegli, Patrizio. Clinical Validity of Serum Antibodies to SARS-CoV-2: A Case-Control Study. *Annals of internal medicine*; 2020.
618. Ripperger, Tyler J, Uhrlaub, Jennifer L, Watanabe, Makiko, Wong, Rachel, Castaneda, Yvonne, Pizzato, Hannah A, Thompson, Mallory R, Bradshaw, Christine, Weinkauff, Craig C, Bime, Christian, Erickson, Heidi L, Knox, Kenneth, Bixby, Billie, Parthasarathy, Sairam, Chaudhary, Sachin, Natt, Bhupinder, Cristan, Elaine, El Aini, Tammer, Rischard, Franz, Campion, Janet, Chopra, Madhav, Insel, Michael, Sam, Afshin, Knepler, James L, Capaldi, Andrew P, Spier, Catherine M, Dake, Michael D, Edwards, Taylor, Kaplan, Matthew E, Scott, Serena Jain, Hypes, Cameron, Mosier, Jarrod, Harris, David T, Lafleur, Bonnie J, Sprissler, Ryan, Nikolich-Zugich, Janko, Bhattacharya, Deepta. Detection, prevalence, and duration of humoral responses to SARS-CoV-2 under conditions of limited population exposure. *medRxiv*; 2020.
619. Seow, Jeffrey, Graham, Carl, Merrick, Blair, Acors, Sam, Steel, Kathryn J. A., Hemmings, Oliver, Bryne, Aoife, Kouphou, Neophytos, Pickering, Suzanne, Galao, Rui, Betancor, Gilberto, Wilson, Harry D., Signell, Adrian W., Winstone, Helena, Kerridge, Claire, Temperton, Nigel, Snell, Luke, Bisnauthsing, Karen, Moore, Amelia, Green, Adrian, Martinez, Lauren, Stokes, Brielle, Honey, Johanna, Izquierdo-Barras, Alba, Arbane, Gill, Patel, Amita, Oconnell, Lorcan, O Hara, Geraldine, MacMahon, Eithne, Douthwaite, Sam, Nebbia, Gaia, Batra, Rahul, Martinez-Nunez, Rocio, Edgeworth, Jonathan D., Neil, Stuart J. D., Malim, Michael H., Doores, Katie. Longitudinal evaluation and decline of antibody responses in SARS-CoV-2 infection. *medRxiv*; 2020.
620. Sekine, Takuya, Perez-Potti, André, Rivera-Ballesteros, Olga, Straling, Kristoffer, Gorin, Jean-Baptiste, Olsson, Annika, Llewellyn-Lacey, Sian, Kamal, Habiba, Bogdanovic, Gordana, Muschiol, Sandra. Robust T cell immunity in convalescent individuals with asymptomatic or mild COVID-19. *BioRxiv*; 2020.
621. Nelde, Annika, Bilich, Tatjana, Heitmann, Jonas S, Maringer, Yacine, Salih, Helmut R, Roerden, Malte, Lübke, Maren, Bauer, Jens, Rieth, Jonas, Wacker, Marcel. SARS-CoV-2 T-cell epitopes define heterologous and COVID-19-induced T-cell recognition. 2020.
622. Blanchard, Sam. Blow to getting Britain back to work after Oxford scientist tasked with evaluating crucial coronavirus antibody tests says it may take a MONTH before one is ready for Britain to use as another expert warns the kits may only be 50% accurate. 2020.
623. Canada, Government,of. Diagnostic devices for use against coronavirus (COVID-19): List of applications received. 2020.
624. NIH. Coronavirus Disease 2019 (COVID-19) Treatment Guidelines. 2020.
625. NIH. COVID-19 Treatment Guidelines Introduction. 2020.
626. IDSA. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19. 2020.
627. Bai, Chunxue, Chotirmall, Sanjay H, Rello, Jordi, Alba, George A, Ginns, Leo C, Krishnan, Jerry A, Rogers, Robert, Bendstrup, Elisabeth, Burgel, Pierre-Regis, Chalmers, James D. Updated guidance on the management of COVID-19: From an american thoracic society/european respiratory society coordinated international task force (29 July 2020). *European Respiratory Review*; 2020.
628. CDC. Interim Clinical Guidance for Management of Patients with Confirmed Coronavirus Disease (COVID-19). 2020.
629. SHEA. Novel Coronavirus 2019 (2019-nCoV) Resources. 2020.
630. ASHP. Assessment of Evidence for COVID-19-Related Treatments: Updated 12/01/2020. 2020.
631. WHO. Clinical management of COVID-19. 2020.
632. WHO. Therapeutics and COVID-19: living guideline. January 14, 2022.
633. American College of Physicians (ACP). Coronavirus disease 2019 (COVID-19): information for internists. 2022.
634. National Institutes of Health (NIH). Prevention of SARS-CoV-2 Infection. February 1, 2022.
635. Derwand, Roland, Scholz, Martin, Zelenko, Vladimir. COVID-19 outpatients—early risk-stratified treatment with zinc plus low dose hydroxychloroquine and azithromycin: a retrospective case series study. 2020.
636. Skalny, A. V., Rink, L., Ajsuvakova, O. P., Aschner, M., Gritsenko, V. A., Alekseenko, S. I., Svistunov, A. A., Petrakis, D., Spandidos, D. A., Aaseth, J., Tsatsakis, A., Tinkov, A. A. Zinc and respiratory tract infections: Perspectives for COVID-19 (Review). *Int J Mol Med*; Jul 2020.

637. Finzi, Eric. Treatment of SARS-CoV-2 with high dose oral zinc salts: A report on four patients. *International Journal of Infectious Diseases*; 2020.
638. Carlucci, Philip, Ahuja, Tania, Petrilli, Christopher M, Rajagopalan, Harish, Jones, Simon, Rahimian, Joseph. Hydroxychloroquine and azithromycin plus zinc vs hydroxychloroquine and azithromycin alone: outcomes in hospitalized COVID-19 patients. *medRxiv*; 2020.
639. Thomas, Suma, Patel, Divyang, Bittel, Barbara, Wolski, Kathy, Wang, Qiuqing, Kumar, Anirudh, Il'Giovine, Zachary J., Mehra, Reena, McWilliams, Carla, Nissen, Steve E., Desai, Milind Y. Effect of High-Dose Zinc and Ascorbic Acid Supplementation vs Usual Care on Symptom Length and Reduction Among Ambulatory Patients With SARS-CoV-2 Infection: The COVID A to Z Randomized Clinical Trial. *JAMA Network Open*; 2021.
640. Alipio, Mark. Vitamin D Supplementation Could Possibly Improve Clinical Outcomes of Patients Infected with Coronavirus-2019 (COVID-19). *Available at SSRN 3571484*; 2020.
641. Ruiz-Irastorza, G, Egurbide, MV, Olivares, N, Martinez-Berriotxo, Aguirre, Aguirre, C. Vitamin D deficiency in systemic lupus erythematosus: prevalence, predictors and clinical consequences. *Rheumatology*; 2008.
642. Lau, Frank H, Majumder, Rinku, Torabi, Radbeh, Saeg, Fouad, Hoffman, Ryan, Cirillo, Jeffrey D, Greiffenstein, Patrick. Vitamin D insufficiency is prevalent in severe COVID-19. *medRxiv*; 2020.
643. D'Avolio, A., Avataneo, V., Manca, A., Cusato, J., De Nicolò, A., Lucchini, R., Keller, F., Cantù, M. 25-Hydroxyvitamin D Concentrations Are Lower in Patients with Positive PCR for SARS-CoV-2. *Nutrients*; May 9 2020.
644. Zhang, Xiao-Jing, Qin, Juan-Juan, Cheng, Xu, Shen, Lijun, Zhao, Yan-Ci, Yuan, Yufeng, Lei, Fang, Chen, Ming-Ming, Yang, Huilin, Bai, Liangjie. In-hospital Use of Statins is Associated with a Reduced Risk of Mortality among Individuals with COVID-19. *Cell metabolism*; 2020.
645. Quesada-Gomez, J., Entrenas-Castillo, M., Bouillon, R. Vitamin D receptor stimulation to reduce acute respiratory distress syndrome (ARDS) in patients with coronavirus SARS-CoV-2 infections: revised Ms SBMB 2020_166. *J Steroid Biochem Mol Biol*; Sep 2020.
646. Oxford University. Low-cost dexamethasone reduces death by up to one third in hospitalised patients with severe respiratory complications of COVID-19. 2020.
647. Ledford, Heidi. Coronavirus breakthrough: dexamethasone is first drug shown to save lives. 2020.
648. Russell, Beth, Moss, Charlotte, Rigg, Anne, Van Hemelrijck, Mieke. COVID-19 and treatment with NSAIDs and corticosteroids: should we be limiting their use in the clinical setting?. *ecancermedicalscience*; 2020.
649. Beigel, John H., Tomashek, Kay M., Dodd, Lori E., Mehta, Aneesh K., Zingman, Barry S., Kalil, Andre C., Hohmann, Elizabeth, Chu, Helen Y., Luetkemeyer, Annie, Kline, Susan, Lopez de Castilla, Diego, Finberg, Robert W., Dierberg, Kerry, Tapsen, Victor, Hsieh, Lanny, Patterson, Thomas F., Paredes, Roger, Sweeney, Daniel A., Short, William R., Touloumi, Giota, Lye, David Chien, Ohmagari, Norio, Oh, Myoung-don, Ruiz-Palacios, Guillermo M., Benfield, Thomas, Fätkenheuer, Gerd, Kortepeter, Mark G., Atmar, Robert L., Creech, C. Buddy, Lundgren, Jens, Babiker, Abdel G., Pett, Sarah, Neaton, James D., Burgess, Timothy H., Bonnett, Tyler, Green, Michelle, Makowski, Mat, Osinusi, Anu, Nayak, Seema, Lane, H. Clifford. Remdesivir for the Treatment of Covid-19 — Preliminary Report. *New England Journal of Medicine*; 2020.
650. Winck, João Carlos, Ambrosino, Nicolino. COVID-19 pandemic and non invasive respiratory management: every Goliath needs a David. An evidence based evaluation of problems. *Pulmonology*; 2020.
651. Lazzeri, Marta, Lanza, Andrea, Bellini, Raffaella, Bellofiore, Angela, Cecchetto, Simone, Colombo, Alessia, D'Abrosca, Francesco, Del Monaco, Cesare, Gaudellio, Giuseppe, Paneroni, Mara. Respiratory physiotherapy in patients with COVID-19 infection in acute setting: a Position Paper of the Italian Association of Respiratory Physiotherapists (ARIR). *Monaldi Archives for Chest Disease*; 2020.
652. Poston, J. T., Patel, B. K., Davis, A. M. Management of Critically Ill Adults With COVID-19. *Jama*; Mar 26 2020.
653. Khan, Fasihul, Fabbri, Laura, Stewart, Iain, Robinson, Karen, Smyth, Alan R, Jenkins, Gisli. A systematic review of Anakinra, Tocilizumab, Sarilumab and Siltuximab for coronavirus-related infections. *medRxiv*; 2020.
654. CHEN, Jun, LIU, Danping, LIU, Li, LIU, Ping, XU, Qingnian, XIA, Lu, LING, Yun, HUANG, Dan, SONG, Shuli, ZHANG, Dandan. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *Journal of Zhejiang University (Medical Science)*; 2020.
655. Ladapo, Joseph A, McKinnon, John E, McCullough, Peter A, Risch, Harvey. Randomized Controlled Trials of Early Ambulatory Hydroxychloroquine in the Prevention of COVID-19 Infection, Hospitalization, and Death: Meta-Analysis. *medRxiv*; 2020.
656. NIH. ClinicalTrials.gov. 2020.

657. Carrouel, F., Valette, M., Gadea, E., Esparcieux, A., Illes, G., Langlois, M., Perrier, H., Dussart, C., Tramini, P., Ribaud, M. Use of an antiviral mouthwash as a barrier measure in the SARS-CoV-2 transmission in adults with asymptomatic to mild COVID-19: A multicentre, randomized, double-blind controlled trial. *CMI*; 2021.
658. Di Domênico, M., Cesca, H., Ponciano, T., Dos Santos, R., Lenz, U., Antunes, V., Godinho, V., Collares, K., Corazza, P. Effectiveness of hydrogen peroxide as auxiliary treatment for hospitalized COVID-19 patients in Brazil: preliminary results of a randomized double-blind clinical trial. *Epidemiol Health*; 2021.
659. Choudhury, M., Shabnam, N., Ahsan, T., Kabir, M., Ahsan, R. Effect of 1% povidone iodine mouthwash/gargle, nasal and eye drop in COVID-19 patient. 2021.
660. Poleti, M., Gregório, D., Bistaffa, A., Fernandes, K., Vilhena, F., da Silva Santos, P., Lozovoy, M., Tatibana, B., Fernandes, T. The use of a mouthwash and a dentifrice containing antimicrobial phthalocyanine derivative on the reduction of clinical symptoms of COVID-19: a randomized triple-blinded clinical trial. 2021.
661. Costa, A., Pontes, F., Souza, L., Lopes, M., Santos-Silva, A., Vargas, P., Andrade, B., Bezerra, K., Romañach, M., Gomez, R., Costa, R., Júnior, D. What is the frequency of floor of the mouth lesions? A descriptive study of 4,016 cases. *Med Oral Patol Oral Cir Bucal*; Nov 1 2021.
662. Guenezan, J., Garcia, M., Strasters, D., Jousselin, C., Lévêque, N., Frasca, D., Mimoz, O. Povidone iodine mouthwash, gargle, and nasal spray to reduce nasopharyngeal viral load in patients with COVID-19: a randomized clinical trial. *JAMA Otolaryngol Head Neck Surg*; Apr 1 2021.
663. Mukhtar, K., Qassim, S., Al Qahtani, S., Danjuma, M., Mohamedali, M., Al Farhan, H., Khudair, M., El Tayeh, A., Al-Dosari, M., Babiker, M. A randomized trial on the regular use of potent mouthwash in COVID-19 treatment. *medRxiv*; 2021.
664. Acuña-Castroviejo, D., Escames, G., Figueira, J., de la Oliva, P., Borobia, A., Acuña-Fernández, C. Clinical trial to test the efficacy of melatonin in COVID-19. *J Pineal Res*; Oct 2020.
665. Brown, G., Pandi-Perumal, S., Pupko, H., Kennedy, J., Cardinali, D. Melatonin as an add-on treatment of COVID-19 infection: current status. *Diseases*; Sep 20 2021.
666. Alizadeh, Z., Keyhanian, N., Ghaderkhani, S., Dashti-Khavidaki, S., Shokouhi Shoormasti, R., Pourpak, Z. A pilot study on controlling coronavirus disease 2019 (COVID-19) inflammation using melatonin supplement. *Iran J Allergy Asthma Immunol*; Aug 7 2021.
667. Farnoosh, G., Akbariomi, M., Badri, T., Bagheri, M., Izadi, M., Saeedi-Boroujeni, A., Rezaie, E., Ghaleh, H., Aghamollaei, H., Fasihi-Ramandi, M., Hassanpour, K., Alishiri, G. Efficacy of a low dose of melatonin as an adjunctive therapy in hospitalized patients with COVID-19: a randomized, double-blind clinical trial. *Arch Med Res*; Jan 2022.
668. Hasan, Z., Atrakji, D., Mehuaiden, D. The effect of melatonin on thrombosis, sepsis and mortality rate in COVID-19 patients. *Int J Infect Dis*; Jan 2022.
669. Mousavi, S., Heydari, K., Mehravaran, H., Saeedi, M., Alizadeh-Navaei, R., Hedayatizadeh-Omran, A., Shamshirian, A. Melatonin effects on sleep quality and outcomes of COVID-19 patients: an open-label, randomized, controlled trial. *J Med Virol*; Jan 2022.
670. García, I., Rodríguez-Rubio, M., Mariblanca, A., de Soto, L., García, L., Villatoro, J., Parada, J., Meseguer, E., Rosales, M., González, J., Arribas, J., Carcas, A., de la Oliva, P., Borobia, A. A randomized multicenter clinical trial to evaluate the efficacy of melatonin in the prophylaxis of SARS-CoV-2 infection in high-risk contacts (MeCOVID Trial): a structured summary of a study protocol for a randomised controlled trial. *Trials*; Jun 3 2020.
671. Arnold, Kevin D., Skillings, Jared. Adapted Treatment Protocol for COVID19-Related Healthcare Professionals: A Holistic Model and Clinical Health Application of Cognitive-Behavioral Therapy to Pandemics. 2020.
672. Trust, Well Being. The COVID Pandemic Could Lead to 75,000 Additional Deaths from Alcohol and Drug Misuse and Suicide. 2020.
673. Arnold, Kevin D., Skillings, Jared L. Treating front-line workers: A step-by-step guide. 2020.
674. Arnold, Kevin D., Skillings, Jared L. Treating anxiety and stress in front-line workers: A step-by-step CBT guide. 2020.
675. Association, American, Psychological. Serious mental illness and COVID-19: How to help your patients right now. 2020.
676. Trivedi, Nidhi. COVID-19 (Coronavirus) Mental Health Guide: Strategies to manage mood effectively in times of Global Distress. 2020.
677. US Department of Veterans Affairs. COVID Coach. 2020.
678. Daily Mail. 28 August 2020 News Archive. 2020.

679. Wilson, Jenna M., Lee, Jerin, Fitzgerald, Holly N., Oosterhoff, Benjamin, Sevi, Baris, Shook, Natalie J. Job Insecurity and Financial Concern During the COVID-19 Pandemic Are Associated With Worse Mental Health. *Journal of Occupational and Environmental Medicine*; 2020.
680. Reiken, S., Sittenfeld, L., Dridi, H., Liu, Y., Liu, X., Marks, A.R. Alzheimer's-like signaling in brains of COVID-19 patients. *Alzheimer's Dement.*; 2022.
681. Tang, Wei, Cao, Zhujun, Han, Mingfeng, Wang, Zhengyan, Chen, Junwen, Sun, Wenjin, Wu, Yaojie, Xiao, Wei, Liu, Shengyong, Chen, Erzhen. Hydroxychloroquine in patients with mainly mild to moderate coronavirus disease 2019: open label, randomised controlled trial. *bmj*; 2020.
682. Tang, Wei, Cao, Zhujun, Han, Mingfeng, Wang, Zhengyan, Chen, Junwen, Sun, Wenjin, Wu, Yaojie, Xiao, Wei, Liu, Shengyong, Chen, Erzhen. Hydroxychloroquine in patients with COVID-19: an open-label, randomized, controlled trial. *medRxiv*; 2020.
683. te Velthuis, Aartjan J. W., van den Worm, Sjoerd H. E., Sims, Amy C., Baric, Ralph S., Snijder, Eric J., van Hemert, Martijn J. Zn²⁺ Inhibits Coronavirus and Arterivirus RNA Polymerase Activity In Vitro and Zinc Ionophores Block the Replication of These Viruses in Cell Culture. *PLOS Pathogens*; 2010.
684. Xue, Jing, Moyer, Amanda, Peng, Bing, Wu, Jinchang, Hannafon, Bethany N., Ding, Wei-Qun. Chloroquine Is a Zinc Ionophore. *PLOS ONE*; 2014.
685. Borba, Mayla Gabriela Silva, Val, Fernando Fonseca Almeida, Sampaio, Vanderson Souza, Alexandre, Marcia Almeida Araújo, Melo, Gisely Cardoso, Brito, Marcelo, Mourão, Maria Paula Gomes, Brito-Sousa, José Diego, Baía-da-Silva, Djane, Guerra, Marcus Vinitius Farias. Effect of High vs Low Doses of Chloroquine Diphosphate as Adjunctive Therapy for Patients Hospitalized With Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection: A Randomized Clinical Trial. *JAMA Network Open*; 2020.
686. Capra, R., De Rossi, N., Mattioli, F., Romanelli, G., Scarpazza, C., Sormani, M. P., Cossi, S. Impact of low dose tocilizumab on mortality rate in patients with COVID-19 related pneumonia. *Eur J Intern Med*; Jun 2020.
687. Lagier, Jean-Christophe, Million, Matthieu, Gautret, Philippe, Colson, Philippe, Cortaredona, Sébastien, Giraud-Gatineau, Audrey, Honoré, Stéphane, Gaubert, Jean-Yves, Fournier, Pierre-Edouard, Tissot-Dupont, Hervé. Outcomes of 3,737 COVID-19 patients treated with hydroxychloroquine/azithromycin and other regimens in Marseille, France: A retrospective analysis. *Travel medicine and infectious disease*; 2020.
688. Gautret, Philippe, Lagier, Jean-Christophe, Parola, Philippe, Meddeb, Line, Mailhe, Morgane, Doudier, Barbara, Courjon, Johan, Giordanengo, Valérie, Vieira, Vera Esteves, Dupont, Hervé Tissot. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents*; 2020.
689. Lover, Andrew A. Quantifying treatment effects of hydroxychloroquine and azithromycin for COVID-19: a secondary analysis of an open label non-randomized clinical trial (Gautret et al, 2020). *medRxiv*; 2020.
690. Raoult, Didier. Abstract. 2020.
691. Raoult, Didier. Table 1. 2020.
692. Davido, Benjamin, Lansaman, Thibaud, Lawrence, Christine, Alvarez, Jean-Claude, Bouchand, Frederique, Moine, Pierre, Perronne, Veronique, Le Gal, Aurelie, Annane, Djillali, Perronne, Christian. Hydroxychloroquine plus azithromycin: a potential interest in reducing in-hospital morbidity due to COVID-19 pneumonia (HI-ZY-COVID)? *medRxiv*; 2020.
693. Rosenberg, E. S., Dufort, E. M., Udo, T., Wilberschied, L. A., Kumar, J., Tesoriero, J., Weinberg, P., Kirkwood, J., Muse, A., DeHovitz, J., Blog, D. S., Hutton, B., Holtgrave, D. R., Zucker, H. A. Association of Treatment With Hydroxychloroquine or Azithromycin With In-Hospital Mortality in Patients With COVID-19 in New York State. *Jama*; May 11 2020.
694. Sbidian, Emilie, Josse, Julie, Lemaitre, Guillaume, Mayer, Imke, Bernaux, Melodie, Gramfort, Alexandre, Lapidus, Nathanael, Paris, Nicolas, Neuraz, Antoine, Lerner, Ivan. Hydroxychloroquine with or without azithromycin and in-hospital mortality or discharge in patients hospitalized for COVID-19 infection: a cohort study of 4,642 in-patients in France. *medRxiv*; 2020.
695. Mahévas, M., Tran, V. T., Roumier, M., Chabrol, A., Paule, R., Guillaud, C., Fois, E., Lepeule, R., Szwebel, T. A., Lescure, F. X., Schlemmer, F., Matignon, M., Khellaf, M., Crickx, E., Terrier, B., Morbieu, C., Legendre, P., Dang, J., Schoindre, Y., Pawlotsky, J. M., Michel, M., Perrodeau, E., Carlier, N., Roche, N., de Lastours, V., Ourghanlian, C., Kerneis, S., Ménager, P., Mouthon, L., Audureau, E., Ravaut, P., Godeau, B., Gallien, S., Costedoat-Chalumeau, N. Clinical efficacy of hydroxychloroquine in patients with covid-19 pneumonia who require oxygen: observational comparative study using routine care data. *Bmj*; May 14 2020.
696. Boulware, David R., Pullen, Matthew F., Bangdiwala, Ananta S., Pastick, Katelyn A., Lofgren, Sarah M., Okafor, Elizabeth C., Skipper, Caleb P., Nascene, Alanna A., Nicol, Melanie R., Abassi, Mahsa, Engen, Nicole W., Cheng, Matthew P., LaBar, Derek, Lothar, Sylvain A., MacKenzie, Lauren J., Drobot, Glen, Marten, Nicole, Zarychanski, Ryan, Kelly, Lauren E.,

- Schwartz, Ilan S., McDonald, Emily G., Rajasingham, Radha, Lee, Todd C., Hullsiek, Kathy H. A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for Covid-19. *New England Journal of Medicine*; 2020.
697. Lane, Jennifer CE, Weaver, James, Kostka, Kristin, Duarte-Salles, Talita, Abrahao, Maria Tereza F, Alghoul, Heba, Alser, Osaïd, Alshammari, Thami M, Biedermann, Patricia, Burn, Edward. Safety of hydroxychloroquine, alone and in combination with azithromycin, in light of rapid wide-spread use for COVID-19: a multinational, network cohort and self-controlled case series study. *medRxiv*; 2020.
698. Geleris, Joshua, Sun, Yifei, Platt, Jonathan, Zucker, Jason, Baldwin, Matthew, Hripcsak, George, Labella, Angelena, Manson, Daniel K, Kubin, Christine, Barr, R Graham. Observational study of hydroxychloroquine in hospitalized patients with Covid-19. *New England Journal of Medicine*; 2020.
699. Magagnoli, Joseph, Narendran, Siddharth, Pereira, Felipe, Cummings, Tammy, Hardin, James W., Sutton, S. Scott, Ambati, Jayakrishna. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. *medRxiv*; 2020.
700. Risch, Harvey A. Early Outpatient Treatment of Symptomatic, High-Risk Covid-19 Patients that Should be Ramped-Up Immediately as Key to the Pandemic Crisis. *American Journal of Epidemiology*; 2020.
701. Oxford University. No clinical benefit from use of hydroxychloroquine in hospitalised patients with COVID-19. 2020.
702. Cavalcanti, Alexandre B., Zampieri, Fernando G., Rosa, Regis G., Azevedo, Luciano C. P., Veiga, Viviane C., Avezum, Alvaro, Damiani, Lucas P., Marcadenti, Aline, Kawano-Dourado, Letícia, Lisboa, Thiago, Junqueira, Debora L. M., de Barros e Silva, Pedro G. M., Tramuja, Lucas, Abreu-Silva, Erlon O., Laranjeira, Ligia N., Soares, Aline T., Echenique, Leandro S., Pereira, Adriano J., Freitas, Flávio G. R., Gebara, Otávio C. E., Dantas, Vicente C. S., Furtado, Remo H. M., Milan, Eveline P., Golin, Nicole A., Cardoso, Fábio F., Maia, Israel S., Hoffmann Filho, Conrado R., Kormann, Adrian P. M., Amazonas, Roberto B., Bocchi de Oliveira, Monalisa F., Serpa-Neto, Ary, Falavigna, Maicon, Lopes, Renato D., Machado, Flávia R., Berwanger, Otavio. Hydroxychloroquine with or without Azithromycin in Mild-to-Moderate Covid-19. *New England Journal of Medicine*; 2020.
703. Voisin, O., Lorc'h, E. L., Mahé, A., Azria, P., Borie, M. F., Hubert, S., Ménage, E., Guillermin, J. C., Mourad, J. J. Acute QT Interval Modifications During Hydroxychloroquine-Azithromycin Treatment in the Context of COVID-19 Infection. *Mayo Clin Proc*; Aug 2020.
704. Mercuro, Nicholas J, Yen, Christina F, Shim, David J, Maher, Timothy R, McCoy, Christopher M, Zimetbaum, Peter J, Gold, Howard S. Risk of QT interval prolongation associated with use of hydroxychloroquine with or without concomitant azithromycin among hospitalized patients testing positive for coronavirus disease 2019 (COVID-19). *JAMA cardiology*; 2020.
705. Ramireddy, A., Chugh, H., Reinier, K., Ebinger, J., Park, E., Thompson, M., Cingolani, E., Cheng, S., Marban, E., Albert, C. M., Chugh, S. S. Experience With Hydroxychloroquine and Azithromycin in the Coronavirus Disease 2019 Pandemic: Implications for QT Interval Monitoring. *J Am Heart Assoc*; Jun 16 2020.
706. Pothén, Lucie, Yildiz, Halil, De Greef, Julien, Penaloza, Andrea, Beauloye, Christophe, Belkhir, Leila, Yombi, Jean Cyr. Safety use of hydroxychloroquine and its combination with azithromycin in the context of Sars-CoV-2 outbreak: Clinical experience in a Belgian tertiary center. *Travel Medicine and Infectious Disease*; 2020.
707. Kuate, Liliane Mfeukeu, Ngatchou, William, Temgoua, Mazou Ngou, Kouanfack, Charles, Lemogoum, Daniel, Tochie, Joel Noutakdie, Zamsi, Armel, Fomete, Lauriane, Lekelem, Skinner, Zamsi, Sylvain. Electrocardiographic safety of daily Hydroxychloroquine 400mg plus Azithromycin 250mg as an ambulatory treatment for COVID-19 patients in Cameroon. *medRxiv*; 2020.
708. Maraj, Ilir, Hummel, James P, Taoutel, Roy, Chamoun, Romy, Workman, Virginia, Li, Cindy, Tran, Lydia, DelVecchio, Alexander, Howes, Christopher, Akar, Joseph G. Incidence and Determinants of QT Interval Prolongation in COVID-19 Patients Treated with Hydroxychloroquine and Azithromycin. *Journal of cardiovascular electrophysiology*; 2020.
709. Chorin, Ehud, Dai, Matthew, Shulman, Eric, Wadhvani, Laila, Cohen, Roi Bar, Barbhaiya, Chirag, Aizer, Anthony, Holmes, Douglas, Bernstein, Scott, Soinelli, Michael. The QT Interval in Patients with SARS-CoV-2 Infection Treated with Hydroxychloroquine/Azithromycin. *medRxiv*; 2020.
710. Chorin, E., Wadhvani, L., Magnani, S., Dai, M., Shulman, E., Nadeau-Routhier, C., Knotts, R., Bar-Cohen, R., Kogan, E., Barbhaiya, C., Aizer, A., Holmes, D., Bernstein, S., Spinelli, M., Park, D. S., Stefano, C., Chinitz, L. A., Jankelson, L. QT interval prolongation and torsade de pointes in patients with COVID-19 treated with hydroxychloroquine/azithromycin. *Heart Rhythm*; May 12 2020.
711. Saleh, M., Gabriels, J., Chang, D., Soo Kim, B., Mansoor, A., Mahmood, E., Makker, P., Ismail, H., Goldner, B., Willner, J., Beldner, S., Mitra, R., John, R., Chinitz, J., Skipitaris, N., Mountantonakis, S., Epstein, L. M. Effect of Chloroquine, Hydroxychloroquine, and Azithromycin on the Corrected QT Interval in Patients With SARS-CoV-2 Infection. *Circ Arrhythm Electrophysiol*; Jun 2020.

712. Omrani, Ali S, Pathan, Sameer A, Thomas, Sarah A, Harris, Tim RE, Coyle, Peter V, Thomas, Caroline E, Qureshi, Isma, Bhutta, Zain A, Al Mawlawi, Naema, Al Kahlout, Reham. Randomized double-blinded placebo-controlled trial of hydroxychloroquine with or without azithromycin for virologic cure of non-severe Covid-19. *EClinicalMedicine*; 2020.
713. Catteau, Lucy, Dauby, Nicolas, Montourcy, Marion, Bottieau, Emmanuel, Hautekiet, Joris, Goetghebeur, Els, van Ierssel, Sabrina, Duysburgh, Els, Van Oyen, Herman, Wyndham-Thomas, Chloé. Low-dose hydroxychloroquine therapy and mortality in hospitalised patients with COVID-19: a nationwide observational study of 8075 participants. *International journal of antimicrobial agents*; 2020.
714. Cipriani, Alberto, Zorzi, Alessandro, Ceccato, Davide, Capone, Federico, Parolin, Matteo, Donato, Filippo, Fioretto, Paola, Pesavento, Raffaele, Previato, Lorenzo, Maffei, Pietro, Saller, Alois, Avogaro, Angelo, Sarais, Cristiano, Gregori, Dario, Iliceto, Sabino, Vettor, Roberto. Arrhythmic profile and 24-hour QT interval variability in COVID-19 patients treated with hydroxychloroquine and azithromycin. *International journal of cardiology*; 2020.
715. Ip, Andrew, Berry, Donald A, Hansen, Eric, Goy, Andre H, Pecora, Andrew L, Sinclair, Brittany A, Bednarz, Urszula, Marafelias, Michael, Berry, Scott M, Berry, Nicholas S. Hydroxychloroquine and Tocilizumab Therapy in COVID-19 Patients- An Observational Study. *medRxiv*; 2020.
716. Marzolini, Catia, Stader, Felix, Stoeckle, Marcel, Franzeck, Fabian, Egli, Adrian, Bassetti, Stefano, Hollinger, Alexa, Osthoff, Michael, Weisser, Maja, Gebhard, Caroline E. Effect of systemic inflammatory response to SARS-CoV-2 on lopinavir and hydroxychloroquine plasma concentrations. *Antimicrobial agents and chemotherapy*; 2020.
717. Nagaraja, Bada Sharanappa, Ramesh, Kalhalli Narayanaswamy, Dhar, Debjyoti, Mondal, Mahammad Samim, Dey, Treshita, Saha, Subhrakamal, Khan, Mumtaz Ali, Rutul, Shah Dhiren, Pratik, Kishore, Manjula, Jayaram. HyPE study: hydroxychloroquine prophylaxis-related adverse events' analysis among healthcare workers during COVID-19 pandemic: a rising public health concern. *Journal of Public Health*; 2020.
718. Yu, Bo, Li, Chenze, Chen, Peng, Zhou, Ning, Wang, Luyun, Li, Jia, Jiang, Hualiang, Wang, Dao-Wen. Low dose of hydroxychloroquine reduces fatality of critically ill patients with COVID-19. *Science China Life Sciences*; 2020.
719. Sekhavati, Ehsan, Jafari, Fatemeh, SeyedAlinaghi, SeyedAhmad, Jamalimoghadamsiahkali, Saeidreza, Sadr, Sara, Tabarestani, Mohammad, Pirhayati, Mohammad, Zendehehdel, Abolfazl, Manafi, Navid, Hajiabdolbaghi, Mahboubbeh, Ahmadinejad, Zahra, Kouchak, Hamid Emadi, Jafari, Siros, Khalili, Hosein, Salehi, Mohamadreza, Seifi, Arash, Golestan, Fereshteh Shahmari, Ghiasvand, Fereshteh. Safety and effectiveness of azithromycin in patients with COVID-19: An open-label randomised trial. *International journal of antimicrobial agents*; 2020.
720. Liu, Jia, Cao, Ruiyuan, Xu, Mingyue, Wang, Xi, Zhang, Huanyu, Hu, Hengrui, Li, Yufeng, Hu, Zhihong, Zhong, Wu, Wang, Manli. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discovery*; 2020.
721. Yao, Xueting, Ye, Fei, Zhang, Miao, Cui, Cheng, Huang, Baoying, Niu, Peihua, Liu, Xu, Zhao, Li, Dong, Erdan, Song, Chunli. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clinical Infectious Diseases*; 2020.
722. Colson, Philippe. Chloroquine for the 2019 novel coronavirus SARS-CoV-2. *International Journal of Antimicrobial Agents*; April 2, 2020.
723. Colson, Philippe, Rolain, Jean-Marc, Lagier, Jean-Christophe, Brouqui, Philippe, Raoult, Didier. Chloroquine and hydroxychloroquine as available weapons to fight COVID-19. *Int J Antimicrob Agents*; 2020.
724. Wang, Manli, Cao, Ruiyuan, Zhang, Leike, Yang, Xinglou, Liu, Jia, Xu, Mingyue, Shi, Zhengli, Hu, Zhihong, Zhong, Wu, Xiao, Gengfu. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell research*; 2020.
725. Vincent, Martin J, Bergeron, Eric, Benjannet, Suzanne, Erickson, Bobbie R, Rollin, Pierre E, Ksiazek, Thomas G, Seidah, Nabil G, Nichol, Stuart T. Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology journal*; 2005.
726. Woodyatt, Amy, Hollingsworth, Julia, Westcott, Ben, Renton, Adam, Wagner, Meg, Hayes, Mike. March 30 coronavirus news. 2020.
727. Savarino, A., Boelaert, J. R., Cassone, A., Majori, G., Cauda, R. Effects of chloroquine on viral infections: an old drug against today's diseases?. *Lancet Infect Dis*; Nov 2003.
728. Rolain, Jean-Marc, Colson, Philippe, Raoult, Didier. Recycling of chloroquine and its hydroxyl analogue to face bacterial, fungal and viral infections in the 21st century. *International journal of antimicrobial agents*; 2007.
729. Lammers, A. J. J., Brohet, R. M., Theunissen, R. E. P., Koster, C., Rood, R., Verhagen, D. W. M., Brinkman, K., Hassing, R. J., Dofferhoff, A., el Moussaoui, R., Hermanides, G., Ellerbroek, J., Bokhizzou, N., Visser, H., van den Berge, M., Bax, H., Postma, D. F., Groeneveld, P. H. P. Early hydroxychloroquine but not chloroquine use reduces ICU admission in COVID-19 patients. *International Journal of Infectious Diseases*; 2020/12/01/.

730. Watanabe, Márcio. Efficacy of Hydroxychloroquine as Prophylaxis for Covid-19. *arXiv preprint arXiv:2007.09477*; 2020.
731. Sire, Alessandro DE, Andrenelli, Elisa, Negrini, Francesco, Lazzarini, Stefano G, Patrini, Michele, Ceravolo, Maria G. Rehabilitation and COVID-19: the Cochrane Rehabilitation 2020 rapid living systematic review. Update as of August 31st, 2020. *Eur J Phys Rehabil Med*; Dec 2020.
732. Blair, Paul W, Brown, Diane M, Jang, Minyoung, Antar, Annukka A R, Keruly, Jeanne C, Bachu, Vismaya S, Townsend, Jennifer L, Tornheim, Jeffrey A, Keller, Sara C, Sauer, Lauren, Thomas, David L, Manabe, Yukari C, Team, Ambulatory, COVID, Study. The Clinical Course of COVID-19 in the Outpatient Setting: A Prospective Cohort Study. *Open Forum Infectious Diseases*; 2021.
733. Moreno-Pérez, Oscar, Merino, Esperanza, Leon-Ramirez, Jose-Manuel, Andres, Mariano, Ramos, Jose Manuel, Arenas-Jiménez, Juan, Asensio, Santos, Sanchez, Rosa, Ruiz-Torregrosa, Paloma, Galan, Irene. Post-acute COVID-19 Syndrome. Incidence and risk factors: a Mediterranean cohort study. *Journal of Infection*; 2021.
734. Cirulli, Elizabeth T., Schiabor Barrett, Kelly M., Riffle, Stephen, Bolze, Alexandre, Neveux, Iva, Dabe, Shaun, Grzymiski, Joseph J., Lu, James T., Washington, Nicole L. Long-term COVID-19 symptoms in a large unselected population. *medRxiv*; 2020.
735. Cellai, Michele, O'Keefe, James B. Characterization of prolonged COVID-19 symptoms in an outpatient telemedicine clinic. *Open forum infectious diseases*; 2020.
736. Sudre, Carole H., Murray, Benjamin, Varsavsky, Thomas, Graham, Mark S., Penfold, Rose S., Bowyer, Ruth C., Pujol, Joan Capdevila, Klaser, Kerstin, Antonelli, Michela, Canas, Liane S., Molteni, Erika, Modat, Marc, Cardoso, M. Jorge, May, Anna, Ganesh, Sajaysurya, Davies, Richard, Nguyen, Long H, Drew, David A., Astley, Christina M., Joshi, Amit D., Merino, Jordi, Tsereteli, Neli, Fall, Tove, Gomez, Maria F., Duncan, Emma L., Menni, Cristina, Williams, Frances M.K., Franks, Paul W., Chan, Andrew T., Wolf, Jonathan, Ourselin, Sebastien, Spector, Tim, Steves, Claire J. Attributes and predictors of Long-COVID: analysis of COVID cases and their symptoms collected by the Covid Symptoms Study App. *medRxiv*; 2020.
737. OfNS. The prevalence of long COVID symptoms and COVID-19 complications. 2020.
738. Zapatero, Diego Castanares, Hanquet, Germaine, Van Den Heede, Koen. Epidemiology of Long COVID: A Pragmatic Review of the Literature. 2021.
739. Goërtz, Yvonne M.J., Van Herck, Maarten, Delbressine, Jeannet M., Vaes, Anouk W., Meys, Roy, Machado, Felipe V.C., Houben-Wilke, Sarah, Burtin, Chris, Posthuma, Rein, Franssen, Frits M.E., van Loon, Nicole, Hajian, Bitá, Spies, Yvonne, Vijlbrief, Herman, van 't Hul, Alex J., Janssen, Daisy J.A., Spruit, Martijn A. Persistent symptoms 3 months after a SARS-CoV-2 infection: the post-COVID-19 syndrome?. *ERJ Open Research*; 2020.
740. Davis, Hannah E., Assaf, Gina S., McCorkell, Lisa, Wei, Hannah, Low, Ryan J., Re'em, Yochai, Redfield, Signe, Austin, Jared P., Akrami, Athena. Characterizing Long COVID in an International Cohort: 7 Months of Symptoms and Their Impact. *medRxiv*; 2020.
741. Vaes, Anouk W, Machado, Felipe VC, Meys, Roy, Delbressine, Jeannet M, Goertz, Yvonne MJ, Van Herck, Maarten, Houben-Wilke, Sarah, Franssen, Frits ME, Vijlbrief, Herman, Spies, Yvonne. Care dependency in non-hospitalized patients with COVID-19. *Journal of Clinical Medicine*; 2020.
742. del Rio, Carlos, Collins, Lauren F., Malani, Preeti. Long-term Health Consequences of COVID-19. *JAMA*; 2020.
743. Mazza, Mario Gennaro, De Lorenzo, Rebecca, Conte, Caterina, Poletti, Sara, Vai, Benedetta, Bollettini, Irene, Melloni, Elisa Maria Teresa, Furlan, Roberto, Ciceri, Fabio, Rovere-Querini, Patrizia. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain, behavior, and immunity*; 2020.
744. Taquet, Maxime, Luciano, Sierra, Geddes, John R., Harrison, Paul J. Bidirectional associations between COVID-19 and psychiatric disorder: retrospective cohort studies of 62–354 COVID-19 cases in the USA. *The Lancet Psychiatry*; 2021.
745. Bellan, Mattia, Soddu, Daniele, Balbo, Piero Emilio, Baricich, Alessio, Zeppegno, Patrizia, Avanzi, Gian Carlo, Baldon, Giulia, Bartolomei, Giuseppe, Battaglia, Marco, Battistini, Sofia. Respiratory and Psychophysical Sequelae Among Patients With COVID-19 Four Months After Hospital Discharge. *JAMA Network Open*; 2021.
746. Iannaccone, Sandro, Castellazzi, Paola, Tettamanti, Andrea, Houdayer, Elise, Brugliera, Luigia, de Blasio, Francesco, Cimino, Paolo, Ripa, Marco, Meloni, Carlo, Alemanno, Federica, Scarpellini, Paolo. Role of Rehabilitation Department for Adult Individuals With COVID-19: The Experience of the San Raffaele Hospital of Milan. *Archives of physical medicine and rehabilitation*; 2020.
747. Salawu, Abayomi, Green, Angela, Crooks, Michael G, Brixey, Nina, Ross, Denise H, Sivan, Manoj. A proposal for multidisciplinary tele-rehabilitation in the assessment and rehabilitation of COVID-19 survivors. *International journal of environmental research and public health*; 2020.

748. Aiyegbusi, O., Hughes, S., Turner, G., Rivera, S., McMullan, C., Chandan, J., Haroon, S., Price, G., Davies, E., Nirantharakumar, K., Sapey, E., Calvert, M. Symptoms, complications and management of long COVID: a review. *J R Soc Med*; Sep 2021.

749. Negrini, Stefano, Donzelli, Sabrina, Negrini, Alberto, Negrini, Alessandra, Romano, Michele, Zaina, Fabio. Feasibility and acceptability of telemedicine to substitute outpatient rehabilitation services in the COVID-19 emergency in Italy: an observational everyday clinical-life study. *Archives of physical medicine and rehabilitation*; 2020.